

**INTERANNUAL VARIABILITY
OF
CHLOROPHYLL CONCENTRATION
IN THE
EASTERN ARABIAN SEA**

A Summer Internship Report

BY
MS. Satlaj Karanje

Under the guidance of

Dr. S. Prasanna Kumar

Scientist, NIO, Goa

**Department of Geography
University of Pune
June 2008**

DECLARATION

I hereby declare that the work incorporated in this dissertation (as a part of the M.Sc course) is original and carried out at National Institute of Oceanography, Dona Paula, Goa under the guidance of Dr. S. Prasanna Kumar (Senior Scientist, Physical Oceanography Department) and it has not been submitted in part or in full any degree, diploma of any other University.

DATE: 6th June 2008

Satlaj Karanje

Place: Panaji-Goa

National Institute of Oceanography
Dona Paula, Goa 403 004, India



Tel: 91-832-2450300
Fax: 91-832-2450608, 2450602

Dr. S. Prasanna Kumar
Scientist F

CERTIFICATE

This is to certify that the dissertation work entitled, “*INTER ANNUAL VARIABILITY OF CHLOROPHYLL CONCENTRATION IN THE EASTERN ARABIAN SEA*” is a bonafide and authentic work carried out by Miss Satlaj Karanje as a part of the summer training programme held at National Institute of Oceanography, Goa, in May-June, 2008 and is according to the results obtained bY her under my guidance and supervision.

S. Prasanna Kumar

ACKNOWLEDGEMENT

I am thankful to Director, N.I.O and HRDG for permitting me to complete my dissertation work at National Institute of Oceanography, Goa.

It gives me immense pleasure to express deep gratitude to my guide Dr. S. Prasanna Kumar, Scientist, Physical Oceanography Department, N.I.O; Goa for his constant and untiring guidance, inspiration, support and encouragement during the course of this dissertation work. I am also grateful to him for motivating me towards new ideas and I take this opportunity to express my indebtedness and respect to him.

Above all, I owe it all to the Almighty for being with me through all the times.

Satlaj Karanje

CONTENTS

1. INTRODUCTION

Chlorophyll and Phytoplankton in ocean and their Importance.

2. DATA DISCRIPTION

Website and system parameters

- Chlorophyll a concentration
- Diffused Attenuation Coefficient

3. RESULTS

- Chlorophyll concentration according to Time Variation
- Climatological Chlorophyll concentration
- Annual Chlorophyll concentration

1. INTRODUCTION

Introduction

The sea can be in many different colours. What colours you see depend not only on the weather and light conditions, but also on what the water contains. This is why ocean colour measurements have become an important tool for studying plant life in the ocean.

Calculating chlorophyll from measurements of watercolour is one of the successes of optical oceanography over the last 30 years or so. As a result we can now use satellite measurements to tell us how well plants in the ocean are growing.

Like plants on land, plants growing in water contain chlorophyll, a molecule that allows the plant to trap the energy in sunlight for photosynthesis. Chlorophyll absorbs blue and red light, so it looks green to us. That is why grass and leaves are green.

The most important plants in the sea are phytoplankton - microscopic plants that float suspended in the water. Where there is a lot of phytoplankton, most of the blue light is absorbed, so the water looks green. Where there are none (or very few), the blue light is not absorbed, so the water looks blue.

The balance between blue and green (the blue-green ratio) to calculate how much chlorophyll in the water contains. This allows them to create worldwide maps of chlorophyll from satellite images of ocean colour

Bright green usually tells you there are millions of microscopic plants in the water, but a very similar green can be seen where rivers wash soil into the sea. Optical oceanographers around the world are hard at work trying to tell the two apart.

In the open ocean away from land, particles that scatter light and colour the water green, are almost certainly tiny plants. In this type of water we can now calculate chlorophyll from ocean colour measurement and be confident about the results.

Near land it is not always so simple. Rivers often bring mud and the remains of dead plants into the sea. This mixture has a yellow-brown colour. Where it mixes with seawater the ocean is coloured green, a colour, which is easily mistaken for the green of chlorophyll.

Chlorophyll a concentrations change with the factors that affect phytoplankton growth. Some of those factors are:

- Amount of sunlight
- Nutrient concentrations (nitrate and phosphate)
- Amount of mixing (stratification)
- Water temperature
- Water quality

Phytoplankton

Phytoplankton is microscopic plants that live in the ocean. There are many species of phytoplankton, each of which has a characteristic shape. Collectively, phytoplankton grows abundantly in oceans around the world and is the foundation of the marine food chain. Small fish, and some species of

whales, eat them as food. Larger fish then eat the smaller fish. Humans catch and eat many of these larger fish. Since phytoplankton depends upon certain conditions for growth, they are a good indicator of change in their environment. For these reasons, and because they also exert a global-scale influence on climate, phytoplankton are of primary interest to oceanographers and Earth scientists around the world.

Like their land-based relatives, phytoplankton requires sunlight, water, and nutrients for growth. Because sunlight is most abundant at and near the sea surface, phytoplankton remains at or near the surface. Also like terrestrial plants, phytoplankton contains the pigment chlorophyll, which gives them their greenish color. Chlorophyll is used by plants for photosynthesis, in which sunlight is used as an energy source to fuse water molecules and carbon dioxide into carbohydrates—plant food. Phytoplankton (and land plants) use carbohydrates as "building blocks" to grow fish and humans consume plants to get these same carbohydrates.

Remote Sensing

Some Earth observing satellites measure the characteristics of light, or radiance, coming from the Earth's surface. To learn about what is in the water using observations from space, we must first know what influences the color of water. Samples of ocean water are taken and their concentrations of phytoplankton and their chlorophyll are analyzed; these concentrations will then be correlated with the measured radiances. As these measurements are made, researchers hope to find consistent relationships between the radiances and the surface variables that are being measured, which will allow them to construct an algorithm. The algorithm will calculate a specific variable, such as chlorophyll concentration, based solely on the radiance data. Satellite data will then be used in these algorithms to calculate the geophysical variables over large areas of the Earth.

The goal of ocean color remote sensing algorithms is to distinguish different types of water, and the constituents that determine a particular color. Ideally, a useful algorithm would calculate the concentration of suspended particulates in the muddy water, and the concentration of chlorophyll in both turbid and clear water.

A part of the ocean with only one kind of phytoplankton will have a fairly uniform color. In these cases, a fairly simple relationship exists between the color that the satellite observes and the density of phytoplankton. Ratios of light intensity detected at various wavelengths of the visible and infrared spectrum have been used in algorithms to calculate vegetation density or chlorophyll concentration. However, if water contains different species of phytoplankton as well as sediments, it becomes much more difficult to find simple relationships between optical properties and geophysical characteristic.

To further complicate the matter, instruments in space tend to change over time, and they usually can't be re-calibrated. Unfortunately, algorithms such as those described above rely on very accurate measurements of radiance. Thus mission operations ensure that a very high level of precision knows calibration of the instrument. Several different ways of calibrating these sensors while the satellite is in space have been devised.

Instruments

The primary instruments NASA uses to investigate ocean color are SeaWiFS and MODIS:

- SeaWiFS

The purpose of the Sea-viewing Wide Field-of-view Sensor (SeaWiFS) Project is to provide useful data on ocean color to the Earth science community. SeaWiFS flies on the OrbView-2 satellite, but beyond the

instrument itself, the SeaWiFS Project has developed and operates a research data system that processes, calibrates, validates, archives and distributes data received from an Earth-orbiting ocean color sensor.

Chlorophyll

With the SeaWiFS instrument, NASA has gathered the first record of photosynthetic productivity in the oceans. The process begins with a measurement of surface chlorophyll concentration. Chlorophyll is the material that allows plant cells to convert sunlight into energy, thus enabling them to grow. It's a green substance, and thus a good indicator of overall plant health: robust forests, lush lawns and vibrant phytoplankton blooms appear green. By measuring chlorophyll concentration, scientists can determine the health and growth of plants in a given area. By extension, healthy color signatures indicate the successful use of carbon, the fundamental building block for life.

Measuring chlorophyll can identify areas rich in nutrients and monitor such processes such as upwelling. As the surface water moves offshore, cold, nutrient-rich water upwells from below to replace it. This upwelling provides nutrients needed for the growth of marine phytoplankton, which, along with larger seaweeds, in turn nourishes the incredible diversity of creatures.

Sensors such as SeaWiFS can "see" the effects of this upwelling-related productivity because the chlorophyll-bearing phytoplankton reflect predominantly green light back into space as opposed to the water itself which reflects predominantly blue wavelengths back to space.

Optical Remote Sensing

Optical remote sensing makes use of visible, near infrared and short wave infrared sensors to form images of the earth's surface by detecting the solar radiation reflected from targets on the ground. Different materials reflect and absorb differently at different wavelengths. Thus, their spectral reflectance signatures in the remotely sensed images can differentiate the targets. There are four types of optical imaging systems commonly in use:

Panchromatic imaging system: The sensor is a single channel detector sensitive to radiation within a broad wavelength range. If the wavelength range coincides with the visible range, then the resulting image resembles a "black-and-white" photograph taken from space. The physical quantity being measured is the apparent brightness of the targets. The spectral information or "colour" of the targets is lost. An example of a panchromatic imaging system is:

- SPOT HRV-PAN

Multispectral imaging system: The sensor is a multichannel detector. Each channel is sensitive to radiation within a narrow wavelength band. The resulting image is a multilayer image, which contains both the brightness and spectral (colour) information of the targets being observed. Examples of multispectral systems are:

- LANDSAT MSS
- LANDSAT TM
- SPOT HRV-XS

Superspectral Imaging Systems: A superspectral imaging sensor has many more spectral channels (typically >10) than a multispectral sensor. The bands have narrower bandwidths, enabling the finer spectral characteristics of the targets to be captured by the sensor. Examples of superspectral systems are:

- MODIS
- MERIS

Hyperspectral Imaging Systems: A hyperspectral imaging system is also known as an "imaging spectrometer". It acquires images in about a hundred or more contiguous spectral bands. The precise spectral information contained in a hyperspectral image enables better characterisation and identification of targets. Hyperspectral images have potential applications in such fields as precision agriculture (e.g. monitoring the types, health, moisture status and maturity of crops), coastal management (e.g. monitoring of phytoplankton, pollution, bathymetry changes). An example of a hyperspectral system is:

- Hyperion on EO1 satellite

2. DATA DESCRIPTION

Ocean Biology Processing Group (OBPG) SeaWiFS global 9-km resolution data products.

- Website Name:
<http://reason.gsfc.nasa.gov/OPS/Giovanni/ocean.seawifs.shtml>
- Name Of Sensor: SeaWiFS (Sea_viewing Wide Field_of_view Sensor)
- Data Analysed: Chlorophyll-a concentration
- Selected Parameters: Diffuse_attenuation_coefficient_at_490_nm
- Selected Area: Latitude=[5.0N, 16.0N]
Longitude=[70.0E, 80.0E]
- Selected Time Period: January, 1998 to December, 2007
- Output Data Resolution: 1.0 Degree
- Undefined/Missing Value: -99999

SeaWiFS

SeaWiFS stands for Sea--viewing Wide Field-of-view Sensor. It is the only scientific instrument on GeoEye's OrbView-2 (AKA SeaStar) satellite, and was a follow-on experiment to the Coastal Zone Color Scanner on Nimbus 7. Launched August 1, 1997 on an Orbital Sciences Pegasus small air launched rocket, the

instrument began scientific operations on 18 September 1997. The sensor resolution is 1.1 km (LAC), 4.5 km (GAC). The sensor records information in the following optical bands:

Instrument Bands

Band = Wavelength

1 = 402-422 nm

2 = 433-453 nm

3 = 480-500 nm

4 = 500-520 nm

5 = 545-565 nm

6 = 660-680 nm

7 = 745-785 nm

8 = 845-885 nm

Mission Characteristics

Orbit Type: Sun Synchronous at 705 km

Equator Crossing: Noon +20 min, descending

Orbital Period: 99 minutes

Swath Width: 2,801 km LAC/HRPT (58.3 degrees)

Swath Width: 1,502 km GAC (45 degrees)

Spatial Resolution: 1.1 km LAC, 4.5 km GAC

Real-Time Data Rate: 665 kbps

Revisit Time: 1 day

Digitization: 10 bits

The instrument has been specifically designed to monitor ocean characteristics such as chlorophyll-a concentration and water clarity. The instrument is able to tilt up to 20 degrees to avoid sunlight from the sea surface. This feature is important at equatorial latitudes where glint from sunlight often obscures watercolour.

Diffuse Attenuation Coefficient

Natural waters have what are often referred to as inherent and apparent optical properties. Inherent optical properties (IOP) are a function of the water and optically active substances in it and are not influenced by the geometric structure of the light fields.

Apparent optical properties (AOP) are derived from measurements of natural light fields in a water body. They depend on the geometry of the light fields and are related to absorption and scattering. The most common of these properties is the diffuse attenuation coefficient for downwelling irradiance (K_d). Irradiance at a given depth (E_z) is a function of the irradiance at the surface (E_0), the diffuse attenuation coefficient, and the depth interval (Z) according to the following relationship, where e is the base of the natural logarithms:

$$E_z = E_0 e^{-k_d Z}$$

From this we can see that when light penetrates water its intensity decreases exponentially with increasing depth.

The diffuse attenuation coefficient can thus be estimated by taking measurements at different depths, and using the above formula to compute K_d . The units for K_d are m^{-1} .

$$K_d = \frac{\ln\left(\frac{E_0}{E_z}\right)}{Z}$$

From this we can see that when light penetrates water its intensity decreases exponentially with increasing depth.


The diffuse attenuation coefficient can thus be estimated by taking measurements at different depths, and using the above formula to compute K_d . The units for K_d are m^{-1}

Using these diffuse attenuation coefficients (obtained from spectral irradiance measurements at different depths in the water), and using the fact that the attenuation in water decreases exponentially with respect to depth (given by the equation discussed earlier), we can plot the spectral curve for the underwater light field at various depths.

3. RESULTS

Chlorophyll Concentration

Monthly mean chlorophyll in January during 1998 to 2007




Period		Chlorophyll Concentration (mg/m**3)			
Year	Month	Min	Max	Mean	SD
1998	January	0.115919	0.506658	0.1962968	0.075480451
1999	January	0.152299	0.583982	0.28768289	0.090701973
2000	January	0.147365	0.574984	0.283160917	0.09709633
2001	January	0.135664	0.581019	0.260069	0.089526
2002	January	0.130446	0.567490	0.259251	0.098126
2003	January	0.128385	0.586411	0.234599419	0.079026732
2004	January	0.127870	0.589086	0.259599	0.088398
2005	January	0.121626	0.555613	0.219225	0.069929
2006	January	0.156893	0.528056	0.304912	0.085191
2007	January	0.130080	0.583513	0.247527	0.086869

Monthly mean chlorophyll in February during 1998 to 2007


Period		Chlorophyll Concentration (mg/m**3)			
Year	Month	Min	Min	Mean	SD
1998	February	0.099486	0.468495	0.193176	0.074271
1999	February	0.168181	0.590033	0.267984	0.076582
2000	February	0.132417	0.598545	0.202186	0.077033
2001	February	0.149711	0.55755	0.238799	0.089441
2002	February	0.134327	0.487888	0.221794	0.074436
2003	February	0.142115	0.503548	0.212643	0.064959
2004	February	0.142343	0.480564	0.21437	0.05619
2005	February	0.121495	0.536176	0.200226	0.072346
2006	February	0.151242	0.561144	0.280731	0.092934
2007	February	0.109215	0.483597	0.202356	0.061072

Chlorophyll Concentration (February) 1998-2007




Monthly mean chlorophyll in March during 1998 to 2007

Chlorophyll Concentration (March) 1998-2007



Period		Chlorophyll Concentration (mg/m ^{**3})			
Year	Month	Min	Min	Mean	SD
1998	March	0.117906	0.585391	0.227546471	0.086823965
1999	March	0.145678	0.546686	0.248522	0.092377
2000	March	0.124375	0.586175	0.201137	0.081929
2001	March	0.121886	0.553682	0.19897	0.070214
2002	March	0.132701	0.351654	0.192731	0.046035
2003	March	0.130464	0.490705	0.195142	0.064701
2004	March	0.136374	0.529755	0.20434358	0.061691087
2005	March	0.128505	0.568783	0.191638	0.072331
2006	March	0.123463	0.444425	0.182925	0.048775

2007	March	0.117638	0.573293	0.19459549	0.067791
Monthly mean chlorophyll in April during 1998 to 2007					




Period		Chlorophyll Concentration (mg/m**3)			
Year	Month	Min	Min	Mean	SD
1998	April	0.107604	0.5346	0.215881941	0.080923024

1999	April	0.120475	0.597984	0.232627	0.099572
2000	April	0.090878	0.541587	0.19655	0.072747
2001	April	0.075458	0.570079	0.192323	0.085772
2002	April	0.085485	0.542821	0.178476	0.085874
2003	April	0.095608	0.539849	0.182737	0.085829
2004	April	0.099326	0.570689	0.18707	0.093668
2005	April	0.171202	0.543039	0.198874	0.078795
2006	April	0.117151	0.451380	0.191284	0.068602
2007	April	0.068534	0.549694	0.18369	0.079331


Monthly mean chlorophyll in May during 1998 to 2007

Chlorophyll Concentration (May) 1998-2007



Period		Chlorophyll Concentration (mg/m**3)			
Year	Month	Min	Min	Mean	SD
1998	May	0.049635	2.179220	0.28035445	0.32225755
1999	May	0.105960	5.217650	0.584843	0.899697
2000	May	0.108675	5.894390	0.663268	1.018087
2001	May	0.098163	4.151780	0.426751	0.738455
2002	May	0.088288	5.467930	0.455939	0.940941
2003	May	0.072984	6.954060	0.488321	1.064592
2004	May	0.127921	7.644380	0.636307	1.278236


2005	May	0.096611	3.353550	0.319542	0.502388
2006	May	0.074505	9.699940	0.455594	1.20824
2007	May	0.078687	3.264950	0.338383	0.569012



Period		Chlorophyll Concentration (mg/m^{**3})			
Year	Month	Min	Max	Avg	Sd
1998	June	0.116510	9.960480	0.702551	1.554645


1999	June	0.101980	7.909830	0.663969	1.343921
2000	June	0.124359	5.983430	0.251472	0.124357
2001	June	0.124926	4.257020	0.641499	0.949435
2002	June	0.124475	9.447480	0.713138	1.45223
2003	June	0.079407	5.504660	0.513164	0.931152
2004	June	0.145367	6.338600	0.714007	1.059723
2005	June	0.106529	4.141230	0.715978	0.932412
2006	June	0.103493	8.941730	0.568587	1.249939
2007	June	0.086221	7.333640	0.430176	0.954468

Chlorophyll Concentration (Jully) 1998-2007



Period		Chlorophyll Concentration (mg/m^3)			
Year	Month	Min	Max	Avg	Sd
1998	July	0.148852	8.927370	1.005028788	1.493174
1999	July	0.109993	9.329340	0.910385	1.536307
2000	July	0.150012	7.327600	0.959887	1.309534
2001	July	0.094109	6.102340	0.846229	1.278345
2002	July	0.114884	7.809880	0.931339	1.631558
2003	July	0.124559	7.270100	0.731167	1.295615
2004	July	0.140112	7.529610	1.092355	1.471404
2005	July	0.122328	8.365060	1.0813	1.568241
2006	July	0.088359	5.913350	0.821535	1.2070276
2007	July	0.118143	9.787330	1.147339	1.834658


Chlorophyll Concentration (August) 1998-2007



Period		Chlorophyll Concentration (mg/m**3)			
Year	Month	Min	Max	Avg	Sd
1998	August	0.139887	5.772570	0.936042	1.07187
1999	August	0.100651	8.981310	0.941887	1.437711
2000	August	0.134453	6.371980	0.936801	1.195823
2001	August	0.140075	9.833300	1.230228	2.068522
2002	August	0.136594	9.810940	1.151681	1.875342
2003	August	0.143606	6.100700	0.832276	1.528585
2004	August	0.133824	8.389730	1.206923	1.939257
2005	August	0.113220	0.578148	0.285445	0.139158

2006	August	0.089490	8.133170	0.866363	1.636326
2007	August	0.092263	6.878150	0.797671	1.469776


Chlorophyll Concentration (September) 1998-2007



Period		Chlorophyll Concentration (mg/m**3)			
Year	Month	Min	Max	Avg	Sd
1998	September	0.102363	5.440920	0.891366	1.218617
1999	September	0.083439	6.286410	0.8543951	1.286841


2000	September	0.125866	6.773980	1.002357	1.540545
2001	September	0.129784	6.266600	0.803025	1.102662
2002	September	0.111535	7.296420	0.853577	1.398779
2003	September	0.127340	5.307120	0.710019	1.022772
2004	September	0.139215	6.154840	0.889526	1.416467
2005	September	0.132310	6.508190	0.828887	1.130858
2006	September	0.104118	5.973020	0.591796	0.840731
2007	September	0.005900	4.477980	0.560096	0.924833

Chlorophyll Concentration (October) 1998-2007




Period		Chlorophyll Concentration (mg/m**3)			
Year	Month	Min	Max	Avg	Sd
1998	October	0.138657	4.282740	0.695143	0.807138
1999	October	0.154496	7.316970	0.716535	1.050352
2000	October	0.135048	7.576120	0.489434	0.884221
2001	October	0.177218	8.702010	0.798498	1.13833
2002	October	0.107792	5.785720	0.512367	0.811163
2003	October	0.143107	3.901210	0.596692	0.759909
2004	October	0.106060	3.776350	0.508421	0.663902
2005	October	0.142916	4.186940	0.570425	0.787981
2006	October	0.096276	4.796300	0.441425	0.766952
2007	October	0.106259	4.814960	0.428883	0.666792

Chlorophyll Concentration (November) 1998-2007



Period		Chlorophyll Concentration (mg/m**3)			
Year	Month	Min	Max	Avg	Sd
1998	November	0.137972	0.598946	0.301951	0.122754
1999	November	0.117512	0.595986	0.306179	0.130291
2000	November	0.128041	0.572296	0.250894	0.093196
2001	November	0.137898	0.570460	0.28703	0.113037
2002	November	0.105268	0.546100	0.227066	0.098782
2003	November	0.131904	0.593916	0.27	0.106021
2004	November	0.117371	0.567034	0.234136	0.099109

2005	November	0.162406	0.518946	0.283577	0.09271
2006	November	0.065441	0.501530	0.171633	0.08674
2007	November	0.134940	0.581641	0.229856	0.082681




Period		Chlorophyll Concentration (mg/m^{**3})			
Year	Month	Min	Max	Avg	Sd
1998	December	0.124142	0.569240	0.294674	0.09568

1999	December	0.121300	0.597664	0.269298	0.113527
2000	December	0.140601	0.598305	0.257348	0.095433
2001	December	0.159178	0.595030	0.274006	0.089492
2002	December	0.122557	0.589638	0.217753	0.076482
2003	December	0.133449	0.583279	0.25119	0.085332
2004	December	0.124093	0.530181	0.225793	0.0920044
2005	December	0.183765	0.581953	0.289334	0.074082
2006	December	0.103493	0.517143	0.203881	0.079621
2007	December	0.137198	0.460043	0.260375	0.069918


Seasonal cycle of Climatological Monthly mean Chlorophyll

Climatic Chlorophyll Concentration



Highest Climatic Chlorophyll Concentration July=0.95266 mg/m**3

Month_Name	Month_No.	Climatic Chlorophyll Concentration (mg/m**3)
January	1	0.255232303
February	2	0.2234265
March	3	0.203755054
April	4	0.195951294
May	5	0.464930245
June	6	0.5914541
July	7	0.952656479
August	8	0.9185317
September	9	0.79850441
October	10	0.5757823
November	11	0.2562322
December	12	0.2543652




Period			Chlorophyll Concentration (mg/m**3)			
Obs.	Year	Month	Min	Max	Avg	Sd
1	1998	January	0.115919	0.506658	0.1962968	0.075480451
2	1999	January	0.152299	0.583982	0.28768289	0.090701973
3	2000	January	0.147365	0.574984	0.283160917	0.09709633
4	2001	January	0.135664	0.581019	0.260069	0.089526
5	2002	January	0.130446	0.567490	0.259251	0.098126
6	2003	January	0.128385	0.586411	0.234599419	0.079026732
7	2004	January	0.127870	0.589086	0.259599	0.088398
8	2005	January	0.121626	0.555613	0.219225	0.069929
9	2006	January	0.156893	0.528056	0.304912	0.085191
10	2007	January	0.130080	0.583513	0.247527	0.086869
11	1998	February	0.099486	0.468495	0.193176	0.074271
12	1999	February	0.168181	0.590033	0.267984	0.076582
13	2000	February	0.132417	0.598545	0.202186	0.077033
14	2001	February	0.149711	0.55755	0.238799	0.089441
15	2002	February	0.134327	0.487888	0.221794	0.074436
16	2003	February	0.142115	0.503548	0.212643	0.064959
17	2004	February	0.142343	0.480564	0.21437	0.05619
18	2005	February	0.121495	0.536176	0.200226	0.072346
19	2006	February	0.151242	0.561144	0.280731	0.092934
20	2007	February	0.109215	0.483597	0.202356	0.061072
21	1998	March	0.117906	0.585391	0.227546471	0.086823965
22	1999	March	0.145678	0.546686	0.248522	0.092377
23	2000	March	0.124375	0.586175	0.201137	0.081929
24	2001	March	0.121886	0.553682	0.19897	0.070214
25	2002	March	0.132701	0.351654	0.192731	0.046035
26	2003	March	0.130464	0.490705	0.195142	0.064701
27	2004	March	0.136374	0.529755	0.20434358	0.061691087
28	2005	March	0.128505	0.568783	0.191638	0.072331
29	2006	March	0.123463	0.444425	0.182925	0.048775
30	2007	March	0.117638	0.573293	0.19459549	0.067791
31	1998	April	0.107604	0.5346	0.215881941	0.080923024
32	1999	April	0.120475	0.597984	0.232627	0.099572
33	2000	April	0.090878	0.541587	0.19655	0.072747
34	2001	April	0.075458	0.570079	0.192323	0.085772
35	2002	April	0.085485	0.542821	0.178476	0.085874
36	2003	April	0.095608	0.539849	0.182737	0.085829
37	2004	April	0.099326	0.570689	0.18707	0.093668

38	2005	April	0.171202	0.543039	0.198874	0.078795
39	2006	April	0.117151	0.451380	0.191284	0.068602
40	2007	April	0.068534	0.549694	0.18369	0.079331
41	1998	May	0.049635	2.179220	0.28035445	0.32225755
42	1999	May	0.105960	5.217650	0.584843	0.899697
43	2000	May	0.108675	5.894390	0.663268	1.018087
44	2001	May	0.098163	4.151780	0.426751	0.738455
45	2002	May	0.088288	5.467930	0.455939	0.940941
46	2003	May	0.072984	6.954060	0.488321	1.064592
47	2004	May	0.127921	7.644380	0.636307	1.278236
48	2005	May	0.096611	3.353550	0.319542	0.502388
49	2006	May	0.074505	9.699940	0.455594	1.20824
50	2007	May	0.078687	3.264950	0.338383	0.569012
51	1998	June	0.116510	9.960480	0.702551	1.554645
52	1999	June	0.101980	7.909830	0.663969	1.343921
53	2000	June	0.124359	5.983430	0.251472	0.124357
54	2001	June	0.124926	4.257020	0.641499	0.949435
55	2002	June	0.124475	9.447480	0.713138	1.45223
56	2003	June	0.079407	5.504660	0.513164	0.931152
57	2004	June	0.145367	6.338600	0.714007	1.059723
58	2005	June	0.106529	4.141230	0.715978	0.932412
59	2006	June	0.103493	8.941730	0.568587	1.249939
60	2007	June	0.086221	7.333640	0.430176	0.954468
61	1998	July	0.148852	8.927370	1.005028788	1.493174
62	1999	July	0.109993	9.329340	0.910385	1.536307
63	2000	July	0.150012	7.327600	0.959887	1.309534
64	2001	July	0.094109	6.102340	0.846229	1.278345
65	2002	July	0.114884	7.809880	0.931339	1.631558
66	2003	July	0.124559	7.270100	0.731167	1.295615
67	2004	July	0.140112	7.529610	1.092355	1.471404
68	2005	July	0.122328	8.365060	1.0813	1.568241
69	2006	July	0.088359	5.913350	0.821535	1.2070276
70	2007	July	0.118143	9.787330	1.147339	1.834658
71	1998	August	0.139887	5.772570	0.936042	1.07187
72	1999	August	0.100651	8.981310	0.941887	1.437711
73	2000	August	0.134453	6.371980	0.936801	1.195823
74	2001	August	0.140075	9.833300	1.230228	2.068522
75	2002	August	0.136594	9.810940	1.151681	1.875342
76	2003	August	0.143606	6.100700	0.832276	1.528585
77	2004	August	0.133824	8.389730	1.206923	1.939257


78	2005	August	0.113220	0.578148	0.285445	0.139158
79	2006	August	0.089490	8.133170	0.866363	1.636326
80	2007	August	0.092263	6.878150	0.797671	1.469776
81	1998	September	0.102363	5.440920	0.891366	1.218617
82	1999	September	0.083439	6.286410	0.8543951	1.286841
83	2000	September	0.125866	6.773980	1.002357	1.540545
84	2001	September	0.129784	6.266600	0.803025	1.102662
85	2002	September	0.111535	7.296420	0.853577	1.398779
86	2003	September	0.127340	5.307120	0.710019	1.022772
87	2004	September	0.139215	6.154840	0.889526	1.416467
88	2005	September	0.132310	6.508190	0.828887	1.130858
89	2006	September	0.104118	5.973020	0.591796	0.840731
90	2007	September	0.005900	4.477980	0.560096	0.924833
91	1998	October	0.138657	4.282740	0.695143	0.807138
92	1999	October	0.154496	7.316970	0.716535	1.050352
93	2000	October	0.135048	7.576120	0.489434	0.884221
94	2001	October	0.177218	8.702010	0.798498	1.13833
95	2002	October	0.107792	5.785720	0.512367	0.811163
96	2003	October	0.143107	3.901210	0.596692	0.759909
97	2004	October	0.106060	3.776350	0.508421	0.663902
98	2005	October	0.142916	4.186940	0.570425	0.787981
99	2006	October	0.096276	4.796300	0.441425	0.766952
100	2007	October	0.106259	4.814960	0.428883	0.666792
101	1998	November	0.137972	0.598946	0.301951	0.122754
102	1999	November	0.117512	0.595986	0.306179	0.130291
103	2000	November	0.128041	0.572296	0.250894	0.093196
104	2001	November	0.137898	0.570460	0.28703	0.113037
105	2002	November	0.105268	0.546100	0.227066	0.098782
106	2003	November	0.131904	0.593916	0.27	0.106021
107	2004	November	0.117371	0.567034	0.234136	0.099109
108	2005	November	0.162406	0.518946	0.283577	0.09271
109	2006	November	0.065441	0.501530	0.171633	0.08674
110	2007	November	0.134940	0.581641	0.229856	0.082681
111	1998	December	0.124142	0.569240	0.294674	0.09568
112	1999	December	0.121300	0.597664	0.269298	0.113527
113	2000	December	0.140601	0.598305	0.257348	0.095433
114	2001	December	0.159178	0.595030	0.274006	0.089492
115	2002	December	0.122557	0.589638	0.217753	0.076482
116	2003	December	0.133449	0.583279	0.25119	0.085332
117	2004	December	0.124093	0.530181	0.225793	0.0920044

118	2005	December	0.183765	0.581953	0.289334	0.074082
119	2006	December	0.103493	0.517143	0.203881	0.079621
120	2007	December	0.137198	0.460043	0.260375	0.069918

Chlorophyll Concentration (May-October) 1998-2007




Chlorophyll Concentration (November To April) 1998-2007



Diffuse Attenuation Coefficient

Diffuse attenuation coefficient January




Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	January	0.033516	0.091622	0.045254	0.012144
1999	January	0.036858	0.092091	0.056072	0.013218
2000	January	0.036587	0.093614	0.056118	0.012385
2001	January	0.035728	0.097765	0.054646	0.011911

2002	January	0.032991	0.097995	0.053752	0.013187
2003	January	0.035396	0.096081	0.051245	0.010157
2004	January	0.034989	0.091911	0.052374	0.010647
2005	January	0.033303	0.090585	0.049904	0.011599
2006	January	0.040731	0.098458	0.061052	0.011682
2007	January	0.036191	0.096535	0.054035	0.012019

Diffuse attenuation coefficient February

Diffuse_attenuation_coefficient (February) 1998-2007




Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	February	0.029388	0.082276	0.043869	0.009341
1999	February	0.041445	0.093926	0.05361	0.009937
2000	February	0.034177	0.095620	0.04597	0.012043
2001	February	0.038898	0.084403	0.049741	0.00942
2002	February	0.035574	0.080716	0.048327	0.008088
2003	February	0.037166	0.094000	0.047915	0.008793
2004	February	0.038477	0.088955	0.048668	0.007615
2005	February	0.033111	0.092872	0.045666	0.009929
2006	February	0.039340	0.097842	0.055183	0.010429
2007	February	0.031238	0.096430	0.047611	0.011201

Diffuse attenuation coefficient March

Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	March	0.032518	0.095208	0.049319	0.010552
1999	March	0.036451	0.091847	0.050551	0.010539
2000	March	0.032099	0.088981	0.045725	0.009996
2001	March	0.032752	0.078189	0.044929	0.007379
2002	March	0.035651	0.089753	0.045371	0.008233
2003	March	0.034882	0.089674	0.045617	0.00933
2004	March	0.037922	0.081536	0.046849	0.00699
2005	March	0.035230	0.080764	0.045204	0.007937
2006	March	0.032544	0.092426	0.043798	0.007967
2007	March	0.034171	0.078335	0.045531	0.007104

Diffuse_attenuation_coefficient (March) 1998-2007




Diffuse attenuation coefficient April

Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	April	0.031085	0.094602	0.04959	0.011888


1999	April	0.033595	0.099899	0.051913	0.017088
2000	April	0.024199	0.079724	0.045826	0.008967
2001	April	0.023400	0.080661	0.043827	0.0103
2002	April	0.025334	0.078767	0.041833	0.010591
2003	April	0.029414	0.095021	0.050196	0.011796
2004	April	0.029809	0.080962	0.043807	0.010354
2005	April	0.028401	0.091210	0.047484	0.011487
2006	April	0.030800	0.071339	0.045438	0.008378
2007	April	0.020600	0.081496	0.044347	0.009887

Diffuse_attenuation_coefficient (April) 1998-2007



Diffuse attenuation coefficient May

Diffuse_attenuation_coefficient (May) 1998-2007




Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	May	0.017801	0.169010	0.052268	0.024371
1999	May	0.027401	0.230912	0.069864	0.040482
2000	May	0.030454	0.250588	0.061086	0.044317
2001	May	0.027701	0.238351	0.058079	0.037954
2002	May	0.026795	0.245813	0.054872	0.043752
2003	May	0.023944	0.302476	0.059405	0.052241

2004	May	0.034400	0.335812	0.071372	0.056301
2005	May	0.028068	0.246199	0.053282	0.03367
2006	May	0.020200	0.224211	0.053225	0.037789
2007	May	0.026726	0.211884	0.052904	0.033612


Diffuse attenuation coefficient June

Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	June	0.032024	0.347483	0.073283	0.058932
1999	June	0.028401	0.299904	0.06688	0.053433
2000	June	0.029599	0.276301	0.075714	0.048845
2001	June	0.026400	0.237208	0.073358	0.048438
2002	June	0.033401	0.404036	0.074487	0.061917
2003	June	0.023201	0.353990	0.064059	0.055214
2004	June	0.035767	0.659813	0.084694	0.086544
2005	June	0.032800	0.216510	0.080089	0.047702
2006	June	0.029824	0.255808	0.061957	0.050515
2007	June	0.025725	0.254331	0.054354	0.037401

Diffuse_attenuation_coefficient (June) 1998-2007



Diffuse_attenuation_coefficient (Jully) 1998-2007




Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	July	0.035452	0.344393	0.094979	0.066646
1999	July	0.031550	0.320324	0.085787	0.05766

2000	July	0.036201	0.503557	0.098274	0.086404
2001	July	0.022400	0.276980	0.0858253	0.056168
2002	July	0.031000	0.349604	0.085596	0.070063
2003	July	0.032134	0.269576	0.075076	0.051516
2004	July	0.035334	0.308716	0.09859	0.061283
2005	July	0.034800	0.333807	0.096698	0.094063
2006	July	0.026695	0.575256	0.087208	0.083338
2007	July	0.031826	0.341732	0.098224	0.076971

Diffuse attenuation coefficient August


Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	August	0.034948	0.269549	0.09572	0.051925
1999	August	0.027753	0.376944	0.090916	0.069292
2000	August	0.034325	0.373959	0.096862	0.069013
2001	August	0.034255	0.379991	0.100624	0.076741
2002	August	0.031500	0.429897	0.107434	0.094608
2003	August	0.031000	0.398130	0.083428	0.073436
2004	August	0.035016	0.338793	0.096233	0.075033
2005	August	0.030997	0.334102	0.087898	0.059761
2006	August	0.026473	0.549478	0.084431	0.084244
2007	August	0.026297	0.320450	0.078997	0.063877

Diffuse_attenuation_coefficient (August) 1998-2007



Diffuse attenuation coefficient September


Diffuse_attenuation_coefficient (September) 1998-2007



Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	September	0.027401	0.246006	0.090774	0.055942
1999	September	0.023201	0.267231	0.0846078	0.0567

2000	September	0.033199	0.275002	0.088561	0.054793
2001	September	0.031999	0.415173	0.091391	0.074713
2002	September	0.030921	0.300316	0.081083	0.057475
2003	September	0.034753	0.382686	0.084567	0.06533
2004	September	0.033199	0.360655	0.087799	0.066506
2005	September	0.035599	0.294651	0.08709	0.051599
2006	September	0.029187	0.384947	0.07705	0.054193
2007	September	0.031905	0.350051	0.075948	0.062009

Diffuse_attenuation_coefficient (October) 1998-2007




Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	October	0.032232	0.214017	0.08023	0.041486
1999	October	0.038602	0.285230	0.08302	0.046128
2000	October	0.034780	0.296683	0.065106	0.03891
2001	October	0.041384	0.278616	0.086102	0.045082
2002	October	0.029874	0.197561	0.064936	0.037951
2003	October	0.036717	0.225159	0.074411	0.040354
2004	October	0.029792	0.228589	0.067665	0.043251
2005	October	0.035900	0.222238	0.073638	0.040456
2006	October	0.029135	0.261367	0.060176	0.040109
2007	October	0.028633	0.287988	0.061082	0.040406


Diffuse attenuation coefficient November

Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	November	0.034627	0.094546	0.059527	0.015895
1999	November	0.033313	0.095198	0.060533	0.017432
2000	November	0.032900	0.096222	0.052836	0.012386
2001	November	0.035438	0.097861	0.059072	0.015917
2002	November	0.029001	0.086602	0.04895	0.012214
2003	November	0.033980	0.088684	0.056234	0.012631
2004	November	0.032317	0.080266	0.049061	0.010976
2005	November	0.039998	0.095095	0.059248	0.014017
2006	November	0.022590	0.088034	0.041636	0.013682
2007	November	0.034800	0.095883	0.049971	0.012143

Diffuse_attenuation_coefficient (November) 1998-2007




Diffuse_attenuation_coefficient (December) 1998-2007



Period		Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Year	Month	Min	Max	Avg	Sd
1998	December	0.032021	0.097967	0.059619	0.014045
1999	December	0.032592	0.095049	0.054238	0.014819
2000	December	0.036627	0.091577	0.053139	0.012558
2001	December	0.038975	0.083850	0.055523	0.010375
2002	December	0.033085	0.094695	0.04851	0.010529
2003	December	0.035791	0.093283	0.053997	0.011535
2004	December	0.033880	0.096421	0.049199	0.012203
2005	December	0.044176	0.095704	0.059201	0.010836


2006	December	0.030439	0.076107	0.045716	0.009322
2007	December	0.035195	0.098207	0.054561	0.011384

Diffuse_attenuation_coefficient (January-December)



Highest Diffuse Attenuation Coefficient at 490 nm August=0.09225(1/m)

Month_Name	Month_no.	Diffuse Attenuation Coefficient at 490 nm (1/m)
January	1	0.05345
February	2	0.04866
March	3	0.04629
April	4	0.04643
May	5	0.05864
June	6	0.07089
July	7	0.09063
August	8	0.09225
September	9	0.08489
October	10	0.07164
November	11	0.05371
December	12	0.05337




Period			Diffuse_attenuation_coefficient_at_490_nm (1/m)			
Obs.	Year	Month	Min	Max	Avg	Sd
1	1998	January	0.033516	0.091622	0.045254	0.012144
2	1999	January	0.036858	0.092091	0.056072	0.013218
3	2000	January	0.036587	0.093614	0.056118	0.012385
4	2001	January	0.035728	0.097765	0.054646	0.011911

5	2002	January	0.032991	0.097995	0.053752	0.013187
6	2003	January	0.035396	0.096081	0.051245	0.010157
7	2004	January	0.034989	0.091911	0.052374	0.010647
8	2005	January	0.033303	0.090585	0.049904	0.011599
9	2006	January	0.040731	0.098458	0.061052	0.011682
10	2007	January	0.036191	0.096535	0.054035	0.012019
11	1998	February	0.029388	0.082276	0.043869	0.009341
12	1999	February	0.041445	0.093926	0.05361	0.009937
13	2000	February	0.034177	0.095620	0.04597	0.012043
14	2001	February	0.038898	0.084403	0.049741	0.00942
15	2002	February	0.035574	0.080716	0.048327	0.008088
16	2003	February	0.037166	0.094000	0.047915	0.008793
17	2004	February	0.038477	0.088955	0.048668	0.007615
18	2005	February	0.033111	0.092872	0.045666	0.009929
19	2006	February	0.039340	0.097842	0.055183	0.010429
20	2007	February	0.031238	0.096430	0.047611	0.011201
21	1998	March	0.032518	0.095208	0.049319	0.010552
22	1999	March	0.036451	0.091847	0.050551	0.010539
23	2000	March	0.032099	0.088981	0.045725	0.009996
24	2001	March	0.032752	0.078189	0.044929	0.007379
25	2002	March	0.035651	0.089753	0.045371	0.008233
26	2003	March	0.034882	0.089674	0.045617	0.00933
27	2004	March	0.037922	0.081536	0.046849	0.00699
28	2005	March	0.035230	0.080764	0.045204	0.007937
29	2006	March	0.032544	0.092426	0.043798	0.007967
30	2007	March	0.034171	0.078335	0.045531	0.007104
31	1998	April	0.031085	0.094602	0.04959	0.011888
32	1999	April	0.033595	0.099899	0.051913	0.017088
33	2000	April	0.024199	0.079724	0.045826	0.008967
34	2001	April	0.023400	0.080661	0.043827	0.0103
35	2002	April	0.025334	0.078767	0.041833	0.010591
36	2003	April	0.029414	0.095021	0.050196	0.011796
37	2004	April	0.029809	0.080962	0.043807	0.010354
38	2005	April	0.028401	0.091210	0.047484	0.011487
39	2006	April	0.030800	0.071339	0.045438	0.008378
40	2007	April	0.020600	0.081496	0.044347	0.009887
41	1998	May	0.017801	0.169010	0.052268	0.024371
42	1999	May	0.027401	0.230912	0.069864	0.040482
43	2000	May	0.030454	0.250588	0.061086	0.044317
44	2001	May	0.027701	0.238351	0.058079	0.037954


45	2002	May	0.026795	0.245813	0.054872	0.043752
46	2003	May	0.023944	0.302476	0.059405	0.052241
47	2004	May	0.034400	0.335812	0.071372	0.056301
48	2005	May	0.028068	0.246199	0.053282	0.03367
49	2006	May	0.020200	0.224211	0.053225	0.037789
50	2007	May	0.026726	0.211884	0.052904	0.033612
51	1998	June	0.032024	0.347483	0.073283	0.058932
52	1999	June	0.028401	0.299904	0.06688	0.053433
53	2000	June	0.029599	0.276301	0.075714	0.048845
54	2001	June	0.026400	0.237208	0.073358	0.048438
55	2002	June	0.033401	0.404036	0.074487	0.061917
56	2003	June	0.023201	0.353990	0.064059	0.055214
57	2004	June	0.035767	0.659813	0.084694	0.086544
58	2005	June	0.032800	0.216510	0.080089	0.047702
59	2006	June	0.029824	0.255808	0.061957	0.050515
60	2007	June	0.025725	0.254331	0.054354	0.037401
61	1998	July	0.035452	0.344393	0.094979	0.066646
62	1999	July	0.031550	0.320324	0.085787	0.05766
63	2000	July	0.036201	0.503557	0.098274	0.086404
64	2001	July	0.022400	0.276980	0.0858253	0.056168
65	2002	July	0.031000	0.349604	0.085596	0.070063
66	2003	July	0.032134	0.269576	0.075076	0.051516
67	2004	July	0.035334	0.308716	0.09859	0.061283
68	2005	July	0.034800	0.333807	0.096698	0.094063
69	2006	July	0.026695	0.575256	0.087208	0.083338
70	2007	July	0.031826	0.341732	0.098224	0.076971
71	1998	August	0.034948	0.269549	0.09572	0.051925
72	1999	August	0.027753	0.376944	0.090916	0.069292
73	2000	August	0.034325	0.373959	0.096862	0.069013
74	2001	August	0.034255	0.379991	0.100624	0.076741
75	2002	August	0.031500	0.429897	0.107434	0.094608
76	2003	August	0.031000	0.398130	0.083428	0.073436
77	2004	August	0.035016	0.338793	0.096233	0.075033
78	2005	August	0.030997	0.334102	0.087898	0.059761
79	2006	August	0.026473	0.549478	0.084431	0.084244
80	2007	August	0.026297	0.320450	0.078997	0.063877
81	1998	September	0.027401	0.246006	0.090774	0.055942
82	1999	September	0.023201	0.267231	0.0846078	0.0567
83	2000	September	0.033199	0.275002	0.088561	0.054793
84	2001	September	0.031999	0.415173	0.091391	0.074713

85	2002	September	0.030921	0.300316	0.081083	0.057475
86	2003	September	0.034753	0.382686	0.084567	0.06533
87	2004	September	0.033199	0.360655	0.087799	0.066506
88	2005	September	0.035599	0.294651	0.08709	0.051599
89	2006	September	0.029187	0.384947	0.07705	0.054193
90	2007	September	0.031905	0.350051	0.075948	0.062009
91	1998	October	0.032232	0.214017	0.08023	0.041486
92	1999	October	0.038602	0.285230	0.08302	0.046128
93	2000	October	0.034780	0.296683	0.065106	0.03891
94	2001	October	0.041384	0.278616	0.086102	0.045082
95	2002	October	0.029874	0.197561	0.064936	0.037951
96	2003	October	0.036717	0.225159	0.074411	0.040354
97	2004	October	0.029792	0.228589	0.067665	0.043251
98	2005	October	0.035900	0.222238	0.073638	0.040456
99	2006	October	0.029135	0.261367	0.060176	0.040109
100	2007	October	0.028633	0.287988	0.061082	0.040406
101	1998	November	0.034627	0.094546	0.059527	0.015895
102	1999	November	0.033313	0.095198	0.060533	0.017432
103	2000	November	0.032900	0.096222	0.052836	0.012386
104	2001	November	0.035438	0.097861	0.059072	0.015917
105	2002	November	0.029001	0.086602	0.04895	0.012214
106	2003	November	0.033980	0.088684	0.056234	0.012631
107	2004	November	0.032317	0.080266	0.049061	0.010976
108	2005	November	0.039998	0.095095	0.059248	0.014017
109	2006	November	0.022590	0.088034	0.041636	0.013682
110	2007	November	0.034800	0.095883	0.049971	0.012143
111	1998	December	0.032021	0.097967	0.059619	0.014045
112	1999	December	0.032592	0.095049	0.054238	0.014819
113	2000	December	0.036627	0.091577	0.053139	0.012558
114	2001	December	0.038975	0.083850	0.055523	0.010375
115	2002	December	0.033085	0.094695	0.04851	0.010529
116	2003	December	0.035791	0.093283	0.053997	0.011535
117	2004	December	0.033880	0.096421	0.049199	0.012203
118	2005	December	0.044176	0.095704	0.059201	0.010836
119	2006	December	0.030439	0.076107	0.045716	0.009322
120	2007	December	0.035195	0.098207	0.054561	0.011384

Diffuse_attenuation_coefficient (May-October) 1998-2007



Diffuse_attenuation_coefficient (November-April) 1998-2007



The study showed a clear seasonal variability with highest chlorophyll concentration during July-August and least in April. The inter-annual variability showed an increasing trend from 1998 to 2007.