Summary and Conclusions

The 3-D circulation of Indian Ocean is very complex and is controlled by both local and remote forcings because of the peculiar geographical and climatological setting. Identification of these forcing functions that control the large scale variability of circulation and numerical simulation of highly time-dependent 3-D circulation should be the objective of any modeling study. To determine the physical and dynamical mechanism underlying the variability of circulation of the Indian Ocean, a variety of ocean circulation models such as diagnostic, semi-diagnostic and prognostic types should be developed for the Indian Ocean area. 3-D circulation models that could be run in diagnostic, semi-diagnostic and prognostic models are the best for the study of circulation in the Indian Ocean. There have been many studies, both observational and modeling, on the circulation and watermass characteristics of the tropical Indian Ocean. Detailed literature survey shows that practically all studies were focused on the theoretical or modeling aspects of a particular current or on a particular basin of the Indian Ocean such as Arabian Sea or Bay of Bengal. These existing models mainly deal with upper layer circulation, and there is hardly any information on the circulation in the intermediate, deep and bottom waters.

One of the main objectives of this thesis is to adapt and configure a fully non-linear, primitive equation type, sigma co-ordinate, 3-dimensional circulation model to the entire Indian Ocean area which can be run on diagnostic, semi-diagnostic and prognostic modes. Such a model could identify both local and remote forcings and also could simulate the highly time-varying circulation of the Indian Ocean.

The thesis basically contains six chapters including the introductory chapter. In the introductory chapter of the thesis, boundaries of Indian Ocean and some of the unique features of Indian Ocean and its seasonal reversing currents because of its peculiar geological, geographical and climatic settings are presented. This chapter also contains information on the various observational and mathematical modeling studies conducted since last five decades.

In chapter-II, a detailed description of the state of art 3-dimensional, sigma co-ordinate model that has been adapted to the tropical Indian Ocean is given. The model description includes details such as boundary and initial conditions, numerical solution procedure and some unique features and parameters of the model including the forcing parameters such as wind field, heat and fresh water fluxes.

Chapter-III addresses the problem of understanding the relative roles of steady local forcing of wind and internal density field in driving the circulation of the Indian Ocean by running the model in semi-diagnostic mode. The primitive equation type, sigma co-ordinate, 3-dimensional model developed at the Institute of Numerical Mathematics, Russia, has been adapted and configured to the Indian Ocean area between the latitudes 36° S - 29.5 N and longitudes 22° E - 142 E . The peculiarities of sigma co-ordinate model in semi-diagnostic mode, forcing parameters of the model, results of calculation of steady state currents and circulation were presented. The spatial variability of mean currents in the upper layers of Arabian Sea, Bay of Bengal, equatorial and southern tropical Indian Ocean were presented under separate headings.

(i) Circulation in the Arabian Sea

Model calculations for the month of January showed that the westward flowing North Equatorial Current (North East Monsoon Current) was present in the latitudinal belt $(2^{\circ}N - 6^{\circ}N)$, and i t impinged on the Somali coast with one branch flowing towards north-northeast and the other branch flowing southwestwards along the coast as Somali Current (SC). The model reproduced the northward flowing East African Coastal Current

(EACC), the West India Coastal Current (WICC) present during the winter season. All the major surface currents were discernable at 50m. depth also, but the strength of the current was considerably reduced.

Model calculations on mean currents in the upper layers of Arabian Sea for the month of April showed that the westward flowing Northeast Monsoon Current (NMC) was present in the latitudinal belt 5^N -10^N, its intensity was considerably reduced as compared to January month. Model reproduced a weak equatorward flowing WICC all along the west coast of India in the upper 50m. water column.

Model calculation for the month of July showed the presence of northeastward flowing Somali Current, the Southern Gyre (SG) and the Great Whirl (GW). The Somali Current was found to be an extension of the EACC which flows northward across the equator. It was found that the Somali Current system was mainly forced by the alongshore wind associated with the Findlater Jet and the internal density field. The South West Monsoon Current (SMC) was found to be primarily wind-driven, although the influence of density gradient on its strength and direction could be seen to a certain extent. The model calculation showed the presence of WICC along the west coast of India during July.

(ii) Circulation in the Bay of Bengal

Model calculations for the month of January showed that large anticyclonic gyre was formed in the entire Bay of Bengal north of 8[°]N. The East India Coastal Current (EICC) associated with the wind-driven anticyclonic gyre in the Bay of Bengal flowed poleward all along the east coast of India. It was inferred that the combined effect of local wind forcing and thermal structure are responsible for the observed circulation in the Bay of Bengal during January. The EICC flowed poleward along the east coast of India during April and the model computed EICC features fully match with the

ship-drift data. Both the thermohaline forcing and wind stress fields are responsible for the formation of poleward flowing EICC during April.

During July, the EICC does not flow poleward as expected along the east coast of India. Model results showed that there was a northward flow north of 10[°]N and southward flow south of it. The probable reasons for the northward surface EICC north of 10[°]N and southward EICC south of it can be explained in terms of the prevailing wind stress curl pattern. During summer, the wind stress curl is cyclonic in the south western part of Bay of Bengal and this cyclonic curl could drive a southward flow south of 10[°]N. Signatures of the poleward flowing EICC could be seen at 50[°]M. and 150[°]M. During the month of October, a weak southward flowing EICC was reproduced by the semi-diagnostic model very near to the coastline. Small cyclonic type of circulation cells were observed in the central Bay of Bengal.

(iii) Mean currents in the equatorial and southern tropical Indian Ocean

The semi-diagnostic model calculations for the month of January showed that the South Equatorial Current (SEC) flowed primarily west with a magnitude of approximately 30 cm/s between 10°S and 20°S. The computed results fully agree with the ship drift data compiled by Cutler and Swallow (1984). The equatorial circulation at 150m. depth and 300 m. depth was dominated by the presence of a strong eastward flowing Equatorial Under Current (EUC) between 5°N and 5°S.

During the month of April, the Wyrtki Jet could be traced across the equatorial regions between 2°S and 2°N and its magn itude varied between 20 cm/s and 30 cm/s. All the permanent currents in the southern tropical Indian Ocean such as SEC, EACC, NEMC and SEMC during April were reproduced by the model.

One important finding during the month of August was the appearance of EUC that flowed eastward in the equatorial belt at 150m. and 300m. depths. The semi-diagnostic model has been successful in reproducing all the permanent current systems in the Indian Ocean including the undercurrents over the entire annual cycle.

Chapter-IV deals with the simulation of the climatic currents in the Indian Ocean. Better understanding of the ocean dynamics can be achieved only by using state-of-art simulation models. The time-varying circulation of the Indian Ocean is controlled by both the local forcing and remote forcing by long waves. The main goal of this chapter is to study the large scale, time dependent, 3-dimensional circulation of the Indian Ocean and to understand the dynamical mechanism responsible for the observed circulation. Like semi-diagnostic model, the simulation model has been adapted and configured to the Indian Ocean area between the latitudinal belt 36°S - 29.5°N and longitude belt $22^{\text{C}} - 142^{\text{C}}$ with a resolution of 1° x 0.5° in longitude and latitude directions respectively. The model was integrated forward with a time step of 3 hours for 10 years using surface climatological forcing which were applied with cyclic condition in time. The solution attained semi-stationary by the 10th year of simulation. The 10th year simulation results were compared with existing observations and with results from SODA (Simple Ocean Data Assimilation).

The circulation obtained from the simulation model are in agreement with the SODA results and also with available observations. All the current systems in the Indian Ocean were well reproduced in the model along with their under currents and gyres. The volume transport of major currents were computed and compared with the transport estimates available in the literature. The volume transport was found to be comparable with the existing estimates for different currents. The dynamics of each of the current systems were explained by comparing the simulation model results with the semi-diagnostic model results. Thus the role of local verses remote forcing on the computed circulation of the Indian Ocean has been brought out in this chapter.

Results of circulation models vary depending on its dynamics, spatial resolution, numerical scheme, the parameterization of physical processes and the forcing parameters of which the later, forcing parameters have significant effect on the model results. Wind stress is the main mechanical forcing term that influences significantly the upper layer circulation. One sensitivity experiment was carried out by forcing the simulation model with the winds derived from the Quikscat Scatterometer. Results of the response of the tropical Indian Ocean to Quikscat winds and NCEP/NCAR wind field are presented in chapter-V.

Large scale zonal wind stress difference was observed between the two wind data sets over the equatorial Indian Ocean. The monthly mean response of the model to two wind data sets were compared for mass transport stream function, and surface velocity for the month of January, July, April and October representing the four seasons. The model driven with Quikscat winds reproduced equatorial circulation features more realistically than the NCEP forcing. The current system in the equatorial Indian Ocean was intensified when the model was driven with Quikscat wind forcing. The Equatorial Jets (EJ's) of the Indian Ocean were strengthened by about 20 cm/s during the month of April and October when the model was driven with Quikscat wind forcing. The sensitivity experiment revealed that the surface circulation was mainly controlled by wind stress forcing and hence the thermodynamic forcing play minor role at the upper levels. Alekseev, V.V. and Zalesny, V.B. (1993). Numerical model of large scale dynamics of ocean : Numerical processes and systems : Edited by G.I. Marchuk, vol.10, Moscow : Nayka, 232-252 (in russian).

Anderson, D. L. T. and Carrington, D. J. (1993). Modeling inter-annual variability in the Indian Ocean using momentum fluxes from the operational weather analyses of the United Kingdom Meteorological Office and European Centre for Medium Range Weather Forecasts. *J. Geophys. Res.*, *98*, 12483–12499.

Anderson, D. L. T. and Moore, D. W. (1979). Cross-equatorial inertial jets with special relevance to very remote forcing of the Somali Current. *Deep-Sea Res.*, *26*, 1–22.

Anderson, D. L. T., Carrington, D. J., Correy, R. and Gordon, C. (1991). Modeling the variability of the Somali Current. *J. Mar. Res.*, *49*, 659–696.

Anderson, D. L. T., Carrington, D. J., Correy, R. and Gordon, C. (1993). The temporal evolution of equatorial currents in the Indian Ocean. Metrological Office Climate Research Tech. Note 23, Bracknell, Berkshire, United Kingdom, 51 pp. [Available from U. K. Meteorological Office, Bracknell, Berkshire RG12 252, United Kingdom].

Antony, M. K. (1990). Northward undercurrent along west coast of India during upwelling — Some inferences. *Ind. J. Mar. Sci.*, *19*, 95–101.

Babu, M.T., Prasanna Kumar, S. and Rao, D.P. (1991). A subsurface cyclonic eddy in the Bay of Bengal, *J. Mar. Res.*, 49, 402-410.

Bagno, A.V., Dianskii, N.A. and Moshoukin, S.N. (1996b). Interaction of ocean surface temperature anomalies with the circulation in the north Atlantic. *Oceanology*, Vol.36, 652-661. (Translated from Oceanologia, Vol. 36, 5, 693-703).

Bagno, A.V., Garashchuk, R.V. and Zalesny, V.B. (1996a). A model of large scale ocean circulation and sea ice evolution, *Oceanology*, Vol. 36, 2, 179-187.

Bahulayan, N. and Shaji, C. (1996). Diagnostic model of 3-D circulation in the Arabian Sea and western equatorial Indian Ocean. *Proc. Indian Nat. Sci. Acad.* 62: 325347.

Bahulayan, N., Shaji, C., Rao, A.D. and Dube, S.K. (1997). Sensitivity experiments with an adaptation model of circulation of western tropical Indian Ocean. *Indian J. Mari. Sci.*, 96: 110.

Bahulayan, N. and Unnikrishnan, A.S. (1992). Simulation of barotropic wind-driven circulation in the upper layers of Bay of Bengal and Andaman Sea during the southwest and northeast monsoon seasons with observed winds. *Proc. Indian Acad. Sci. (Earth and Planet. Sci.)*, vol.101, 1, 47-66.

Bahulayan, N. and Varadachari, V.V.R. (1986). Numerical model of wind driven circulation in the Bay of Bengal. *Ind. J. Mar. Res.*, vol.15, 8-12.

Banse, K. (1968). Hydrography of the Arabian Sea shelf of India and Pakistan and effects on demersal fishes. *Deep Sea Res.*, 15, 45-79.

Bleck, R. and Smith, L.T. (1990). A wind driven isopycnic coordinate model of the north and equatorial Atlantic ocean. 2. Model development and supporting experiments. *J. Geophys. Res.*, 95, 3273-3285.

Blumberg, A. F. and Mellor, G.L. (1987). A description of a threedimensional coastal ocean circulation model, edited by N. Heaps, 208 pp., American Geophysical Union.

Bohm, E., Morrison, M., Manghani, V.J.A., Kim, H.S., Flagg, C.N. (1999). Remotely sensed and acoustic Doppler observations of the Ras al Hadd Jet in 1994-95. *Deep-Sea Res. II*, 46, 1531-1549.

Brandt, P., Stramma, L., Schott, F., Fischer, J., Dengler, M., Quadfasel, D., (2002). Annual Rossby waves in the Arabian Sea from TOPEX/POSEIDON altimeter and in-situ data. *Deep-Sea Res. II*, 49 (7), 1197-1210.

Brown, O.B., Bruce, J.C. and Evans, R. H. (1980). Evolution of sea surface temperature in the Somali basin during the southwest monsoon of 1979. *Science*, 209, 595-597.

Bruce, J.G., (1973). Large scale variations of the Somali Current during southwest monsoon, 1970. *Deep-Sea Res.* 20, 837-846.

Bruce, J.G., (1979): Eddies off the Somali coast during the southwest monsoon. *J. Geophys. Res.* Vol. 84, 7742-7748.

Bruce, J. G., Johnson, D. R., and Kindle, J. C., (1994). Evidence for eddy formation in the eastern Arabian Sea during the northeast monsoon. *J. Geophys. Res.*, *99*, 7651–7664.

Bruce, J.G., Kindle, J.C., Kantha, L. H., Kerling, J. L. and Bailey, J. F. (1998). Recent observations and modeling in the Arabian Sea Laccadive High region, *J. Geophys. Res.*, 103, 7593-7600.

Bryan, K. (1969). A numerical method for the study of the circulation of the world ocean, *J. Comput. Phys.*, 4, 347-376.

Bryden, D., San, S. and Bleck, R. (1999). A new approiximation of the equation of state for sea water suitable for numerical models. *J. Geophys. Res.*, 104, c1, 1537-1540.

Cane, M. (1980). On the dynamics of equatorial currents, with application to the Indian Ocean. *Deep-Sea Res.*, *27A*, 525–544.

Cane, M., and Gent, P. (1984). Reflections of low-frequency equatorial waves at arbitrary western boundaries. *J. Mar. Res.*, *4*2, 487–502.

Cane, M. A., and Moore, D. W. (1981). A note on low-frequency equatorial basin modes. *J. Phys. Oceanogr.*, 11, 1578–1584.

Carton, J.A., Chepurin,G., Cao, X. and Giese, B. S. (2000). A Simple Ocean Data Assimilation analysis of the global upper ocean 1950-1995, Part 1: methodology, *J. Phys. Oceanogr.*, 30, 294-309.

Clarke, R. A., and Liu, X. (1993). Observations and dynamics of semiannual and annual sea levels near the eastern equatorial Indian Ocean boundary. *J. Phys. Oceanogr.*, *23*, 386–399.

Cox, M.D. (1970). A mathematical model of Indian Ocean. *Deep-Sea Res.* 17, 4775.

Cox, M.D. (1976). Equatorially trapped waves and the generation of the Somali Current. *Deep- Sea Res.*, Vol. 23., 1139-1152.

Cox, M.D. (1979). A numerical study of the Somali Current eddies. J. *Phys. Oceanogr.*, Vol. 9, 311-326.

Cox, M.D. (1984). A primitive equation, 3-dimensional model of the ocean, GFDL Ocean Group Technical Report No. 1, Geophysical Fluid Dynamics Laboratory, Princeton, NJ.

Cresswell, G. R. and Golding, T. J. (1980). Observations of a southflowing current in the southeastern Indian Ocean. *Deep-Sea Res.,* 27, pp. 449-466

Cutler, A. N., and Swallow, J. C. (1984). Surface currents of the Indian Ocean. (To 25^oS, 100^oE). I. O. S. Technical Rep. 18 7pp.

Dabyshire, M. (1967). The surface waters off the coast of Kerala, South west India. *Deep Sea Res.*, 14, 295-320.

Dengler, M., Quadfasel, D., Schott, F. and Fischer, J. (2002). Abyssal circulation in the Somali Basin. *Deep-Sea Res. Part II*, 9, 1297-1322.

de Ruijter, W.P.M., Ridderinkhof, H. Lutjeharms, J. R. E. Schouten, M. W. and Veth, C. (2002). Observations of the flow in the Mozambique Channel. *Geophys. Res. Lett.*, 29, 140.1-140.3.

Deutsches Hydrographisches Institut. (1960). Monatskarten für den Indischen Ozean. Deutsches Hydrographisches institut, Hamburg, Public. 2422.

Diansky N.A., Bagno A.V., Zalesny V.B., (2002). Sigma model of global ocean circulation and its sensitivity to variations in wind stress. Izv. Atmos. Ocean. Phys. (Engl. Transl.), V.38, No. 5, pp. 477-494.

Donguy, J. R., and Meyers, G. (1995). Observations of geostrophic transport variability in the western tropical Indian Ocean. *Deep- Sea Research*, *42*, 1007–1028.

Dube, S.K., Rao, A.D. and Rao, G.S. (1988). Wind induced circulation of the Bay of Bengal. Computational methods in flow analysis. (H. Nikki and M. Kawahara, eds.), Okayama University of Science Press, 2, 1278-1284.

Dueing, W., and Schott, F. (1978). Measurements in the source region of the Somali Current during the monsoon reversal. J. Phys. Oceanogr., 8, 278-289.

Duing, W. (1970). The monsoon regime of the currents in the Indian Ocean. East west Centre Press, university of Hawai, Honolulu, 68pp.

Eigenheer, A., and Quadfasel, D. (2000). Seasonal variability of the Bay of Bengal circulation inferred from TOPEX/POSEIDON altimetry. *J. Geophys. Res.*, *105*, 3243–3252.

Eriksen, C. C. (1979). An equatorial transect of the Indian Ocean, *J. Mar. Res.*, 37, 215-232.

Esenkov, O. E. and Olson, D. B. (2002). A numerical study of the Somali Coastal Undercurrent. *Deep-sea Res. II*, Topical Studies in Oceanography, Vol. 49, 7-8, 1253-1277.

Fieux, M., Andrie[´], C., Delecluse, P., Ilahude, A. G., Kartavtseff, A., Mantisi, F., Molcard, R., and Swallow, J. C. (1994). Measurements within the Pacific-Indian Oceans Throughflow region. *Deep-sea Res. I*, *41*, 1091–1130.

Fieux, M., Molcard, R., and Ilahude, A. G. (1996). Geostrophic transport of the Pacific-Indian Oceans Throughflow. *J. Geophys. Res.*, 101, 12421–12432.

Fischer, J., Schott, F., and Stramma, L. (1996). Currents and Transports of the Great Whirl-Socotra Gyre System during the Summer Monsoon August 1993. *J. Geophys. Res.*, 101, 3573–3587.

Flagg, C. A., and Kim, H.-S. (1998). Upper ocean currents in the northern Arabian Sea from shipboard ADCP measurements collected during the 1994–1996 U.S. JGOFS and ONR Programs. *Deep-Sea Res. II*, 45, 1917–1959.

Fu, L. (1986). Mass, heat and freshwater fluxes in the South Indian Ocean. *J. Phys. Oceanogr.*, 16, 1683-1693.

Gent, P.R., O'Neill. K. and Cane, M. A. (1983). A model of the semiannual oscillation in the Equatorial Indian Ocean. *J. Phys. Oceanogr.*, Vol. 13, 2148-2160.

Godfrey, J. S., and Ridgeway, K. R. (1985). The large scale environment of the poleward flowing Leeuwin Current, western Australia: longshore steric height gradients, wind stresses and geostrophic flow. *J. Phys. Oceanogr.*, *15*, 481–495.

Gopalakrishna, V.V., Pednekar, S.M. and Murty, V. S. N. (1996). T-S variability and volume transport in the central Bay of Bengal during south west monsoon. *Ind. J. Mar. Sci.*, 1996, Vol.25, 50-55.

Gopalakrishna, V.V., Sastry, J.S. (1985). Surface circulation over the shelf off the east coast of India during the south west monsoon. *Indian J. Mar. Sci.*, 14, 62-65.

Gordon, A. L., and Susanto, R. D. (1999). Makassar Strait transport: initial estimate based on Arlindo results. *MTS Journal*, *32*, 34–45.

Gründlingh, M. L. (1988). Review of cyclonic eddies in the Mozambique Ridge Current. *South African J. Mar. Sci.*, 6, 193-206.

Gründlingh, M. L., Carter, R.A., and Stanton, R.C. (1991). Circulation and water properties of the Southwest Indian Ocean, Spring 1987. *Progr. Oceanogr.*, 28, 4, 305-342.

Hacker, P., Firing, E., Hummon, J., Gordon, A. L., and Kindle, J.C. (1998). Bay of Bengal currents During the Northeast Monsoon. *Geophys. Res. Lett.*, 25, 2769-2772.

Han, W., McCreary, J. P. Jr., Anderson, D. L. T., and Mariano, A. J. (1999). On the dynamics of the eastward surface jets in the equatorial Indian Ocean. *J. Phys. Oceanogr.*, *29*, 2191–2209.

Harris, T. F. W. (1972). Sources of the Agulhas Current in the spring of 1964. *Deep-Sea Res.*, 19, 633-650.

Hastenrath S. (1989). The monsoonal regimes of upper-hydrospheric structure in the tropical Indian Ocean. *Atmosphere-Ocean.* Vol. 27, 478-507.

Hastenrath, S., and Greischar, L. (1991). The monsoonal current regimes of the tropical Indian Ocean: observed surface flow fields and their geostrophic and wind-driven components. *J. Geophys. Res.*, 96, 12619-12633.

Hellerman, S., and Rosenstein, M. (1983). Normal monthly wind stress over the world ocean with error estimates. J. Phys. Oceanogr., 13, 1093-1104.

Holloway, P. E. (1995). Leeuwin Current observations on the Australian North West Shelf, May–June 1993. *Deep-Sea Res. I*, *42*, 285–305.

Hurlburt, H. E., and Thompson, J. D. (1976). A numerical model of the Somali Current. *J. Phys. Oceanogr.*, 6, 646–664.

Jensen, T. G. (1991). Modeling the seasonal undercurrents in the Somali Current system. *J. Geophys. Res.*, *96*, 22151–22167.

Jensen, T. G. (1993). Equatorial variability and resonance in a winddriven Indian Ocean Model. *J. Geophys. Res.*, *98*, 22533–22552.

Johannessen, O.M., Subbaraju, G. and Blindheim, J. (1981). Seasonal variation of the oceanographic conditions off the sw coast of India during 1971-75. Fisk. Dir. Skr. Ser. HavUnders, 18, 247-261.

Kalnay. E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, L., Saha, S., White, G., Woollen, J., Zhu, Y., Chellah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Roy Jenne and Dennis Joseph, (1996). The NCEP/NCAR 40-year reanalysis project. *Bull. of American. Met. Soc.*, 77, 437-471.

Kindle, J. C., and Thompson, J. D. (1989). The 26- and 50-day oscillations in the western Indian Ocean: model results. *J. Geophys. Res.*, *94*, 4721–4736.

Knox, R. (1976). On a long series of measurements of Indian Ocean equatorial currents near Addu Atoll. *Deep-Sea Res.*, 23, 211–221.

Koninklijk Nederlands Meteorologisch Instituut. (K.N.M.I.) (1952). Indian Ocean, oceanographic and meteorological data. Koninklijk Nederlands Meteorologisch Institut, publication 135.

La Fond, E. C. (1957). Oceanographic studies in the Bay of Bengal. *Proc. Indian Acad. Sci.(B)*, 46(1):1-46

Le Blond, P.H. and Mysak, L. A. (1980). Waves in the Ocean, Elseveir Scientific Publishing Co., Amsterdam.

Leetmaa, A. (1982). Development of the flow field during the onset of the Somali Current. *J. Phys. Oceanogr.* 12, 23-28.

Leetmaa, A., and Stommel, H. (1980). Equatorial current observations in the western Indian Ocean during 1975 and 1976. *J. Phys. Oceanogr.*, *10*, 258–269.

Legeckis, R. (1987). Observation of a western boundary current in the Bay of Bengal. *J. Geophys. Res.*, Vol. 92, 12974-12978.

Levitus, S. (1994). World Ocean Atlas 1994-CD-ROM dataset : U. S. Department of Commerce, National Oceanic and Atmospheric Administration, National Environmental Satellite Data and Information Service, National Oceanographic Data Centre, Ocen Climate Laboratory.

Lighthill, M.J., (1969). Dynamic response of the Indian Ocean to onset of the southwest monsoon. Phil.Trans. A., 265,1159,45-92.

Lin, L. B., and Hurlburt, H. E. (1981). Maximum simplification of nonlinear Somali Current dynamics. In M. J. Lighthill, and R. P. Pearce (Eds.), *Monsoon dynamics*. Cambridge: Cambridge University Press.

Lisan Yu, O'Brien, J. J., and liayan Yang. (1991). Oon the remote forcing of the circulation in the Bay of Bengal. *J. Geophys. Res.*, Vol. 96, C2, 20,449-20,454.

Lutjeharms, J. R. E., Bang, N.D., and Duncan, C. P. (1981). Characteristics of the currents east and south of Madagascar. *Deep-Sea Res.*, 28, 879-899.

Lutjeharms, J.R.E and Van Ballegooyen, R. C. (1988). The Retroflection of the Agulhas Current. J. Phys. Oceanogr., 18, 11, 1570-1583.

Luther, M. E., and O'Brien, J. J. (1985). A model of the seasonal circulation in the Arabian Sea forced by observed winds. *Progr. Oceanogr.*, *14*, 353–385.

Luyten, J.R. and Swallow, J. C. (1976). Equatorial Under currents. *Deep-Sea Res.*, 23, 999-1001.

Marchuk, G.I. and Sarkisyan, A. S. (1988). Mathematical modeling of ocean circulation. Springer-Verlg, Berlin, 292 pp.

Mariano, A.J., Ryan, E. H., Perkins, B. D., and Smithers, S. (1995). The Mariano Global Surface Velocity Analysis 1.0. USCG Report CG-D-34-95, 55 pp.

Mc Creary, J. P., Kohler, K.E., Hood, R.R., Olson, D.B. (1996). A fourcomponent ecosystem model of biological activity in the Arabian Sea. *Prog. Oceanogr.*, 37: 193-240.

Mc Creary, J. P., and Kundu, P. K. (1985). Western boundary circulation driven by an alongshore wind: with application to the Somali Current system. *J. Mar. Res.*, *43*, 493–516.

Mc Creary, J. P., and Kundu, P. K. (1988). A numerical investigation of the Somali Current during the Southwest Monsoon. *J. Mar. Res.*, *46*, 25–58.

Mc Creary, J. P. Jr., Kundu, P. K., and Molinari, R. L. (1993). A numerical investigation of dynamics, thermodynamics and mixedlayer processes in the Indian Ocean. *Progr. Oceanogr.*, *31*, 181–244.

Mc Phaden, M. (1982). Variability in the central Indian Ocean. Part I: Ocean dynamics. *J. Mar. Res.*, *40*, 157–176.

Mesinger, F. and Arakawa, A. (1976). Numerical methods used in atmospheric models, 1, 66 pp. GARP Publications Series 17. WMO, Geneva, Switzerland

Meyers, G., Bailey, R. J., and Worby, A. P. (1995). Geostrophic transport of Indonesian Throughflow. *Deep-Sea Res.*, *4*2, 1163–1174.

Michaelis, G. (1923). Die Wasserbewegung an der Oberfläche des Indischen Ozeans im Januar und Juli. Veröffentlichungen des Instituts für Meereskunde an der Universität Berlin, n.f. A8(16): 32 pp.

Molinari, R. L., Olson, D., and Reverdin, G. (1990). Surface current distributions in the tropical Indian Ocean derived from compilations of surface buoy trajectories. *J. Geophys. Res.*, *95*, 7217–7238.

Möller, L. (1929). Die Zirkulation des Indischen Ozeans; auf Grund von Temperatur- und Salzgehaltstiefenmessungen und Oberflächenstrombeobachtungen. Veröffentlich-ungen des Instituts für Meereskunde an der Universität Berlin, n.f. A21: 48 pp. Morrow, R., and Birol, F. (1998). Variability in the southeast Indian Ocean from altimetry: Forcing mechanism for the Leeuwin Current. *J. Geophys. Res.*, *103*, 18529–18544.

Murty, V.S.N., Sarma, Y.V.B., Rao, D.P., Murty, C.S. (1992a). Water characteristics, mixing and circulation in the Bay of Bengal during southwest monsoon. *J. Mar. Res.*, 40, -228

Murty, V.S.N., Sarma, Y.V.B., Babu, M.T., Rao, D.P. (1992b). Hydrography and circulation in the northwestern Bay of Bengal during the retreat of southwest monsoon. *Proc. Ind. Acad. Sci., Earth and Planet. Sci.,* 167-75

O'Brien, J. J., and Hurlburt, H. E. (1974). An equatorial jet in the Indian Ocean theory. Science, *184*, 1075–1077.

Pacanowski, R.L., Griffins, S.M. (1999). The MOM3 Mannual. Geophysical Fluid Dynamics Laboratory/NOAA, Priceton, USA : pp.680.

Pacanowski, R. C., and Philander, S. G. H. (1981). Parameterization of vertical mixing in numerical models of the tropical ocean. *J. Phys. Oceanogr.*, 11, 1442-1451.

Paech, H. (1926). Die Oberflächenströmungen um Madagascar in ihrem Jährlichen Gang. Veröffentlichungen des Instituts für Meereskunde an der Universität Berlin, n.f. A(16): 39 pp.

Park, Y.-H. and Gamberoni, L. (1995). Large scale circulation and its variability in the south Indian Ocean from TOPEX/POSEIDON altimetry. *J. Geophys. Res.*, 100, 24911-24929

Potemra, J. T., Luther, M. E., and O'Brien, J. J. (1991). The seasonal circulation of the upper ocean in the Bay of Bengal. *J. Geophys. Res.*, *96*, 12667–12683.

Quadfasel, D., and Schott, F. (1983). Southward subsurface flow below the Somali Current. *J. Geophys. Res.*, *88*, 5973–5979.

Rao, A.D., Dube, S.K., and Sinha, P.C. (1992). Wind-driven seasonal ocean circulation around southern Indian peninsula. Indian J. Mar. Sci., 21, 196-200.

Rao, R. R., Molinari, R. L., and Festa, J. F. (1989). Evolution of the climatological near-surface thermal structure of the tropical Indian Ocean: 1. Description of mean monthly mixed layer depth, and sea surface temperature, surface current, and surface meteorological fields. *J. Geophys. Res.*, *94*, 10801–10815. Rao, R. R., Molinari, R. L., Festa, J. (1991). Surface meteorological and near surface oceanographic atlas of the tropical Indian Ocean. NOAA Technical Memorandum, ERL west Monsoon of 1979. Science 209, 595 597. AOML-69.

Reppin, J., Schott, F. A., Fischer, J., and Quadfasel, D. (1999). Equatorial currents and transports in the upper central Indian Ocean: annual cycle and interannual variability. *J. Geophys. Res.*, 104, 15495–15514.

Reverdin, G. (1987). The Upper Equatorial Indian Ocean. The climatological Seasonal Cycle. *J. Phys. Oceanogr.*, Vol. 17, 7, 903-927.

Reverdin, G. and Fieux, M. (1987). Sections in the western Indian Oceanvariability in the temperature structure. *Deep-Sea Res.*, 34, 60162.

Reverdin, G., Luyten, J. (1986). Meanders in the near-surface equatorial jet in the Indian Ocean. *J. Phys. Oceanogr.*, **16**(6), 1088-1100.

Rochford, D.J. (1964). Salinity maxima in the upper 1000 metres of the North Indian Ocean. *Aust. J. Mar. Freswat. Res.*, 15, 1-24.

Salvekar, P. S., Behra, S. K., Ganer, D. W., Deo, A. D., and Rahul Reddy, P. (2000). Behavior pf north Indian Ocean during the onset phases of southwest monsoon using satellite winds. IITM-SAC collaborative project Research Report, 11-19.

Sarkisyan, A.S. and Demin, Yu. L. (1983). A semi-diagnostic method of sea current calculation. Large scale oceanographic experiments in the WCRP publication series, Tokyo, 2(1), 201-214.

Sætre, R. and Jorge da Silva, A. (1984). The circulation of the Mozambique Channel. *Deep-Sea Res.*, 31(5): 485-508.

Scherbinin, A.D., Arse'yev, V. S., and Golddin, YU. A. (1979). The western boundary current of the Bay of Bengal, Dok6. Acad. Sci. USSR, Earth Sci. Sect., English Translation, 244, 167-168.

Schott, F. (1983). Monsoon response of the Somali Current and associated upwelling. *Progr. Oceanogr.*, Vol. 12, 357-382.

Schott, F. (1986). Seasonal variation of cross-equatorial flow in the Somali Current. *J. Geophys. Res.*, *91*, 10581–10584.

Schott, F. and Fieux, M. (1985). The Somali Current in autumn 1984, before the onset of the northeastern monsoon. *Nature*, 315, 50-52.

Schott, F., Fieux, M., Kindle, J., Swallow, J., and Zantopp, R. (1988). The boundary currents east and north of Madagascar. Part II: Direct

measurements and model comparisons. J. Geophys. Res., 93, 4963-4974.

Schott, F., and Fischer, J. (2000). Winter monsoon circulation of the northern Arabian Sea and Somali Current. *J. Geophys. Res.*, 105, 6359–6376.

Schott, F., Fischer, J., Garternicht, U., and Quadfasel, D. (1997). Summer monsoon response of the Northern Somali Current, 1995. *Geophys. Res. Lett.*, *24*, 2565–2568.

Schott, F., and Quadfasel, D. (1982). Variability of the Somali Current and associated upwelling. *Progr. Oceanogr.*, *12*, 357–381.

Schott, F., J. Reppin, D. Quadfasel and J. Fischer, 1994: Currents and transports in the Monsoon Current south of Sri Lanka. *J. Geophys. Res.*, **99**, 25,127-25,141.

Schott, F., Swallow, J. C., and Fieux, M. (1990). The Somali Current at the equator: annual cycle of currents and transports in the upper 1000 m and connection to neighboring latitudes. *Deep-Sea Res.*, *37*, 1825–1848.

Semtner, Jr., A.J. (1974). An oceanic general circulation model with bottom topography. In: Numerical Simulation of Weather and Climate, Tech. Rept. No. 9, UCLA Department of Meteorology.

Senan, R., Sengupta, D. and Goswami, B. N. (2003). Intraseasonal "Monsoon Jets" in the equatorial Indian Ocean. *Geophys. Res. Lett.*, 30, 14,1750, doi:10.1029/2003GL017583.

Sewell, R.B.S. (1929). Temperature and salinity of the surface waters of the by of Bengal and Andaman Sea with reference to the Laccadive Sea. *Memoirs of the Asiatic Society of Bengal*, 9, 207-356.

Shaji, C., Bahulayan, N., Dube, S.K. and Rao, A.D. (1997). A semidiagnostic calculation in the western Indian Ocean, *Indian J. Mar. Sci.*, 26, 241 249.

Shankar, D., Mc Creary, J. P., Han, W., and Shetye, S. R. (1996). Dynamics of theEast India Coastal Current. 1. Analytic solutions forced by interior Ekman pumping and local alongshore winds. *J. Geophys. Res.*, Vol. 101, C6, 13975-13991.

Shankar, D., and Shetye, S. R. (1997). On the dynamics of the Lakshadweep high and low in the southeastern Arabian Sea. *J. Geophys. Res.*, 102, 12551–12562.

Shankar, D. Vinaychandran, P. N., and Unnikrishnan, A.S., (2002). The monsoon currents in the North Indian Ocean, *Progr. Oceanogr.*, 52(1),63-120.

Shetye, S. R., Gouveia, A. D., Shenoi, S. S. C., Sundar, D., Michael, G. S., Almeida, A. M., and Santanam, K. (1990). Hydrography and circulation off the west coast of India during the Southwest Monsoon 1987. *J. Mar. Res.*, *48*, 359–378.

Shetye, S. R., Gouveia, A. D., Shenoi, S. S. C., Sundar, D., Michael, G. S., and Nampoothiri, G. (1991a). The western boundary current of the seasonal subtropical gyre in the Bay of Bengal. *J. Geophys. Res.*, *98*, 945–954.

Shetye, S. R., Gouveia, A. D., Shenoi, S. S. C., Sundar, D., Michael, G. S., and Nampoothiri, G. (1993). The western boundary current of the seasonal subtropical gyre in the Bay of Bengal. *J. Geophys. Res.*, *98*, 945–954.

Shetye, S. R., Gouveia, A. D., Shenoi, S. S. C., Vinayachandran, P. N., Sundar, D., Michael, G. S., and Nampoothiri, G. (1996). Hydrography and circulation in the western Bay of Bengal during the northeast monsoon. *J. Geophys. Res.*, 101, 14011–14025.

Shenoi, S. S. C., Saji, P. K., and Almeida, A. M. (1999). Near-surface circulation and kinetic energy in the tropical Indian Ocean derived from Lagrangian drifters. *J. Mar. Res.*, *57*, 885–907.

Shetye, S. R., and Shenoi, S. S. C. (1988). The seasonal cycle of surface circulation in the coastal North Indian Ocean. *Proc. Ind. Acad. Sci. (Earth and Planet. Sci.)*, *97*, 53–62.

Shi, W., Morrison, J. M., Bo⁻hm, E., and Manghnani, V. (1999). Remotely sensed features in the US JGOFS Arabian Sea Process Study. *Deep-Sea Res. II*, *46*, 1551–1575.

Simmons, R.C., Luther, M.E., O'Brien, J. J., and Legler, D. M. (1988). Verification of a numerical ocean model of the Arabian Sea. *J. Geophys. Res.*, C12, 15437-15453.

Smith, R. L., Huyer, A., Godfrey, J. S., and Church, J. A. (1991). The Leeuwin Current off western Australia, 1986–1987. *J. Phys. Oceanogr.*, 21, 322–345.

Sprintall, J., Gordon, A., Murtugudde, R., and Susanto, D. (2000). A semiannual Indian Ocean forced Kelvin wave observed in the Indonesian Seas in May 1997. *J. Geophys. Res.*, 105, 17217-17230. Stramma, L. (1992). The South Indian Ocean Current, *J. Phys. Oceanogr.*, 22, 412-430.

Stramma, L., Brandt, P., Schott, F., Quadfasel, D., and Fischer, J. (2002). Winter and summer monsoon water mass and heat transport changes in the Arabian Sea near 8^oN. *Deep-Sea Res. II*, 49, 1173-1195.

Stramma, L., Fischer, J., and Schott, F. (1996). The flow field off southwest India at 8[°]N during the southwest monsoon of August 1993. *J. Mar. Res.*, *54*, 55–72.

Stramma, L. and Lutjeharms, J. R. E. (1997). The flow field of the subtropical gyre of the South Indian Ocean, *J. Geophys. Res.*, 102(C3), 5513-5530.

Suryanarayana, A., Murty, V.S.N., Rao, D.P. (1993). Hydrography and circulation of the Bay of Bengal during early winter, 1983. *Deep-Sea Res.*, 40, 1, 205-217.

Swallow, J. (1967). The equatorial undercurrent in the western Indian Ocean in 1964. *Studies in Tropical Oceanography*, *5*, 15–36.

Swallow, J. C., and Bruce, J. C. (1966). Current measurements off the Somali coast during the southwest monsoon of 1964. *Deep-Sea Res.*, 13, 861–888.

Swallow, J. and Fieux, M. (1982). Historical evidence for two gyres in the Somali Current. *J. Mari. Res.*, 40, 747-755.

Swallow, J. C., Fieux, M., and Schott, F. (1988). The boundary currents east and north of Madagascar, Part I: Geostrophic currents and transports. *J. Geophys. Res.*, 93, 4951–4962.

Swallow, J. C., Molinari, R. L., Bruce, J. G., Brown, O. B., and Evans, R. H. (1983). Development of near-surface flow pattern and water mass distribution in the Somali Basin in response to the southwest monsoon of 1979. *J. Phy. Oceanogr.*, 13, 1398–1415.

Swallow, J. C., Schott, F., and Fieux, M. (1991). Structure and transport of the East African Coastal Current. *J. Geophys. Res.*, *96*, 22254–22267.

Thompson, R.O.Y.R. (1984). Observation of the Leeuwin current off western Australia. *J. Phys. Ocenogr.* 14, 623-628.

Toole, J. M., and Warren, B. (1993). A hydrographic section across the subtropical South Indian Ocean. *Deep-Sea Res. I*, *40*, 1973–2019.

Traft, B. A. (1967). Equatorial undercurrent of the Indian Ocean, (1963). *Studies in Tropical Oceanography*, *40*, 3–14.

UNESCO (1969). Discharge of selected rivers of the world, Vol.1, Paris, NUESCO.

Unnikrishnan, A.S., Murty, V. S. N., Babu, M. T., Gopinathan, C. K., and Charyulu, R. J. K. (2001). Anomalous current structure in the eastern equatorial Indian Ocean during the south-west monsoon of 1994. *Mar. Freshwater Res.*, 52, 5, 727 - 734

Vinayachandran, P. N., Masamoto, Y., Mikawa, T., and Yamagata, T. (1999a). Intrusion of the Southwest Monsoon Current into the Bay of Bengal. *J. Geophys. Res.*, *104*, 11077–11085.

Vinayachandran, P. N., Saji, N. H., and Yamagata, T. (1999b). Response of the equatorial Indian Ocean to an unusual wind event during 1994. *Geophys. Res. Lett.*, *26*, 1613–1616.

Vinayachandran, P. N., and Yamagata, T. (1998). Monsoon response of the sea around Sri Lanka: generation of thermal domes and anticyclonic vortices. *J. Phys. Oceanogr.*, *28*, 1946–1960.

Visbeck, M., and Schott, F. (1992). Analysis of seasonal current variation in the western equatorial Indian Ocean: direct measurements and GFDL model comparison. *J. Phys. Oceanogr.*, *22*, 1112–1128.

Walter, M. (1997). Die Zirkulation des westlichen tropischen Indischen Ozeans im Sommer 1995. Dipl. Thesis. University of Kiel.

Warren, B., Stommel, H. and Swallow, J.C. (1966). Water masses and patterns of flow in the Somali Basin during the Southwest Monsoon of 1964. *Deep-sea Res.* 13, 825-860.

Woodberry, K. E., Luther, M. E., and O'Brien, J. J. (1989). The wind-driven seasonal circulation in the southern tropical Indian Ocean. *J. Geophys. Res.*, *94*, 17985–18002.

Wyrtki, K. (1961). Physical Oceanography of the southeast Asian waters, vol. 2, 1-195, NAGA Rep., Scripps Institute of Oceanography, University of California, San Diego.

Wyrtki, K. (1971). Oceanographic Atlas of the International Indian Ocean Expedition. National Science Foundation, Washington D.C., 531 pp.

Wyrtki, K. (1973a). An equatorial jet in the Indian Ocean. *Science*, 181, 262–264.

Wyrtki, K. (1973b). Physical oceanography of the Indian Ocean. Vol. 3 of Ecological studies: analysis and synthesis. In B. Zeitschel, and S. A. Gerlach (Eds.), Biology of the Indian Ocean (pp. 1836). London: Chapman and Hall Ltd.

Zalesny, V.B., Dianskii, N.A., Moshoukin, S.N. and Bagno, A.V. (2000). Report on the model of large scale dynamics of ocean, MONAR-1. Russian Academy of Sciences, Institute of Numerical Mathematics, Moscow (in russian).

Zalesny, V.B. and Galkin (1996). Initialisation of the temperature filed in a model of Arabian Sea thermodynamics. *Oceanology*, English translation, Vol.35, 4, 472-484.