

ACTINIARIAN SEA ANEMONE FAUNA OF INDIA

Arun Parulekar
National Institute of Oceanography
Dona Paula, Goa: 403 004.

ABSTRACT

The actiniarian sea anemones, inhabiting marine and estuarine ecosystems, from upper littoral to abyssal depths, from burrowing to adhesive forms and from free living to commensal types, though accounting for a small part of the diverse marine biota of India, yet from an ecological point of view, these anthozoans represent a thoroughly successful faunal group. Out of 40 species, belonging to 33 genera under 17 families, as many as 13 species are reported here for the first time.

The habitat preference was: 13 brackishwater, 24 marine and 3 cosmopolitan. Dominance of sedentary, both attached and adhesive forms, over the burrowing species was evident. Only two species, namely Boloceroides mcmurrichi and Boloceractis gopalayi displayed dual habitat of partly sedentary and partly pelagic. While almost all the burrowing species were represented by solitary forms, the gregariousness, whenever observed, was a redeeming feature of most of the attached and adhesive forms, either free living or commensals.

Though marine in origin, many of the burrowing, sedentary and pelagic species of actiniarian sea anemones have successfully adapted to unpredictable salinity conditions of brackish water as also to the extreme conditions of desiccation within the tide marks. A distinct substrate specificity, whether for attachment to biogenic and/or any other object or burrowing in bottom deposits, is clearly discernible. Biotic associations, either symbiosis or commensalism is found to have developed to perfection in actiniarian sea anemones.)

✓ Inadequacy of knowledge about the magnitude and diversity of sea anemones is a reflection on lack of exploration along and within the extensive marine and estuarine ecosystems of India.

Introduction

The sea anemone fauna of the world oceans, comprising of 42 families, 200 genera and about 800 species, though contributes a very small part of the marine biota as a whole, yet ecologically is a successful group of marine invertebrates (Carlgren 1949). The actiniarian fauna of India, but for the contributions of Annandale (1907 & 1915), Panikkar (1936, 1937, 1937a, 1937b, 1938 & 1939) and Parulekar (1956, 1967, 1968, 1969a, 1969b & 1971) has rather discontinuously been investigated and is better known through new species and new genera, rather than the magnitude and diversity of the faunal itself.

Though the term sea anemone encompasses order Actiniaria, Corallimorpha and Ptychodactiaria, however, due to their wide distribution and sheer numbers, especially in the intertidal region, the

actiniarian sea anemones are far more better studied than the other two. In recent years, the actiniarian sea anemones have received specific attention, for their potential uses in drugs and pharmaceuticals as well as the sentinel organisms for ecological monitoring of estuarine and marine environment (Ross 1967).

As a part of the assessment and evaluation of the biological resources of the Exclusive Economic Zone (EEZ) of India and further under an INDO-US collaborative research programme on bioactive substances from the Indian Ocean, a detailed study on the distribution, abundance, ecology and bioactive characteristics in space and time is being carried out, for the last few years and a lot of information about the sea anemone fauna of India is available by now.

In the present communication, an account of the occurrence, taxonomy and ecobiological features of a number of actiniarian sea anemones from estuarine, mangrove, intertidal, coral, oceanic island, shallow and deep water ecosystems of India is presented and reviewed.

Fauna and Species

The present account enumerates 40 species belonging to 33 genera under 17 families (Table 1). The family Actiniidae represented by 10 genera and 15 species is the most common group of actiniarian sea anemones.

While 31 species were reported by earlier workers, 9 species, namely Halcampa capensis Carlgren 1938; Alicia sansibarensis, Carlgren 1900; Glyphoperidium bursa, Roule 1909; Glyphostylum calyx, Roule 1909; Parabunodactis (Bunodactis) inflexibilis, Carlgren 1928; Stoichactis (Priapus) giganteum, Forskal 1775; Phymanthus (Ixalactis) Ioligo, H and E Ehrenberg, 1834; Bathydactylus valdiviae, Carlgren 1929, and Paraphellia sanzoi, Colabresi 1926, are recorded for the first time from the Indian seas. Besides the identified 40 species, the material collected presumably contain a large number of species, some of which may be new to science.

The actiniarian sea anemone fauna of India was so far known from few places viz. Port Canning (W.Bengal), Chilka lake (Orissa); Adyar backwaters and Krusadai (Tamil Nadu); Cochin backwaters and Ashtamudi lake (Kerala);, Bombay and part of Maharashtra, Goa and northern Karnataka along the Central West coast of India. In last two decades, two new species, namely Edwardsia jonesii, Seshaiya and Cutress 1969 and Paracondylactis sagarensis, Bhattacharya 1979, from Porto Novo along Tamilnadu coast and Sagar Island (Sunderbans) along the West Bengal coast, respectively, have been reported.

Besides a number of unidentified forms from the areas explored so far, 3 sea anemones viz. Bunodeopsis, Cribrinopsis and Aiptasia (Table 1), reported by earlier workers, have still remained undescribed upto species level. Similarly, the status of Boloceractis gopalayi as an independent species than Boloceroides mc murrichi has been seriously doubted (Uchida and Muramatsu, 1958).

Ecological Features

All the 40 species reported here (Table 2) represent diverse ecosystems, like, fringing mangroves, estuaries, coastal lagoon, rocky foreshores, sandy and muddy intertidal area, continental and oceanic islands, neritic and oceanic waters upto abyssal depths. Earlier reports by Annandale (op. cit), Panikkar (op. cit) and Parulekar (op. cit) dealt with only the intertidal and swimming species, majority of which were from brackish water - lagoon and backwaters - and the littoral region of open sea.

In the present report, eight species, inhabiting subtidal regions from shallow to abyssal depths, have been documented. These are Halcampa capensis: 120-300 m; Alicia sansibarensis: 20-40 m; Glyphoperidium bursa: upto 500 m; Glyphostylum calyx: 100-180 m; Bunodactis nicobarica: upto 200 m; Parabunodactis inflexibilis upto 450 m; Bathydactylus valdiviae: 4000 m and Paraphellia sanzoi: upto 40 m depth.

Similarly, some of the species, like Stephensonactis ornata, Phytococopsis ramunii, Bunodeopsis sp., Nevadne glauca, Anemonia indicus, Bunodosoma granulifera, Anthopleura panikkarii, Cribriopsis robertii and Stoichactis giganteum, which earlier were reported only from the intertidal biotopes, were found to be distributed even in the shallow depths of the subtidal region.

An analysis of the habit and substrate preference, as deduced from Table 2, reveals that 20 species were of burrowing type; 18 species comprised of adhesive and attached forms with 2 amphibious species, exhibiting partly sedentary and partly swimming habit. While majority of the adhesive and attached forms were observed to be restricted to intertidal region, especially at mid littoral and lower littoral tide levels, the burrowing forms were mostly confined to lower littoral and subtidal waters and subtidal waters. Most of the burrowing forms, prefer silty sand substratum, except, Edwardsia jonesii, Metapeachia tropica, Alicia sansibarensis, Actinogeton sultana and Bunodactis nicobarica, which prefer fine to medium grain sand. Further, almost all the burrowing species, comprise of solitary forms.

The substrate preference in adhesive and attached species was varied and diverse (Table 2). While few of the species were found attached, mostly singly or rarely in groups, to rock - crevices and fissures - and boulder, majority of the species display substrate preference for living objects. Two types of associations, namely commensalism and symbiosis, were discernible.

Majority of the attached species, occur in group and in species of genus Anthopleura and Cribriopsis, distinct tendencies of colony formation are evident. Similarly, most of the species, belonging to family Actiniidae and Stoichactiidae, harbour a number of symbiotic species, both algae and invertebrates. Many of the species live as commensal with diverse partners like, zoantharians: Anthopleura pacifica and Cribriopsis robertii; corals: Bunodactis nicobarica, Stoichactis giganteum and Phymanthus loligo; bivalve and gastropod molluscs: Anemonia indicus, Anthopleura asiatica, A. panikkarii, Glyphoperidium bursa Cribriopsis sp. and Aiptasiomorpha luciae; bivalve and barnacles:

Anthopleura panikkarii Cribrinopsis robertii and Acontiophorum bombayensis; gastropod and hermit crab: Aiptasia sp. and Neoaiptasia commensali; sessile and stalked cirripedes; Diadumene schilleriana, Commensallic association with polychaete worms, crabs and fishes is also evident in sea anemones viz. Anemonia indicus, Paracondylactis indicus, P. sagarensis, Stoichactis giganteum and Phymanthus loligo.

All the commensal species, either adhesive or swimming forms, show a preferential behaviour pattern with its association. The sea anemones, essentially non-locomotory, its association with other animals is of more benefit to actiniarian. Most sea anemones pass their life attached to host object and take as food organisms or detritus coming within reach of its tentacles. Ecological records show that most sea anemones occupy preferred location in particular ecological situation. There is now a great deal of evidence that actiniarian sea anemones possess capacities for recognising specific chemical factors associated not only with its commensal or symbiont, but also those occurring in its surroundings (Ross op. cit). This particular aspect has opened up various opportunities for using sea anemones, not only as sentinel organisms but as a promising source of bioactivity.

Literature Cited

- ANNANDALE, N. 1907. The fauna of brackish ponds at Port Canning, Lower Bengal. P. 1. Records Indian Museum, 1(4): 46-74.
- ANNANDALE, N. 1915. Fauna of the Chilka lake. The Coelenterates. Mem. Indian Museum V: 21-55.
- CARLGREN, O. 1949. A survey of Ptychodactiaria, Coralliomorpha and Actiniaria, pp. 121 (Almqvist & Wiksell, Stockholm).
- PANIKKAR, N.K. 1936. The structure, bionomics and systematic position of two new brackish water actiniaria from Madras. Proc. Zool. Soc. London, 106: 39-52.
- PANIKKAR, N.K. 1937. A study of the actinian, Phytocoetes gangeticus Zool. Jahb. Abt. Anat. 3: 62-71.
- PANIKKAR, N.K. 1937a. The morphology and systematic relationship of a new boloceroidean from brackish water near Madras. Proc. Indian Acad. Sci. 5(2) : 33-41.
- PANIKKAR, N.K. 1937b. Notes on Nevadne galuca. Records Indian Museum, XXXIX, P.IV.
- PANIKKAR, N.K. 1939. Studies on the brackish water anemone, Pelocoetes exul, Annandale and on a new marine species from Madras. Proc. Zool. Soc. London, (B), 108: 4-7.
- PARULEKAR, A.H. 1966. Cnidae in the actinians of Maharashtra. J. biol. Sci., 9(1 & 2): 36-42.
- PARULEKAR, ARUN. 1967. Two new species of sea anemones (Actiniaria) from Maharashtra. J. Bombay nat. Hist. Soc., 64(3): 524-529.
- PARULEKAR, ARUN. 1968. Sea anemones (Actiniaria) of Bombay J. Bombay nat. Hist. Soc. 65(1): 138-147.
- PARULEKAR, ARUN. 1969a. Neoaiptasia commensali gen et sp nov: An actiniarian commensal of hermit crab. J. Bombay nat. Hist. Soc. 66(1): 57-62.
- PARULEKAR, ARUN. 1969b. On a new species of sea anemone from Maharashtra, India. J. Bombay nat. Hist. Soc. 66(3), 590-595.
- PARULEKAR, A.H. 1971. A new sea anemone, Cribrinopsis robertii (Endomyaria: Actiniidae) from Maharashtra and Goa coast J. Bombay

nat. Hist. Soc. 68(1): 291-295.

ROSS, D.M. 1967. Behavioural and ecological relationships between sea anemones and other invertebrates. Oceanogr. Mar. Biol. Ann. Rev., 5: 291-316.

UCHIDA, T., AND MURAMATSU, S. 1958. Notes on some Japanese sea anemones. J. Fac. Sci. Hokkaido Univ. Series VI (Zoology), 14(1): 111-119.

Table 1. Actiniarian Sea Anemones of India

Genus & Species	Family
1. <u>Edwardsia tinctrix</u> , Annandale 1915	... Edwardsiidae, Andres, 1880
2. <u>E. jonesii</u> , Seshaiya & Curtress 1969	... " "
3. <u>Metapeachia tropica</u> , Panikkar 1938	... Haloclavidae, Verrill 1899
4. <u>Halcampa capensis</u> , Carlgren 1938	... Halcampidae, Andreas 1883
5. <u>Mena limnicola</u> , Annandale 1915	... " "
6. <u>Pelocoetes exul</u> , Annandale 1915	... Haliactidae, nov. nom
7. <u>P. minima</u> , Panikkar 1939	... " "
8. <u>Stephensonactis ornata</u> , Panikkar 1936	... " "
9. <u>Phytocoetes gangeticus</u> , Annandale 1915	... " "
10. <u>Phytocoetopsis ramunni</u> , Panikkar 1936	... " "
11. <u>Boloceroides mc murrichi</u> Kwiteriewski	1898 Boloceroidae, Carlgren 1924
12. <u>Boloceractis gopalayi</u> , Panikkar 1937	... " "
13. <u>Bunodeopsis</u> sp.	... " "
14. <u>Nevadne glauca</u> , Annandale 1915	... Nevadneidae, Carlgren 1925
15. <u>Alicia sansibarensis</u> , Carlgren 1900	... Aliciidae (Duerden 1897)
16. <u>Anemonia indicus</u> , Parulekar 1968	... Actiniidae (Gosse 1858)
17. <u>Bunodosoma granulifera</u> , Leseur 1917	... " "
18. <u>Anthopleura midori</u> , Uchida 1958	... " "
19. <u>A. asiatica</u> , Uchida 1958	... " "
20. <u>A. pacifica</u> , Uchida 1958	... " "
21. <u>A. panikkarii</u> , Parulekar 1969	... " "
22. <u>Paracondylactis indicus</u> , Dave 1957	... " "
23. <u>P. sagarensis</u> , Bhattacharya 1979	... " "
24. <u>Glyphoperidium bursa</u> , Roule 1909	... " "
25. <u>Glyphostylum calyx</u> , Roule 1909	... " "
26. <u>Actinogeton sultana</u> , Carlgren 1900	... " "
27. <u>Cribrinopsis robertii</u> , Parulekar 1971	... " "
28. <u>Cribrinopsis</u> sp.	... " "
29. <u>Bunodactis nicobarica</u> , Carlgren 1928	... " "
30. <u>Parabunodactis inflexibilis</u> , Carlgren 1928	" "
31. <u>Stoichactis giganteum</u> , Forskal 1775	... Stoichactiidae Carlgren 1900
32. <u>Phymanthus loligo</u> , H and E Ehrenberg 1834	Phymanthiidae (Andres 1883)
33. <u>Bathydactylus valdiviae</u> , Carlgren 1928	... Actinostoiidae, Carlgren 1938
34. <u>Acontiophorum bombayensis</u> , Parulekar 1968	Acontiophoriidae, Carlgren 1938
35. <u>Paraphellia sanzoi</u> , Colabresi 1926	... Hormathiidae, Carlgren 1925
36. <u>Metridium senile</u> var <u>fimbriatum</u> (Verrill)	Metridiidae (Carlgren 1893)
37. <u>Aiptasia</u> sp.	... Aiptasiidae, Carlgren 1924
38. <u>Neoaiptasia commensali</u> , gen et sp. nov, Parulekar 1969	... " "
39. <u>Aiptasiomorpha luciae</u> , Verrill 1899	... Aiptasiomorphidae (Stephenson 1920)
40. <u>Diadumene schilleriana</u> , Stoliczka 1863	... Diadumeneidae (Stephenson 1920)

Table 2. Ecological Features of the Actinian Sea Anemones of India.

Sl. No.	Species	Habitat	Habit	Distribution
1	2	3	4	5
1.	<u><i>Edwardsia tinctrix</i></u>	Estuarine/marine Intertidal-mid to lower littoral	Burrowing-silty-sand solitary form	Chilka lake, Madras, Bombay, Malvan, Goa
2.	<u><i>E. Jonesii</i></u>	Estuarine-Intertidal-Littoral	Burrowing-Fine to sand-Lower Littoral	Silty sand-solidary Porto Novo (Tamil Nadu)
3.	<u><i>Metapeachia tropica</i></u>	Marine-Intertidal-Lower Littoral	Burrowing-Fine sand-Solitary form	Krusadai, Bombay, Malvan, Goa.
4.	<u><i>Halocampa capensis</i></u>	Marine-Subtidal (100-300 m)	Burrowing-Silty sand Solitary form	Northwest coast of India.
5.	<u><i>Mena limnicola</i></u>	Estuarine-Intertidal-Lower littoral	Burrowing-silty sand growing in groups	Chilka Lake, Balram-gadhi (Orissa)
6.	<u><i>Pelocoetes exul</i></u>	Estuarine-Intertidal-lower littoral	Burrowing/Attached-silty sand/concrete structures-solidary but at times found aggregating on jetty pilings	Gangetic delta, Madras Cochin backwaters, Goa estuaries, Malvan, Gulf of Cambay.
7.	<u><i>P. minima</i></u>	Estuarine-Intertidal-mid to lower littoral	Burrowing-silty sand	Nadras, Sagar Island
8.	<u><i>Stephensonactis ornata</i></u>	Estuarine-Intertidal/subtidal-lower littoral/ upto 15 m depth	Burrowing-Silty sand-solidary	East & West coast of India.
9.	<u><i>Phytocoetes gangeticus</i></u>	Estuarine-Intertidal	Burrowing/Attached-silty sand/concrete structures-solidary to aggregation	East & West coast of India.

1	2	3	4	5
10.	<u>Phytocoetops ramunni</u>	Estuarine-Intertidal/subtidal	Burrowing-silty sand- Madras solitary	
11.	<u>Boloceroides mc murrichi</u>	Estuarine/marine-pelagic	Sedentary/swimming-Hard substratum/ Pelagic-solidary	Central west coast of India
12.	<u>Boloceractis gopalayi</u>	Estuarine-pelagic	Sedentary/swimming-Hard substratum/ Pelagic-solidary	Madras, Cochin back-waters, Asthamudi Lake
13.	<u>Bunodeopsis sp.</u>	Estuarine-Intertidal/subtidal	Burrowing-silty sand	Madras
14.	<u>Nevadne glauca</u>	Estuarine-Intertidal/subtidal	Burrowing-silty sand/ Chilka Lake, Goa	
15.	<u>Alicia sansibarensis</u>	Marine-subtidal (20-40 m)	Burrowing-fine sand-solidary form	Southwest & Central West coast of India
16.	<u>Anemonia indicus</u>	Marine-Lower littoral/subtidal	Adhesive form-attached to bivalve shells, Goa, North Karnataka & polychaete tubes of Gulf of Kutch.	Coast of Maharashtra
			<u>Onuphis</u> sp. and concrete structures-solidary to gregarious	
17.	<u>Bunodosoma granulifera</u>	Marine-Lower littoral/subtidal	Adhesive form-found attached in crevices/ fissures of rock	Bombay
18.	<u>Anthopleura midori</u>	Marine-mid littoral to lower littoral	Adhesive form-attached to rock or under-surface of boulders & stones-gregarious in habit	Maharashtra, Goa and north Karnataka coast.
19.	<u>A.asiatica</u>	Marine-Lower littoral	Adhesive form-firmly attached, singly to rock or oyster shell in rock pools	Bombay, Malvan, Anjiv Island

	1	2	3	4	5
20.	<u>A. pacifica</u>	Marine-mid-littoral to lower littoral	Adhesive form-found in shallow rock pools -solitary	Bombay, Malvan, Goa.	
21.	<u>A. Panikkarii</u>	Marine-lower littoral to subtidal	Adhesive form-attach-ed in large numbers to rocky slopes also crevices alongwith barnacles and oysters Nicobar).	Maharashtra, Goa, north Karnataka coast, Vizhinjam (Kerala), Port Blair (Andaman-	
22.	<u>Paracondylac indicus</u>	Marine/Estuarine-intertidal lower	Burrowing-silty sand/ slushy mud/silty clay delta. -solitary	Bombay, Sikka, Gangetic	
23.	<u>P. sagrensis</u>	Estuarine-inter-tidal	Burrowing-slushy mud (Sunder bans). -solitary	Sagar Island (Sunder	
24.	<u>Glyphoperidium hursa</u>	Marine-subtidal (upto 500 m depth)	Adhesive form-attach-ed singly or in group to molluscan shell debris	Off southwest coast of India	
25.	<u>Glyphostylum calyx</u>	Marine-subtidal (100-180 m depth)	Adhesive form-colonial-attached to shell and coral debris	Off southwest coast of India & Lakshadweep sea	
26.	<u>Actinogenetosultana</u>	Marine-Intertidal-Lower littoral	Burrowing-fine sand-solitary	Cuffe Parade (Bombay).	
27.	<u>Cribrinopsis robertii</u>	Marine-Lower littoral	Adhesive form-attach-ed to oyster (<i>C. cucullata</i>) and barnacle shells. Also found in association with live corals	Maharashtra & Goa coast Mandapam, Tuticorin.	
28.	<u>Cribrinopsis</u> sp.	Marine-intertidal-mid littoral to lower littoral	Adhesive form-attach-ed to rock and oyster shells-solitary to group	Bombay.	
29.	<u>Bunoedactis nicobarica</u>	Marine-subtidal (upto 200 m depth)	Burrowing-fine to medium sand/coral debris-solitary	Andaman Sea.	

1	2	3	4	5
30.	<u>Parabunodactis</u> <u>inflexibilis</u>	Marine-subtidal (upto 450 m depth) to fine sand-solitary	Burrowing-silty sand	Andaman Sea.
31.	<u>Stoichactis</u> <u>giganteum</u>	Marine-Intertidal- Lower littoral to subtidal	Adhesive form-found attached singly to live corals	Gulf of Kutch & Gulf of Mannar
32.	<u>Phymanthus</u> <u>loligo</u>	Marine-Intertidal- mid to lower littoral	Adhesive form at times found burrowing Mannar.	Gulf of Kutch & Gulf of Mannar
33.	<u>Bathydactylus</u> <u>valdiviae</u>	Marine-subtidal- abyssal depth of 4000 m	Burrowing-silty clay- solitary	Off west coast of India (previously reported from the northern sector of the Southern ocean).
34.	<u>Acontiophorum</u> <u>bombayensis</u>	Marine-Subtidal- upper to mid littoral	Adhesive form-found attached singly or in group to boring bivalve, <u>Martesia</u> sp. and barnacle shells	Bombay, Malvan, Karwar.
35.	<u>Paraphellia</u> <u>sanzoi</u>	Marine-subtidal- upto 40 m depth	Burrowing-silty clay-	Gulf of Kutch
36.	<u>Metridium</u> <u>senile</u> var <u>fimbriatum</u>	Marine-Intertidal- lower littoral	Adhesive form attach- ed in groups inside rock crevices, fissu- res and to concrete structures	West & East coast of India, Port Blair
37.	<u>Aiptasia</u> sp.	Marine-Intertidal- Lower littoral at times mid littoral	Adhesive form-attach- ed singly or in pair to gastropod shells inhabited by hermit crabs	Bombay

1	2	3	4	5
38.	<u><i>Neoaiptasia commensali</i></u>	Marine-Intertidal- Lower littoral	Adhesive form-attach- ed singly or even upto 4 individuals to 73° 28'E) a gastropod shell viz <u><i>Babylonia spirata</i></u> in- habited by hermit crab with coinhabitant, <u><i>Ergoea walshii</i></u> and polynoid <u><i>Gatyania deludens</i></u>	Padmagad (16° 03'N,
39.	<u><i>Aiptasiomorpha luciae</i></u>	Marine/Estuarine- Intertidal-Lower littoral	Adhesive form-attach- ed generally in pair to gastropod shell <u><i>Nassarius ornata</i></u> , rarely <u><i>Cerithidium fluviatilis</i></u>	Bombay, Gulf of Kutch, Central & Southwest coast of India.
40.	<u><i>Diadumene schilleriana</i></u>	Marine-Pelagic/ mid littoral	Adhesive form-attach- ed to drifting wood in association with stalked barnacle, <u><i>Lepas</i></u> sp. Also found attached to barnacle shells	West coast of India, Tuticorin, Mandapam, Chandipore, Digha.