

Author version: J. Mar. Biol. Assoc. UK, vol.90(6); 2010; 1241-1247

What do barnacle larvae feed on? Implications in biofouling ecology

CHETAN A. GAONKAR AND ARGHA CHANDRASHEKAR ANIL

National Institute of Oceanography (Council of Scientific and Industrial Research), Dona-Paula, Goa - 403
004, India

*Barnacles are one of the dominant macrofouling organisms found in the intertidal region throughout the world. Among the different species of barnacles *Balanus amphitrite* (= *Amphibalanus amphitrite*) is a favoured candidate organism used in experimental studies. Larval development in this barnacle includes planktotrophic naupliar stages followed by pre-settling cyprid instar. Studies have shown that availability of food during naupliar development is of critical importance to successful metamorphosis of the cypris larva. Traditionally barnacle larvae are raised in the laboratory providing mono-algal cultures of diatoms as food organisms. Such a luxury is not a reality in the wild. Observations to quantify the food available for the nauplii deliberated by monitoring the fecal pellets egested by freshly captured larvae from a tropical estuarine environment (Dona Paula bay, Goa, West Coast of India) influenced by monsoon and characteristic temporal variations in the phytoplankton abundance and diversity indicated that the percentage of defecating larvae (an indicator of food consumed) was comparatively higher during the pre-monsoon season. Generally this season is characterized by lower chlorophyll a concentration. However, the average number of fecal pellets defecated by a larva remained constant irrespective of the season. Earlier work in the study area depicts temporal changes in phytoplankton community structure; diatoms dominate during the post-monsoon season whereas dinoflagellates dominate during the pre-monsoon season. These observations indicate a possible shift in the food available for the larvae. As the fecal pellets did not always have remnants of diatom frustules, it is possible to say that the larvae survived on food material other than diatoms. Settlement of barnacles on panels of aluminium in the vicinity was monitored round the year and is peaked during pre-monsoon season of the year. It is thus possible to infer successful larval development and metamorphosis in this barnacle species on varying forms of food.*

Keywords: Barnacle larvae, *Balanus amphitrite* (= *Amphibalanus amphitrite*), fecal pellets, phytoplankton, diatoms, dispersion, settlement and recruitment

Corresponding author:

A. C. Anil

E-mail: acanil@nio.org

INTRODUCTION

Barnacles are an important intertidal organisms and one of the dominant components of the macrofouling community. Larval development of this organism includes planktonic naupliar stages followed by a pre-settling cyprid instar. Successful recruitment of barnacles is governed by the naupliar developmental history. Their sessile nature and planktotrophic larval development give them the dual capability of dispersion either as a fouling organism or through their sojourn inside the dark ballast tanks. There has been a growing body of evidence showing that decreases in larval energetic reserves can strongly impose upon post-metamorphic performance of the larvae (Jarrett, 1997; Pechenik *et al.*, 1998; Miron *et al.*, 2000; Anil *et al.*, 2001). Another facet to larval dispersion is their capability to survive such dispersion as they are dependent on the availability of food. Experiments carried out earlier have shown the starvation capability of larvae of the barnacle *Balanus amphitrite* (= *Amphibalanus amphitrite*) is influenced by temperature and the type of food available (Anil *et al.*, 1995; Desai & Anil, 2000; 2004).

Barnacle *Balanus amphitrite* (= *Amphibalanus amphitrite*) is a favoured candidate organism in antifouling assay and its larvae are raised with a rich supply of mono-algal food coupled with optimum incubating conditions whereas larvae in the natural habitat has to survive wide variations in environmental parameters as well as the availability of food. Information on the food consumed by the barnacle nauplii in the wild is very much lacking in spite of its importance in larval energetics, dispersion, settlement and recruitment (Turner *et al.*, 2001; Vargas *et al.*, 2006). Technical difficulties when working with these small nauplii may also explain the scarcity of information on their feeding habits. Work on laboratory rearing of nauplii, fed on unialgal cultures demonstrates that nauplii feed efficiently on cells over a different size range (Vargas *et al.*, 2006). However, our knowledge is derived from incubation experiments, most of them using laboratory cultured phytoplankton as food (Qiu & Qian, 1997; Anil *et al.*, 2001; Desai & Anil, 2004; Desai *et al.*, 2006; Nasrolahi *et al.*, 2007).

Present study deals with the food consumed by the barnacle nauplii in a natural environment, assessed by the quantification of fecal pellets egested by a freshly captured individual, which were reckoned to represent a measure of feeding status at the time of capture. The method provides information on *in situ* feeding rates or conditions immediately prior to collection. Until now, this

method has not been used with nauplii of barnacles collected from the field mostly due to technical difficulties like handling these small size nauplii. Studies related to the use of this method in case of crustaceans are mostly available with adult copepods which are collected from the field as well as from laboratory reared copepods fed with different mono-algal cultures (Honjo & Roman, 1978; Morales, 1987; Urban-Rich *et al.*, 1998; Fleddum *et al.*, 2001; Frangoulis *et al.*, 2001; Schnetzer & Steinberg, 2002; Bathmann & Liebezeit, 2008; Ploug *et al.*, 2008). The aim of the present study was to assess the food consumed by the barnacle nauplii in the natural environment by quantifying the larval defecation rates and their variations over a temporal scale and also to assess the contents of the fecal pellets using scanning electron microscopy. Further, the consequences of changes in the food availability as indexed through fecal pellets egested are discussed with respect to settlement and recruitment variations of the barnacle *Balanus amphitrite* (= *Amphibalanus amphitrite*) in a tropical estuarine environment influenced by monsoons.

MATERIALS AND METHODS

In order to ascertain the food consumed by the barnacle nauplii in the natural environment, a study on temporal variations of larval defecation and its contents was carried out over a period of two years (June 2005- May 2007) in a tropical estuarine environment influenced by monsoons (Dona Paula bay, Goa, West Coast of India) which is known for its characteristic temporal variations in the phytoplankton abundance and diversity. Nauplii collected from the Dona Paula bay were incubated in the laboratory for the evaluation of food consumed.

Larval samples were collected monthly using 100 μ m mesh net towed horizontally with the help of boat. They were immediately brought to the laboratory after collection in live conditions and IV-VI instar nauplii (n=24) were isolated individually in to a multi-wells (24 well) containing 0.22 μ m filtered seawater. The number of defecated pellets by a larva after 12 hours of incubation at room temperature and the proportion of the larvae defecating was quantified. After the observations, pellets were preserved in Ethanol for Scanning Electron Microscope (SEM) photography to find the signatures of food. Ethanol preserved samples were filtered on to 0.22 μ m polycarbonate membrane filters, rinsed with deionised water to remove salt and ethanol and then air dried. Filters were then placed on SEM stubs and sputtered with gold/palladium for 3 minutes and the photographs were taken at different magnifications.

Observations were also carried out with the laboratory hatched larvae by providing phytoplankton cultures of *Chaetoceros calcitrans* and *Skeletonema costatum* as food and the pellets defecated by the larvae were collected and processed for SEM photography for the evidence of ingested food and to compare them with the pellets defecated by field collected larvae.

Settlement and recruitment observations of the barnacles in the vicinity of the study area were also carried out on a monthly basis from June 2005 to May 2007 by immersing aluminium panels. Every month new set of panels were immersed in triplicate and observations were taken from six marked quadrats within the panels of size 25cm² each. Settled and recruited barnacles on the panels were counted at the end of a month with the help of a magnifying glass. 0-2mm size classes were considered as settlers and above 2mm size classes were considered as recruits. Settlement and recruitment of the barnacles is expressed in terms of numbers per decimetre square (No./dm²).

Data analysis

Percentage of defecating larvae collected during different sampling months was subjected to two-way analysis of variance (ANOVA) (Sokal and Rohlf, 1981) to evaluate the variance between different sampling months. Settlement and recruitment of the barnacles monitored during different sampling months was also subjected to two-way analysis of variance. Data was log transformed before subjecting to ANOVA to ensure normality of means and homogeneity of variance.

RESULTS

Fecal pellets

Average number of fecal pellets defecated by a larva remained more or less constant throughout the period of observation (July 2005-May 2007) (Figure 1). The % of larvae defecating was comparatively higher during February-May (pre-monsoon) period of the year and the highest was observed during March 2006 (Figure 2). Analysis of variance also indicated a significant variation among different sampling months (ANOVA: $p \leq 0.0005$).

SEM photographs of the pellets collected during October-January (post-monsoon) period indicated the presence of pennate diatoms. The pellets produced during February-May (pre-monsoon) period did not show any diatom frustules. They were mostly in the form of particulate matter or in some cases unidentifiable (Figure 3 b & c).

Laboratory experiments

The pellets produced from the laboratory raised larvae with centric diatoms as food (*Chaetoceros calcitrans* and *Skeletonema costatum*) indicated the presence of diatom frustules in the remnants of food provided (Figure 3 a).

Settlement and recruitment of barnacles

Settlement and recruitment of the barnacles varied throughout the sampling period. Settlement was less during the monsoon season (June-September). Peak in settlement was observed during February-06 followed by November-06 (Figure 4). Analysis of variance also indicated a significant variation in barnacle settlement among different months. (ANOVA: $p \leq 0.05$). Recruitment of barnacles was also less during the monsoon season except for August month and it peaked during pre-monsoon season (February-May) of the year (Figure 5). Analysis of variance also indicated a significant variation in barnacle recruitment among different sampling months (ANOVA: $p \leq 0.05$).

DISCUSSION

The mode of nutrition in larval forms plays an important role in their capability to survive and disperse. In case of larval forms, which have a feeding requirement essential for their growth and development, the rate of their survival is dependent on food availability, life span and their capability to tolerate starvation. Barnacles have both feeding and non feeding stages in their larval life span. The chances of survival of non-feeding stage larvae (cyprid stage) depend upon the stored energy of the larvae or the naupliar developmental history. On the other hand survival of the feeding stage larvae depends upon the food available for the larva in the environment or their capability to tolerate starvation (Desai & Anil, 2004).

As the success of population of barnacles in any given environment is determined by the events during larval life period it is important to ascertain the type of food the nauplii consume. Such an effort made in this study through larval defecation experiments showed that the average number of fecal pellets defecated by a larva remained constant in spite of possible variations in the type of phytoplankton available in the environment. Results also indicated that the percentage of defecating larvae (an indicator of food consumed) was comparatively higher during the pre-monsoon season (Figure 2). However, the average number of fecal pellets defecated by a larva remained constant irrespective of the season (Figure 1).

The signatures of food found during the pre-monsoon season in the fecal pellets were elusive, indicating altered food consumption (i.e. other than diatoms) as compared to the post-monsoon months. SEM photographs of the pellets which were collected during post-monsoon season indicated the presence of pennate diatoms not among the common food organisms employed in the laboratory rearing. The pellets produced during the pre-monsoon season did not have any diatom remains whereas the pellets produced during the post-monsoon season showed some frustules of diatoms. This indicates that during the pre-monsoon season the larvae survived on food material other than diatoms. It has been reported that the phytoplankton community in the study area changes with the season and dinoflagellates form a substantial part (~30%) during the pre-monsoon season (Patil & Anil, 2008).

Earlier studies have also indicated that the food preferred by barnacle larvae in the wild includes small flagellates at relatively high rates along with diatoms (Turner *et al.*, 2001). In accordance with their feeding mechanisms and body size, barnacle nauplii were also able to feed on autotrophic picoplankton (<5 μm) and did not consume the largest phytoplankton cells (Vargas *et al.*, 2006). Experiments by Nejstgaard *et al.*, (2007) also indicated that they can feed on *Phaeocystis* but was pointed out by authors that this needs to be treated with caution and with further investigation.

Microcosm rearing of the larvae in this tropical bay by Desai & Anil (2002) indicated that the naupliar duration of larvae is prolonged by only two days as compared to those raised in the laboratory. It was also observed that chlorophyll *a* content of the sea water in the bay during the experiment was in the range of 0.3-4.0 $\mu\text{g L}^{-1}$ (Desai & Anil, 2002). Even at the lowest concentration of food provided in the laboratory rearing, the chlorophyll *a* content is 15 times higher than the maximum values observed in the field. This shows the capability of larvae to survive in the

field in spite of observed low chlorophyll *a* concentration in the water column, though the larvae raised in the microcosm experienced nutritional stress which was shown by lower ratios of RNA:DNA (Desai & Anil, 2002).

This study further indicates their capability to feed on food materials other than diatoms. As the fecal pellets did not always have remnants of diatom frustules, it is possible to say that the larvae survived on food material other than diatoms. This indicates a shift in the food available for the larvae in the environment and such shifts can influence development and metamorphosis capability of the barnacles. Settlement and recruitment of the barnacles in the vicinity indicated a seasonal variation in the rate of settlement and recruitment. Settlement and recruitment during the monsoon season was inconsistent, whereas during the pre-monsoon season, settlement and recruitment occurred throughout the season. The consistency in settlement and recruitment observed during the pre-monsoon season coincides with the higher percentage of defecating larvae and the absence of diatom frustules in the fecal pellet.

Success of larval development in the natural environment in spite of altered food availability can have greater implication from the point of survival and translocation of larvae to different habitats through dispersion and other physical processes. Settlement behaviour of the larvae also has the potential to affect larval dispersion. 'Desperation' of the larvae is one source of variation in larval settlement behaviour. It has been proposed that as larvae grow old, they become less discriminatory in their selection of settlement substrate (Knight-Jones, 1953; Toonen & Pawlik 2001). A number of workers have suggested that the maximum planktonic period of non-feeding larvae is determined by the energetic reserves (Lucas *et al.*, 1979; Anil & Kurian, 1996; Anil *et al.*, 2001; Thiyagarajan *et al.*, 2002; Desai & Anil, 2004 and references therein). Other source of variation in settlement behaviour may be the size of larvae. There are some evidences that larval size does influence larval life span in some invertebrates, with larger larvae remaining active for longer than smaller larvae (Marshall *et al.*, 2003; Marshall & Keough, 2003; Isomura & Nishihira, 2001). However, these issues have not been addressed in case of barnacles from the point of settlement of larvae or dispersal of larvae to different bioregions. If larval size affects larval settlement behaviour, then variation in larval size could also indirectly affect the dispersal potential. Larger larvae will have greater nutritional reserves than smaller larvae and they can swim for longer duration than smaller larvae and would be less desperate to settle. In this context, it is possible for the larger larvae to remain planktonic for longer duration, hence a greater dispersal potential.

Most of the limited information on feeding characteristics of the barnacle nauplii comes from experimental or rearing studies where laboratory cultured phytoplankton are offered as food. Also, primarily herbivorous feeding by barnacle nauplii is in agreement with numerous observations using laboratory cultured diatoms as food, but there is in fact very limited information on feeding by barnacle nauplii on natural plankton assemblages. The attempt made here in this study indicates the capability of nauplii to feed on food materials other than diatoms and also shows that the rate of larval defecation remained constant throughout the year in spite of reported seasonal variations in phytoplankton community in the water column by Patil & Anil (2008). In view of this it is important to consider the consequences of such changes to the availability of food for the nauplii when one postulates the risk of translocation of organisms with such planktotrophic naupliar developmental pathways.

Quantitative analysis of fecal pellet production and its content would also provide insights about the preference of larval food in the wild. This is a difficult task to be accomplished and requires molecular techniques to identify the signatures of food organisms consumed. Nevertheless, this observation shows the need to study in detail the larval food preference in the natural environment.

ACKNOWLEDGEMENTS

We are grateful to Dr. S.R. Shetye, Director, National Institute of Oceanography for his support and constant encouragements. The Ballast Water Management Programme, India (Directorate General of Shipping, Government of India) and the project on Molecular Characterization of Microbial and Invertebrate Diversity of Indian West Coast (Department of Biotechnology, Government of India) supported this work. We thank Dr. S.S. Sawant and Mr. K. Venkat and other members of the project for their ever-willing help. The first author acknowledges the Research Fellowship provided by the Council of Scientific and Industrial Research (CSIR), India. This is a NIO contribution #####.

REFERENCES

- Anil A.C., Chiba K., Okamoto K. and Kurokura H.** (1995) Influence of temperature and salinity on the larval development of *Balanus amphitrite*: Implications in the fouling ecology. *Marine Ecology Progress Series*, 118, 159-166.
- Anil A.C. and Kurian J.** (1996) Influence of food concentration, temperature and salinity on the larval development of *Balanus amphitrite*. *Marine Biology*, 127, 115-124.
- Anil A.C., Desai D. and Khandeparker L.** (2001) Larval development and metamorphosis in *Balanus amphitrite* Darwin (Cirripedia: Thoracica): significance of food concentration, temperature and nucleic acids. *Journal of Experimental Marine Biology and Ecology*, 263(2), 125-141.
- Bathmann U. and Liebezeit G.** (2008) Chlorophyll in copepod fecal pellets: changes in pellet numbers and pigment content during a declining Baltic spring bloom. *Marine Ecology*, 7(1), 59-73.
- Desai D.V. and Anil A.C.** (2000) Influence of temperature on the starvation threshold of nauplii of the barnacle *Balanus amphitrite* (Cirripedia: Thoracica). *Indian Journal of Marine Science*, 29, 69-72.
- Desai D. V. and Anil A. C.** (2002) Comparison of nutritional status of field and laboratory reared *Balanus amphitrite* Darwin (Cirripedia: Thoracica) larvae and implication of starvation. *Journal of Experimental Marine Biology and Ecology*, 280, 117-134.
- Desai D.V. and Anil A.C.** (2004) The impact of food type, temperature and starvation on larval development of *Balanus amphitrite* Darwin (Cirripedia: Thoracica). *Journal of Experimental Marine Biology and Ecology*, 306, 113-137.
- Desai D.V., Khandeparker L. and Shirayama Y.** (2006) Larval development and metamorphosis of *Balanus albicostatus* (Cirripedia: Thoracica): implications of temperature, food concentration and energetics. *Journal of the Marine Biological Association of the United Kingdom*, 86 (2), 335-343.
- Fleddum A., Kaartvedt S. and Ellertsen B.** (2001) Distribution and feeding of the carnivorous copepod *Paraeuchaeta norvegica* in habitats of shallow prey assemblages and midnight sun. *Marine Biology*, 139, 719-726.
- Frangoulis C., Belkhiria S., Goffart A. and Hecq J.H.** (2001) Dynamics of copepod fecal pellets in relation to a *Phaeocystis* dominated phytoplankton bloom: Characteristics, Production and Flux. *Journal of Plankton Research*, 23(1), 75-88.

- Honjo S. and Roman M.R.** (1978) Marine copepod fecal pellets: production, preservation and sedimentation. *Journal of Marine Research*, 36, 45-56.
- Isomura N. and Nishihira M.** (2001) Size variation of planulae and its effect on the lifetime of planulae in three pocilloporid corals. *Coral Reefs*, 20, 309-315.
- Jarrett J.N.** (1997) Temporal variation in substrate specificity of *Semibalanus balanoides* (Linnaeus) cyprids. *Journal of Experimental Marine Biology and Ecology*, 211, 103-114.
- Knight-Jones E.W.** (1953) Laboratory experiments on gregariousness during settling in *Balanus balanoides* and other barnacles. *Journal of Experimental Biology*, 30, 584-599.
- Lucas M.I., Walker G., Holland D.L. and Crisp D.J.** (1979) An energy-budget for the free-swimming and metamorphosing larvae of *Balanus balanoides* (Crustacea: cirrepedia). *Marine Biology*, 55, 221-229.
- Marshall D.J. and Keough M.J.** (2003) Variation in the dispersal potential of non-feeding invertebrate larvae: the desperate larva hypothesis and larval size. *Marine Ecology Progress Series*, 255, 145-153.
- Marshall D.J., Pechenik J.A. and Keough M.J.** (2003) Larval activity levels and delayed metamorphosis affect post-larval performance in the colonial ascidian *Diplosoma listerianum*. *Marine Ecology Progress Series*, 246, 153-162.
- Miron G., Walters L.J., Tremblay R. and Bourget E.** (2000) Physiological condition and barnacle larval behavior: a preliminary look at the relationship between TAG/DNA ratio and larval substratum exploration in *Balanus amphitrite*. *Marine Ecology Progress Series*, 198, 303-310.
- Morales C.E.** (1987) Carbon and nitrogen content of copepod fecal pellets: effect of food concentration and feeding behaviour. *Marine Ecology Progress Series*, 36, 107-114.
- Nasrolahi A., Sari A., Saifabadi S. and Malek M.** (2007) Effects of algal diet on larval survival and growth of the barnacle *Amphibalanus (=Balanus) improvisus*. *Journal of the Marine Biological Association of the United Kingdom*, 87, 1227-1233.
- Nejstgaard J.C., Tang K.W., Steinke M., Dutz J., Koski M., Antajan E. and Long J.D.** (2007) Zooplankton grazing on Phaeocystis: a quantitative review and future challenges. *Biochemistry*, 83, 147-172.
- Patil J.S. and Anil A.C.** (2008) Temporal variation of diatom benthic propagules in a monsoon-influenced tropical estuary. *Continental Shelf Research*, 28, 2404-2416.
- Pechenik J.A., Wendt D.E. and Jarrett J.N.** (1998) Metamorphosis is not a new beginning. *Bioscience*, 48, 901-910.

- Ploug H., Iversen M.H., Koski M. and Buitenhuis E.T.** (2008) Production, oxygen respiration rates and sinking velocity of copepod fecal pellets: direct measurements of ballasting by opal and calcite. *Limnology Oceanography*, 53(2), 469-476.
- Qiu J.W. and Qian P.Y.** (1997) Effects of food availability, larval source and culture method on larval development of *Balanus amphitrite amphitrite* Darwin: implications for experimental design. *Journal of Experimental Marine Biology and Ecology*, 217, 47-61.
- Schnetzer A. and Steinberg D.K.** (2002) Active transport of particulate organic carbon and nitrogen by vertically migrating zooplankton in the Sargasso Sea. *Marine Ecology Progress Series*, 234, 71-84.
- Sokal R.R. and Rohlf F.J.** (1981) In: Biometry, second ed. Freeman WH and Company, San Francisco.
- Thiyagarajan V., Nair K.V.K., Subramoniam T. and Venugopalan V.P.** (2002) Larval settlement behaviour of the barnacle *Balanus reticulatus* in the laboratory. *Journal of the Marine Biological Association of the United Kingdom*, 82(4): 579-582.
- Toonen R.J. and Pawlik J.R.** (2001b) Settlement of the gregarious tube worm *Hydroides dianthus* (Polychaeta: Serpulidae) II: testing the desperate larvae hypothesis. *Marine Ecology Progress Series*, 224, 115-131.
- Turner J.T., Levinsen H., Nielsen T.G. and Hansen B.W.** (2001) Zooplankton feeding ecology: grazing on phytoplankton and predation on protozoans by copepod and barnacle nauplii in Disko Bay, West Greenland. *Marine Ecology Progress Series*, 221, 209-219.
- Urban-Rich J., Hansell D.A. and Roman M.R.** (1998) Analysis of copepod fecal carbon using a high temperature combustion method. *Marine Ecology Progress Series*, 171, 199-208.
- Vargas C.A., Manriquez P.H. and Navarrete S.A.** (2006) Feeding larvae of intertidal invertebrates: assessing their position in pelagic food webs. *Ecology*, 87 (2), 444-457.

LEGENDS TO FIGURE


Fig. 1. Average number of fecal pellets defecated by a larva during different sampling months. * sign indicates the absence of sample collection during that particular month. Error bar indicates the standard deviation from the mean.


Fig. 2. Percentage of larvae defecating during different sampling months. * sign indicates the absence of sample collection during that particular month.


Fig. 3. SEM photographs: (a) fecal pellets produced by the laboratory reared larvae, (b) fecal pellets produced by the field collected larvae (Pre-monsoon, Feb-May), (c) fecal pellet contents from the field collected larvae (Post-monsoon, Oct-Jan).

Fig. 4. Settlement of the barnacle *Balanus amphitrite* (= *Amphibalanus amphitrite*) during different sampling months. Error bar indicates the standard deviation from the mean.


Fig. 5. Recruitment of the barnacle *Balanus amphitrite* (= *Amphibalanus amphitrite*) during different sampling months. Error bar indicates the standard deviation from the mean.

**Fig. 1.**


**Fig. 2.**



a.





b.



c.

Fig.3

**Fig. 4.**

**Fig. 5.**