

## **Non-destructive foraminiferal paleoclimatic proxies: A brief insight**

**Rajeev Saraswat**

Micropaleontology Laboratory, National Institute of Oceanography, Dona Paula, Goa, India

(Email: rsaraswat@nio.org; Phone: 91-832-2450622)

### **Abstract**

The knowledge of past climate can help us to understand imminent climatic changes. Oceans are the vast archives of past climate. Various indirect techniques termed as proxies are used to infer key components of the past climate from the marine sediments. The fossil remains of marine microorganisms, namely foraminifera, are widely used to reconstruct past climate from marine archives, as foraminifera are highly abundant and extensively present in almost all marine realms. Foraminiferal paleoclimatic proxies can be classified in two broad categories, viz. A) destructive (the stable isotopic and trace element composition of foraminifera), those wherein the foraminiferal remains are analyzed to measure its chemical composition and it leads to destruction of the foraminiferal test, and B) non-destructive, wherein changes in the abundance and morphology of foraminifera are used to reconstruct past climate. Here the non-destructive foraminiferal paleoclimatic proxies have been discussed with its recent developments as well as applications. Despite the extensive application of the stable isotopic and trace element composition of foraminifera to infer past climate, non-destructive techniques are still quite helpful to reconstruct past climate as well as to substantiate inferences drawn from destructive techniques.

**Keywords:** Foraminifera, proxy, morphology, paleoclimate, benthic, planktic, species

## **Introduction**

Climate is one of the important factors that affect the fate of mankind on earth. Climate changes can lead to significant cultural shifts, large-scale migration or even demise of civilizations (Polyak and Asmerom, 2001; Haug *et al.*, 2003; Gupta, 2004; Robbins *et al.*, 2013). Therefore precise information of imminent climatic variations is necessary to secure the fate of mankind on earth and for long and short-term policy planning. Continuous efforts are made to develop models to predict future climatic variations. The training and validation of climate prediction models requires knowledge of inter-relationship among various climatic parameters, as well as short and long-term changes in the climate. In view of absence of written records of climate variations prior to the last ~100-150 yr, indirect techniques called proxies are used to infer past climate changes. An index sensitive to changes in the physico-chemical characteristics of the ambient environment and having good preservation potential (resistant to any pre- or post depositional change, especially the changes that can alter the signatures of environmental conditions) can be used to infer past climatic conditions.

The aquatic bodies, with continuous accumulation of sediments are among the ideal places to retrieve paleoclimatic proxies and records. The sediments accumulating at the bottom of aquatic bodies contain various indices that incorporate signatures of environmental conditions. In the aquatic bodies, the indices containing climatic signatures are immediately buried below subsequent sediments, to be preserved for long time. Even though the characteristics of sediments, including its physical and chemical properties, have been used to infer past climate (Bassinot, 1993; Weber *et al.*, 1997), the variations in the characteristics of the biological component of the sediments have been used most extensively. The preferred application of biological components over the abiogenic remains arises due to the high sensitivity and enhanced response of biological components to the changes in ambient environment. Since marine sediments contain plenty of remains of marine microorganisms (like coccolithophores, radiolarians, pteropods, ostracodes, foraminifera, and others), such remains serve as potential proxy to infer past climatic conditions. Out of these remains of various marine microorganisms, the hard outer skeletal remains of foraminifera, termed as tests, are among the most widely used paleoclimatic proxy.

As foraminifera are extremely sensitive to ambient conditions, and ubiquitous in almost all the marine realms, they are extensively used to reconstruct past climate. Ever since their discovery, numerous characteristics of foraminifera have been used to infer different seawater parameters. The initially developed techniques made use of changes in abundance and morphology of foraminifera and thus were non-destructive. The stable isotopic analysis of foraminifera were initiated in the early

fifties (Urey *et al.*, 1951; Emiliani and Epstein, 1953; Emiliani, 1955), while the first successful attempts to analyze trace element ratio of foraminiferal tests were made in early eighties (Boyle, 1981; Boyle and Keigwin, 1985) leading to the development of destructive techniques to reconstruct past climate by using foraminiferal tests. Despite the growing application of stable isotopic and trace element ratio of foraminifera to reconstruct paleoclimate, non-destructive techniques are still extensively applied not only to independently infer past climate but also to substantiate inferences drawn from the stable isotopic and trace element ratio. In this review, the traditional non-destructive foraminiferal proxies are summarized.

## **Foraminifera**

Foraminifera are unicellular, preferentially marine microorganisms with the single cell encased by a hard outer covering, called test. The foraminiferal test is made up either of calcium carbonate secreted by the cell (Calcareous), or various sized sediments adhered together with cement (Agglutinated). Foraminifera inhabit both the sediments on the sea floor (benthic foraminifera) (Figure 1) as well as the upper few hundred meters of the seawater column (planktic foraminifera) (Figure 2). Benthic foraminifera live both on the seafloor (epifaunal) and up to a few cm in the upper sediments (infaunal). The epifaunal and infaunal benthic foraminifera has a distinct morphology. Most of the infaunal forms are rectilinear, as it aids in burrowing, while the epifaunal forms are round in shape (Boltovskoy and Wright, 1976; Murray, 2006). A few benthic foraminifera, belonging to the family Allogromidae, thrive in fresh water and do not have hard exoskeleton like their marine counterparts. By using molecular systematic analysis, foraminifera have also been reported in soils collected from different terrestrial regions (Lejzerowicz *et al.*, 2010). The planktic foraminifera are depth stratified with a few species inhabiting the top mixed layer while a few others being abundant in the thermocline region. A vertical migration to cooler deep waters during the later phase of life for gametogenesis is reported in a few planktic foraminifera. The near coastal waters are almost exclusively inhabited by benthic foraminifera while both benthic and planktic foraminifera are found in oceanic regions deeper than ~75-100 m but above carbonate compensation depth (CCD). The agglutinated benthic foraminifera (Figure 3) are abundant in regions where carbonate dissolution is prevalent, including marshes, deep shelves or below the carbonate compensation depth (Murray, 2006; Schröder-Adams and Rooyen, 2011). Even though the planktic foraminifera live in the surface water column in deeper oceanic regions, their tests almost completely dissolve, before reaching the seafloor due to over saturation of deeper oceanic waters by carbon dioxide (Berger, 1967).

While the benthic foraminifera are tremendously diverse with thousands of species reported from different parts of the world oceans, there are only ~40 species of planktic foraminifera found in modern world oceans. The tests of foraminifera, especially benthic foraminifera, show wide variations in size, ornamentation and shape, from being single chambered (unilocular) to chambers arranged in one, two or many linear rows (uniserial, biserial and multiserial, respectively) as well as chambers arranged in circular fashion (spiral forms). The planktic foraminiferal tests in general, have globular and highly perforated chambers and also comparatively large aperture.

### **Foraminiferal Proxies**

The variation in the abundance and morphology of the tests is an adaptive response of the foraminifera to the changes in environment inhabited by foraminifera. Thus the changes in the foraminiferal tests in response to climatic changes, makes it possible to apply various foraminiferal characteristics to infer past environmental conditions. Various non-destructive characteristics of the foraminiferal tests that have been used to infer past climatic variations include:

1. Total Foraminiferal Number
2. Benthic:Planktic Ratio
3. Foraminiferal Fragmentation index
4. Foraminiferal Size Fraction
5. Species Assemblages
6. Species Abundance
7. Test Size/Diameter
8. Coiling Direction
9. Average Number of Chambers
10. Proloculus Size
11. Test Abnormalities
12. Test Weight

### **Total Foraminiferal Number**

The total foraminiferal number (TFN) is defined as the number of intact foraminiferal tests in unit dry sediment. All the intact benthic and planktic foraminiferal tests are picked from a pre-weighed representative fraction and the TFN is calculated by extrapolating the foraminiferal number in unit

gram of dry sediment. The representative fraction is taken either by quartering and coning the entire coarse fraction or by using a splitter. Although, initially it was proposed that the water depth or water mass characteristics are the major factors that control foraminiferal population, later it was realized that the food availability has the dominant control on the living benthic foraminiferal population (Jorissen *et al.*, 2007). In addition to food, benthic foraminiferal population also depends on dissolved oxygen content of the water. The organic matter flux and dissolved oxygen content are inter-related and a model was proposed to infer the relative role of these factors on benthic foraminiferal population (Jorissen *et al.*, 2007). Besides the food and dissolved oxygen, the foraminiferal population has definite environmental control, depending upon the water depth, turbidity, sediment characteristics, energy conditions in the region and other physico-chemical characteristics (Boltovskoy and Wright, 1976; Murray, 1991; Lee and Anderson, 1991). Dissolution during the sinking of planktic foraminiferal tests from the surface to the sea bottom as well as the post-depositional dissolution at the sea bottom, alters the foraminiferal population (Berger, 1967; 1968; Peterson and Prell, 1985; de Villiers, 2005). The agglutinated benthic foraminifera readily dissolve after death while a subsequent change in pore-water pH leads to partial or full dissolution of calcareous forms, thus altering the original foraminiferal population. Dissolution of calcareous foraminiferal tests also depends on the amount of organic matter settling in the sediments (Berkeley *et al.*, 2007). Since the planktic and benthic foraminifera represent the top several hundred meters of the water column, and bottom water as well as sediment characteristics, respectively, separate estimates of changes in population of benthic and planktic foraminifera are more efficient proxy than the combined population, i.e. total foraminiferal number. The variations in planktic, benthic or total foraminiferal population have been used for stratigraphic correlation (Berggren and Boersma, 1969; Reiss *et al.*, 1980), changes in the concentration of pollutants (see Nigam *et al.*, 2006 for review), bathymetric changes (Bandy, 1956; Sharma and Takayanagi, 1982; Nigam and Henriques, 1992) and shifts in lysocline depth (Cullen and Prell, 1984).

### **Benthic: Planktic Ratio**

Planktic foraminifera are rare in turbid shallow waters. Such regions also do not support vertical movement of planktic foraminifera, as required during gametogenesis in a few species, thus restricting their population. Therefore, the abundance of planktic foraminifera in general increases with depth away from the coast, till foraminiferal lysocline. Below the foraminiferal lysocline, significant dissolution of the foraminiferal tests leads to decreased foraminiferal abundance. Therefore, the changes in the relative abundance of benthic and planktic foraminifera indicate, changes in the bathymetry (Stehli and Creath, 1964; Wright, 1977), changes in the position of

foraminiferal lysocline as well as carbonate compensation depth (Pinxian *et al.*, 1995; Chen *et al.*, 1997), and productivity changes (Berger and Diester-Haass, 1988). Although the planktic foraminiferal population in general decreases away from the coast in all world oceans, the exact relationship between changes in relative abundance of planktic foraminifera and water depth varies from region to region. A paleodepth model for the eastern Arabian Sea was suggested on the basis of study of planktic foraminiferal percentage in surface sediments collected from 13 m to 1050 m water depth, thus allowing prediction of paleodepth in samples with 5-93% planktic foraminifera (Nigam and Henriques, 1992). Benthic to planktic ratio can also be used as productivity proxy in deep-sea regions devoid of differential dissolution. The productivity in such pelagic conditions is the product of the benthic/planktic ratio in percent, and the depth of deposition, whereas in hemipelagic conditions, the index has to be corrected for the distance from the land by multiplying it by the square-root of its distance from the land (Berger and Diester-Haass, 1988).

### Foraminiferal Fragmentation Index

The foraminiferal fragmentation index is defined as the number of fragments of foraminiferal tests per gram dry sediment. A part of the test which is less than two-thirds of the original test is defined as a foraminiferal fragment (Berger *et al.*, 1982). The foraminiferal fragmentation index (FFI) can be calculated by several ways.

A simple foraminiferal fragmentation index is defined as

$$\text{FFI (\%)} = [\text{Foraminiferal Fragments}/(\text{Foraminiferal Fragments} + \text{Whole Tests})]*100$$

Here tests with more than 2/3<sup>rd</sup> portion intact are considered whole tests, while any test with less than 2/3<sup>rd</sup> part intact is counted as fragment (Berger, 1970; Berger *et al.*, 1982).

According to Williams *et al.* (1985) and Malmgren (1987) the FFI can be calculated by the following equation

$$\text{FFI} = (\text{F}/8)/[(\text{F}/8) + \text{Whole Planktic Foraminifera}]$$

where F is number of fragments. As a foraminiferal test on an average usually breaks into 8 fragments and the number of fragments with respect to intact tests provides a better idea about dissolution, the number of fragments is divided by 8 plus the number of intact foraminiferal tests (Le and Shackleton, 1992). As the benthic foraminifera are more resistant to dissolution their tests are not included in the whole foraminifer count to assess changes in dissolution susceptibility.

Stuut *et al* (2002) proposed carbonate fragmentation index (CFI) based on grain size distributions of the bulk calcareous ooze. It is defined as the ratio of the 25-90  $\mu\text{m}$  and >90  $\mu\text{m}$

fractions. The CFI relies on the fact that the complete shells and fragments of juvenile foraminiferal shells dominate the 25-90  $\mu\text{m}$  of the bulk sediments, while  $>90 \mu\text{m}$  fraction primarily consists of adult foraminifera shells and fragments. The ratio of the 25-90  $\mu\text{m}$  and  $>90 \mu\text{m}$  fractions is suggested as a measure of the extent of carbonate dissolution induced fragmentation of the foraminifera shells (Stuut *et al.*, 2002).

A few other foraminiferal dissolution indices make use of relative ranking of species in order of their dissolution susceptibility as inferred from both the field samples as well as controlled experiments, while a few others make use of dissolution resistance of benthic foraminifera as compared to dissolution susceptibility of planktic foraminiferal tests (Conan *et al.*, 2002).

The fragmentation of the foraminiferal tests increases under high-energy conditions (increased turbidity), or increased dissolution susceptibility due to shoaling of lysocline or carbonate compensation depth. The variation in the amount of dissolved carbon-dioxide and carbonate ions also affects the rate of fragmentation of foraminiferal tests. Thus the foraminiferal fragmentation index provides information about turbidity currents as well as changes in the concentration of dissolved  $\text{CO}_2$  and carbonate ions (Peterson and Prell, 1985; Pinxian *et al.*, 1995; Chen *et al.*, 1997). Dittert and Henrich (2000) proposed, ultrastructure breakdown in *Globigerina bulloides* as proxy to infer the position of lysocline and carbonate compensation depth. Similarly, the degree of attrition of tests of planktic foraminiferal species *Globorotalia menardii* and *Globorotalia tumida* has been applied to estimate amount of calcium carbonate dissolved from the sediments (Ku and Oba, 1978).

### Foraminiferal Size Fraction

The foraminiferal tests, especially the planktic foraminiferal tests of different species fall within a particular size range. Thus the sediments of different size fractions are dominated by different species (Bé and Hutson, 1977; Peeters *et al.*, 1999; Kandiano and Bauch, 2002). The total planktic foraminiferal composition tremendously changes with size; the species richness and diversity increases with decreasing sieve size (Peeters *et al.*, 1999). The foraminiferal composition in different size fractions, of the water and sediment samples, differs significantly, suggesting large scale modulation of foraminiferal population during its post mortem settling on the sea floor (Peeters *et al.*, 1999). Therefore, the variation in weight of different sized foraminiferal tests in the sediments where sand fraction ( $>63 \mu\text{m}$ ) consists almost entirely of foraminiferal tests can be applied to document past variation in relative abundance of different species and to infer past climatic variations. While applying the test size/diameter to reconstruct past climate, however the

evolutionary changes in size of certain foraminiferal tests must also be considered, especially in studies including long time frame of several million years.

### Species Assemblages

Certain foraminiferal species prefer similar restricted environmental conditions, while a few others dwell in a wide range of seawater conditions. The group of species that inhabit similar and restricted environment is referred to as species assemblage. The species assemblages are often identified with cluster analysis and the characteristic species of an assemblage representing a particular environmental parameter like organic matter flux, dissolved oxygen, grain-size, water depth, may vary from region to region. Such species assemblages are thus representative of specific environmental conditions in a region (Murray, 1973; SenGupta, 1977; Lutze and Coulbourn, 1984; SenGupta *et al.*, 1993; Gooday, 1994; Mackensen *et al.*, 1995; Jorissen *et al.*, 1998; Loubere and Fariduddin, 1999; Gooday, 2003; Jorissen *et al.*, 2007; Kirci-Elmas, 2013). Therefore, temporal variation in such species assemblages is applied to infer variation in environmental conditions and is one of the most extensively used foraminiferal proxies to infer paleoclimatic variations (Nigam *et al.*, 1992; Mackensen *et al.*, 1994; Gupta *et al.*, 2001a; Gupta and Thomas, 2003). Application of species assemblages is preferred over that of the relative abundance of individual species due to a usually statistically significant number of individuals representing a species assemblage than individual species.

### Species Abundance

The relative abundance of any species can be counted by picking a minimum of ~300 specimens of foraminifera from a representative sample and then by dividing the number of specimens of the species by the total number of foraminiferal tests picked from the entire representative fraction. The relative abundance of benthic foraminiferal species is counted in fraction  $>63 \mu\text{m}$ , while that of planktic foraminiferal species in either  $>125 \mu\text{m}$  or  $>150 \mu\text{m}$  fraction. A relatively coarser fraction is taken to count the relative abundance of planktic foraminifera as the species identification is possible in only these coarser fractions. The juveniles of several planktic foraminifera are similar until they attain maturity and an average size of usually  $>125 \mu\text{m}$ . The foraminiferal studies from the surface sediments and sediment traps show significant changes in the abundance of foraminiferal species with changing ambient environmental conditions (Bé and Tolderlund, 1971; Boltovskoy and Wright, 1976; Bé and Hutson, 1977; Corliss, 1985; Thunnel and Honjo, 1987; Murray, 1973; 1991; Lee and Anderson, 1991; Naidu, 1993; Guptha *et al.*, 1997; Kawahata *et al.*, 2002; Fraile *et al.*, 2008; Panchang and Nigam, 2012; Rao *et al.*, 2013). Therefore, the fact that the abundance of certain



planktic as well as benthic foraminiferal species changes with a change in physico-chemical conditions, has been applied to infer paleoclimatic changes. Besides the use in paleoclimatic reconstruction, appearance and disappearance of foraminiferal species, termed as 'first appearance datum' and 'last appearance datum' are widely used to assign stratigraphy of a marine sedimentary sequence. Benthic foraminifera *Asterorotalia trispinosa*, is abundant in fine grained low salinity regions of the Bay of Bengal (Panchang and Nigam, 2012; Rao *et al.*, 2013) and thus its relative abundance has been used to reconstruct late Holocene monsoon changes (Panchang and Nigam, 2012). The variation in the abundance of benthic foraminiferal species *Epistominella exigua* has been applied to infer past changes in organic matter production (Gupta and Melice, 2003; Saraswat *et al.*, 2005). Among the planktic foraminifera, the temporal variation in the abundance of *Globigerina bulloides* has very often been used to infer past upwelling/productivity changes from the Indian Ocean (Prell and Curry, 1981; Naidu and Malmgren, 1996; Anderson *et al.*, 2002; Bassinot *et al.*, 2011; Naik *et al.*, 2014). The abundance and diameter of planktic foraminiferal species *Orbulina universa* is influenced by the ambient seawater temperature and salinity and thus, has been used to infer past temperature and salinity from the Indian Ocean (Colombo and Cita, 1980; Haenel, 1987; Nigam, 1990). The disappearance of pink variety of *Globigerinoides ruber* from the Indian and Pacific Ocean at 120,000 yr B.P. has been used as a biostratigraphic datum (Thompson *et al.*, 1979). While applying the relative abundance of species to reconstruct past climate or oceanography, it should however be kept in mind that its depth habitat may vary latitudinally (Saraswat and Khare, 2010). Additionally, the species abundance also depends on the size fraction taken for picking foraminifera. In the polar north Atlantic region, polar species *Neogloboquadrina pachyderma* (s) forms almost monospecific assemblages in larger size fractions (125-250  $\mu\text{m}$ , >150  $\mu\text{m}$ ) of MIS 8 to 7 whereas, the smaller size fractions (100-150  $\mu\text{m}$ , 80-150  $\mu\text{m}$ ) are dominated by the subpolar species *Turborotalita quinqueloba* (Kandiano and Bauch, 2002).

### Test Size/Diameter

The morphological changes, especially the variation in size (Figure 4), in response to the changes in the physico-chemical conditions like temperature, salinity, upwelling intensity, have long been recorded from both field as well as laboratory culture studies of foraminifera (Bé *et al.* 1973; Hecht, 1976; Caron *et al.*, 1987a; 1987b; Bijma *et al.*, 1990; Naidu and Malmgren, 1996; Schmidt *et al.*, 2004). Bé *et al.* (1973) showed that the test diameter of *Orbulina universa* from tropical and subtropical Indian Ocean shows a latitudinal variation and that the mean diameter of the test of *O. universa* in surface sediments and surface waters is strongly associated with the distribution of water masses in the southern Indian Ocean and northern Atlantic Ocean. Except physico-chemical controls

on the size of foraminifera, evolutionary changes have also been reported to result in an increase in the size of planktic foraminifera (Arnold *et al.*, 1995). A concurrent change in test diameter of several dominant deep-sea benthic foraminifera and dissolved oxygen content of the seawater, during the last 120 m yr, suggests a dominant control of climatic changes on benthic foraminiferal test diameter (Kaiho, 1999). The size and abundance of planktic foraminifera fluctuates with environmental factors such as temperature, primary productivity, and environmental variability, and reaches its maximum during optimum conditions suggesting a prominent effect of climatic conditions on planktic foraminiferal test size (Schmidt *et al.*, 2003; 2004). Abundance of *Epistominella exigua* is inversely related to its size suggesting increased instances of asexual reproduction during favourable conditions (Saraswat *et al.*, 2011). Thus, the temporal changes in foraminiferal test dimensions have been applied to infer past variations in physico-chemical parameters as well as in evolutionary studies (Naidu and Malmgren, 1996; Arnold *et al.*, 1995; Kaiho, 1999; Schmidt *et al.*, 2004). It should however be kept in mind that the close correspondence between planktic foraminiferal test size and climatic parameters is confined only to the regions of large climate variability like frontal systems and upwelling zones, whereas such a relationship is obscure in regions with relatively stable climate (Schmidt *et al.*, 2004).

### Coiling Direction

The foraminiferal tests, especially those of benthic foraminifera, show a wide variation in the number and arrangement of chambers. Out of several thousands of species of foraminifera, chambers are arranged trochospirally, in a few foraminiferal species. In the species belonging to this group, the chambers may be arranged either in dextral (clockwise) or sinistral (anticlockwise) manner (Figure 5). The coiling direction is determined by placing the tests in spiral view and following the direction of addition of chambers beginning from the proloculus and moving towards the last added chamber. If possible, a minimum of 50 specimens of a species should be picked from each interval while assessing temporal changes in coiling direction, in order to arrive at a statistically significant interpretation. Various studies have shown that the ambient environmental conditions influence the direction in which the chambers are arranged in certain species of foraminifera (Bolli, 1950; Ericson, 1959; Nigam and Khare, 1992; Naidu and Malmgren, 1996). This preferred coiling pattern of both benthic and planktic foraminiferal species has been applied to infer past climatic conditions (Ericson *et al.*, 1954; Bé, 1960; Thiede, 1971; Keany and Kennett, 1972; Williams, 1976; Nigam and Khare, 1992; West *et al.*, 2004). The dextral *Rotalidium annectens* are preferentially microspheric suggesting that mode of reproduction influences the coiling pattern (Nigam and Khare, 1992). The relative abundance of dextral and sinistral forms of planktic foraminifera *Neogloboquadrina*

*pachyderma* was frequently used to reconstruct past seawater temperature, with the sinistral form being dominant in colder waters (Bandy, 1960; Keany and Kennett, 1972; Naik *et al.*, 2013). Even though a similar possible control of regional seawater properties on the coiling direction of *Globorotalia menardii* was suggested by Bolli (1971), but no such relationship between coiling direction and sea water properties during the late Quaternary was found in the northern Indian Ocean, wherein only an insignificant percentage of specimens were dextrally coiled (Bhonsale and Saraswat, 2012). However, of late, with the application of molecular biology techniques, it has been established that the differently coiled specimens, formerly considered to belong to the same species, are in fact different species (Bauch *et al.*, 2003; Darling *et al.*, 2006).

### Average Number of Chambers

Many species of foraminifera have bimodal reproduction with morphologically distinct sexual and asexual phases. The tests belonging to offsprings of sexual and asexual mode of reproduction have several differences, including different number of chambers. The environmental conditions influence the mode of reproduction in foraminifera (Boltovskoy and Wright, 1976; Nigam and Rao, 1987; Nigam and Caron, 2000) and thus the number of chambers in the individuals belonging to the same foraminiferal species. This characteristic of variation in average number of chambers, in certain foraminiferal species as a result of changing mode of reproduction under different environmental conditions, has been used to understand ecology of foraminifera and further proposed as a tool to infer paleoclimatic variations. The number of chambers in a test is counted starting from the proloculus and moving along the direction of addition of chambers till the last formed chamber (Figure 6). On the basis of inverse relationship between *Epistominella exigua* abundance and the number of chambers, it was inferred that *E. exigua* prefers an asexual mode of reproduction during favorable conditions (Saraswat *et al.*, 2011).

### Proloculus Size

Cushman (1905) coined the term proloculus to represent the first formed chamber of the foraminiferal test. The changing mode of reproduction as discussed in the previous section, also influences the size of the proloculus (Figure 4) with megalospheric specimens having larger proloculus as compared to microspheric specimens (Boltovskoy and Wright, 1976). In a few benthic foraminiferal species, the distinction between microspheric and megalospheric forms is difficult only from the size of the proloculus. In such cases, the number of chambers (more in microspheric forms as compared to megalospheric forms) and the size of the tests (microspheric being larger than megalospheric) can help substantiate the differentiation between the two forms. The mean proloculus

size of benthic foraminifera *Cavarotalia annectens*, calculated as the ratio of microspheric to megalospheric forms is inversely proportional to salinity and temperature in the shallow water regions of the eastern Arabian Sea (Nigam and Rao, 1987). As noted above, that the mode of reproduction in foraminifera is environmentally controlled, the mean proloculus size of selected foraminiferal species have been used to infer past climatic variations (Nigam and Rao, 1987; Nigam and Sarkar, 1993; Nigam and Khare, 1995; Saraswat *et al.*, 2005).

### Test Abnormalities

Majority of the foraminifera follow a well defined pattern to add chambers during its growth till it reaches maturity. This normal course of addition of chambers for any species, can be assessed by looking at the general population of that particular species in a region. Any deviation from this normal course of addition of chambers is termed as abnormality. The regions where deformed tests comprise upto 1% of the total population, are termed as normal while locations with >1% abnormal tests are termed as stressed (Alve, 1991). A comparatively large number of abnormal foraminiferal tests have been reported in sediments collected from ecologically stressed (both naturally as well as anthropogenically) environments (such as very high or low salinity, temperature, pH, or high concentration of pollutants like heavy metal, sewage) as well as in laboratory culture experiments wherein foraminifera were subjected to pollutants (Boltovskoy *et al.* 1991; Yanko *et al.*, 1998; Alve, 1995; Stouff *et al.*, 1999; Scott *et al.*, 2001; Saraswat *et al.*, 2004; Nigam *et al.*, 2006a; 2006b; 2009; Caruso *et al.*, 2011). The abnormalities in foraminiferal tests from ecologically stressed environments include stunted tests, comparatively smaller or bigger chambers, chambers oriented away from the normal plane of addition of chambers, tests with regressed or completely absent ornamentation. Thus the unusual presence of abnormal foraminiferal tests is applied as a proxy to infer ecologically-stressed environmental conditions (Nagy and Alve, 1987; Schafer *et al.*, 1991; Scott *et al.*, 1995; Panchang *et al.*, 2005). However, the application of test abnormalities to infer 'specific' ecologically-stressed environment is hampered due to the report of nearly similar test abnormalities from differently stressed environments and further by no correlation between abundance of deformed tests and known sources of pollution (Cosentino *et al.*, 2013).

### Test Weight

A majority of the foraminiferal tests are calcareous. This shell calcite is a mixture of primary calcite and secondary calcite. Most of the planktic foraminifera grow in size by adding chambers secreted from primary calcite. In contrast, secondary calcification thickens existing chamber walls, which can double a shell's mass without significantly affecting its size (Lohmann, 1995). The different effect

that primary and secondary calcification has on shell size allow their relative proportions to be estimated. The secondary calcite is added to the shell when there is excess  $[\text{CO}_3^{2-}]$  available, *i.e.*, the more the  $[\text{CO}_3^{2-}]$  concentration, more the secondary calcification will occur (Broecker and Clark, 1999). Laboratory culture studies under controlled conditions indicate that an increase in ambient  $[\text{CO}_3^{2-}]$  increases the calcification rates and size-normalized weight (SNW) of planktic foraminifera suggesting a dominant control of carbonate ions which are two orders of magnitude smaller than that of calcium ions in seawater, during the calcification (Bijma *et al.*, 2002). Hence, if we can determine the secondary calcite amount then we can infer the  $[\text{CO}_3^{2-}]$  concentration of the seawater. Therefore, the species in which secondary calcification changes only shell mass (e.g. *Pulleniatina obliquiloculata* and *Globoquadrina dutertrei*) and not the size, the amount of secondary calcite added to a shell reflects the change in the  $[\text{CO}_3^{2-}]$  concentration of the seawater.

The size normalized weight of a foraminiferal species can be determined either by taking the average weight of specimens (with a precision of 1  $\mu\text{g}$ ) picked from a narrow size range (usually 50  $\mu\text{m}$ ) obtained by sieving the coarse fraction (Broecker and Clark, 2001) or by measuring the diameter of individual tests and by dividing the weight of the tests by the average measured size (Barker and Elderfield, 2002). Prior to weighing, the specimens should be thoroughly cleaned to remove organic matter/clay adhering to the tests. The changes in shell weight have been used to infer decrease in biogenic marine carbon sequestration due to increasing ocean acidification as a result of high atmospheric green-house gas concentration (de Moel *et al.*, 2009). The shell weight is often used to assess the degree of dissolution and its potential effect on the trace element composition of the tests (Naik *et al.*, 2014).

The use of shell weight as carbonate ion proxy however has a few potential problems. The shell weight may not always increase but decrease due to decrease in the carbonate ion concentration of sediment pore water. Additionally, the shell weight can also vary due to changes in environmental factors unrelated to carbonate ion concentration (Broecker and Clark, 2001). Only a weak covariance of shell weight with climate indicators and a smaller than expected offset between down-core records at a shallow and deeper sites in the Ontong-Java Plateau, suggests that the controls on shell weight are more complicated and also questions the use of shell weight to reconstruct past carbonate saturation and carbon dioxide levels (de Villiers, 2003; 2005). Large interspecies- and intraspecies-specific variations were reported not only in the strength and gradient of the relationship between  $[\text{CO}_3^{2-}]$  and the shell weight but also the sign of this relationship, suggesting that shell weight is not a reliable proxy to determine past  $[\text{CO}_3^{2-}]$  changes (Beer *et al.*, 2010). A few other studies reported a dominant control of phosphate ion concentration in seawater on shell weight in addition to  $[\text{CO}_3^{2-}]$

and seawater temperature and minimal effect of optimum growth conditions, thus implying that further work is required before the shell weight can be used as a reliable proxy for  $[\text{CO}_3^{2-}]$  (Aldridge *et al.*, 2012).

## Conclusions

Reconstructing the past climate is important and characteristics of marine microfossils, foraminifera offer a wealth of information about past climate. Changes in abundance, morphology and chemical composition of the foraminiferal tests are used to reconstruct past climate. The non-destructive foraminiferal proxies help in understanding seawater temperature, salinity, productivity, dissolved oxygen content, lysocline movement, sea level changes, organic matter flux, monsoon changes, circulation and several others. The non-destructive foraminiferal proxies not only help to reconstruct qualitative changes in past climate but are also effective to quantify a few parameters like paleodepth and temperature. The foraminiferal proxies are region specific and work best if calibrated regionally.

## Acknowledgements

The review was written as Prof. S.C. Lakhotia, the Editor-in-Chief, Proceedings of the Indian National Science Academy, provided an opportunity to the INSA Young Scientist Medal winners to contribute a review of the work related to their field of specialization. The author acknowledges the financial support by the Council of Scientific and Industrial Research, Department of Science and Technology, and Ministry of Earth Sciences, New Delhi.

## References

- Alve E (1991) Benthic foraminifera in sediment cores reflecting heavy metal pollution in Sørøfjord, Western Norway *J Foraminiferal Res* **21** 1-19
- Alve E (1995) Benthic foraminiferal responses to estuarine pollution: a review *J Foraminiferal Res* **25** 190-203.
- Aldridge D, Beer C J and Purdie D A (2012) Calcification in the planktonic foraminifera *Globigerina bulloides* linked to phosphate concentrations in surface waters of the North Atlantic Ocean *Biogeosciences* **9** 1725-1739
- Anderson D M, Overpeck J T and Gupta A K (2002) Increase in the Asian southwest monsoon during the past four centuries *Science* **297** 596-599
- Arnold A J, Kelly D C and Parker W C (1995) Causality and Cope's rule: evidence from the planktonic foraminifera *J Paleontol* **69** 203-210
- Bandy O L (1956) Ecology of foraminifera in the northeastern Gulf of Mexico *U S Geol Surv Prof Pap* **724-G** 179-204

- Bandy O L (1960) The geologic significance of coiling ratios in the foraminifera *Globigerina pachyderma* *J Paleontol* **4** 671-681
- Barker S and Elderfield H (2002) Foraminiferal calcification response to glacial-interglacial changes in atmospheric CO<sub>2</sub> *Science* **297** 833-836
- Bassinot F C (1993) Sonostratigraphy of tropical Indian Ocean giant piston cores: toward a rapid and high-resolution tool for tracking dissolution cycles in Pleistocene carbonate sediments *Earth Planet Sci Lett* **120** 327-344
- Bassinot F C, Marzin C, Braconnot P, Marti O, Mathien-Blard E, Lombard F and Bopp L (2011) Holocene evolution of summer winds and marine productivity in the tropical Indian Ocean in response to insolation forcing: data-model comparison *Clim Past* **7** 815-829
- Bauch D, Darling K, Simstich J, Bauch H A, Erlenkeuser H and Kroon D (2003) Palaeoceanographic implications of genetic variation in living North Atlantic *Neogloboquadrina pachyderma* *Nature* **424** 299-302
- Bé A W H (1960) Ecology of Recent planktonic foraminifera: Part 2.-Bathymetric and seasonal distributions in the Sargasso Sea off Bermuda *Micropaleontology* **6** 373-392
- Bé A W H and Tolderlund D S (1971) Distribution and ecology of living planktonic foraminifera in surface waters of the Atlantic and Indian Ocean. In: *The Micropaleontology of Oceans* (Eds: Funnel B M and Riedel W R) pp 105-149
- Bé A W H, Harrison S M and Lott L (1973) *Orbulina universa* d'Orbigny in the Indian Ocean *Micropaleontology* **19** 150-192
- Bé, A W H and Hutson A (1977) Ecology of planktonic foraminifera and biogeographic patterns of life and fossil assemblages in the Indian Ocean *Micropaleontology* **23** 269-414
- Beer C J, Schiebel R and Wilson P A (2010) Testing planktic foraminiferal shell weight as a surface water [CO<sub>3</sub><sup>2-</sup>] proxy using plankton net samples *Geology* **38** 103-106
- Berger W H (1967) Foraminiferal ooze: solution at depths *Science* **156** 383-385
- Berger W H (1968) Planktonic Foraminifera: selective solution and paleoclimatic interpretation *Deep-Sea Res* **15** 31-43.
- Berger W H (1970) Planktonic foraminifera: selective solution and the lysocline *Mar Geol* **8** 111-138.
- Berger WH, Bonneau M-C and Parker F L (1982) Foraminifera on the deep-sea floor: lysocline and dissolution rate *Oceanol Acta* **5** 249-258
- Berger W H and Diester-Haass L (1988) Paleoproductivity: The benthic planktonic ratio in foraminifera as a productivity index *Mar Geol* **81** 15-25
- Berggren W A and Boersma A (1969) Late Pleistocene and Holocene planktonic foraminifera from the Red Sea. In: *Hot Brines and Recent Heavy Metal Deposits in the Red Sea* (Eds: Degens E T and Ross D A) pp 282-298, Springer-Verlag, Berlin
- Berkeley A, Perry C T, Smithers S G, Horton B P, Taylor K G (2007) A review of the ecological and taphonomic controls on foraminiferal assemblage development in intertidal environments *Earth-Sci Rev* **83** 205-230

- Bhonsale S and Saraswat R (2012) Abundance and size variation of *Globorotalia menardii* in the northeastern Indian Ocean during the Late Quaternary *J Geol Soc India* **80** 771-782
- Bijma J, Faber W W and Hemleben C (1990) Temperature and salinity limits for growth and survival of some planktonic foraminifers in laboratory cultures *J Foraminiferal Res* **20** 95-116.
- Bijma J, Hönisch B and Zeebe R E (2002) Impact of the ocean carbonate chemistry on living foraminiferal shell weight: Comment on “Carbonate ion concentration in glacial-age deep waters of the Caribbean Sea” by W. S. Broecker and E. Clark *Geochem Geophys Geosyst* **3** 1064, doi: 10.1029/2002GC000388
- Bolli H (1950) The direction of coiling in the evolution of some Globorotaliidae *Contr Cushman Found Foraminiferal Res* **1** 82-89
- Bolli H M (1971) The direction of coiling in planktonic foraminifera. In: Micropaleontology of Oceans (Eds: Funnel B M and Riedel W R) pp 639-648, Cambridge University Press, Cambridge
- Boltovskoy E and Wright R (1976) Recent Foraminifera. Dr. W. Junk b.v. Publishers The Hague
- Boltovskoy E, Scott D B and Medioli F S (1991) Morphological variations of benthic foraminiferal tests in response to changes in ecological parameters: a review *J Paleontol* **65** 175-185
- Boyle E A (1981) Cadmium, zinc, copper, and barium in foraminiferal tests *Earth Planet. Sci Lett* **53** 11-35
- Boyle E A and Keigwin L D (1985) Comparison of Atlantic and Pacific paleochemical records for the Last 215,000 years: Changes in deep ocean circulation and chemical inventories *Earth Planet Sci Lett* **76** 135-150
- Broecker W and Clark E (1999) CaCO<sub>3</sub> size distribution: a paleo carbonate ion proxy *Paleoceanography* **14** 596-604
- Broecker W S and Clark E (2001) An evaluation of Lohmann's foraminifera-weight dissolution index *Paleoceanography* **16** 531-534
- Caron D A, Faber W W J and Bé A W H (1987a) Effects of temperature and salinity on the growth and survival of the planktonic foraminifer *Globigerinoides sacculifer* *J Mar Biol Assoc UK* **67** 323-342
- Caron D A, Faber W W J and Bé A W H (1987b) Growth of the spinose planktonic foraminifer *Orbulina universa* in laboratory culture and the effect of temperature on life processes *J Mar Biol Assoc UK* **67** 343-358
- Caruso A, Cosentino C, Tranchina L and Brai M (2011) Response of benthic foraminifera to heavy metal contamination in marine sediments (Sicilian coasts, Mediterranean Sea) *Chem Ecol* **27** 9-30
- Chen M T, Huang CY and Wei K Y (1997) 25,000-year late Quaternary records of carbonate preservation in the South China Sea *Palaeogeogra Palaeoclimatol Palaeoecol* **129** 155-169
- Colombo M R and Cita M B (1980) Changes in size and test porosity of *Orbulina universa* d'orbigny in the Pleistocene record of Cape Bojador (DSDP site 397, eastern north Aatlantic) *Mar Micropaleontol* **5** 13-29
- Conan S M-H, Ivanova E M and Brummer G-J A (2002) Quantifying carbonate dissolution and calibration of foraminiferal dissolution indices in the Somali Basin *Mar Geol* **182** 325-349



- Corliss B H (1985) Microhabitats of benthic foraminifera within deep-sea sediments *Nature* **314** 435-438
- Cosentino C, Pepe F, Scopelliti G, Calabrò M and Caruso A (2013) Benthic foraminiferal response to trace element pollution-the case study of the Gulf of Milazzo, NE Sicily (Central Mediterranean Sea) *Environ Monit Assess* **185** 8777-8802
- Cullen J L and Prell WL (1984) Planktonic foraminifera of the northern Indian Ocean: distribution and preservation in surface sediments *Mar Micropaleontol* **9** 1-52
- Cushman J A (1905) Developmental stages in the Lagenidae *Am Naturalist* **39** 537-553
- Darling K F, Kucera M, Kroon D and Wade C M (2006) A resolution for the coiling direction paradox in *Neogloboquadrina pachyderma* *Paleoceanography* **21** doi:10.1029/2005PA001189
- de Moel H, Ganssen G M, Peeters F J C, Jung S J A, Kroon D, Brummer G J A and Zeebe R E (2009) Planktic foraminiferal shell thinning in the Arabian Sea due to anthropogenic ocean acidification? *Biogeosciences* **6** 1917-1925
- de Villiers S (2003) A 425 kyr record of foraminiferal shell weight variability in the western equatorial Pacific *Paleoceanography* **18** 1080, doi:10.1029/2002PA000801
- de Villiers S (2005) Foraminiferal shell-weight evidence for sedimentary calcite dissolution above the lysocline *Deep-Sea Res* **52** 671-680
- Dittert N and Henrich R (2000) Carbonate dissolution in the South Atlantic Ocean: evidence from ultrastructure breakdown in *Globigerina bulloides* *Deep-Sea Res* **47** 603-620
- Emiliani C (1955) Pleistocene temperatures *J Geol* **63** 538-578
- Emiliani C and Epstein S (1953) Temperature variations in the Lower Pleistocene of southern California *J Geol* **61** 171-181
- Ericson D B (1959) Coiling direction of *Globigerina pachyderma* as a climatic index *Science* **130** 219-220
- Ericson D B, Wollin G and Wollin J (1954) Coiling direction of *Globorotalia truncatulinoides* in deep-sea cores *Deep-Sea Res* **2** 152-158
- Fraile I, Schulz M, Mulitza S and Kucera M (2008) Predicting the global distribution of planktonic foraminifera using a dynamic ecosystem model *Biogeosciences* **5** 891-911
- Gooday A J (1994) The biology of deep-sea foraminifera: A review of some advances and their applications in paleoceanography *Palaios* **9** 14-41
- Gooday A J (2003) Benthic foraminifera (protista) as tools in deep-water palaeoceanography: Environmental influences on faunal characteristics *Adv Mar Bio* **46** 1-90
- Gupta A K (2004) Origin of agriculture and domestication of plants and animals linked to early Holocene climate amelioration *Curr Sci* **87** 54-59
- Gupta A K and Srinivasan M S (1990) Response of northern Indian Ocean deep-sea benthic foraminifera to global climates during Pliocene-Pleistocene *Mar Micropaleontol* **16** 77-91
- Gupta A K and Srinivasan M S (1992) *Uvigerina proboscidea* abundances and paleoceanography of the northern Indian Ocean DSDP Site 214 during the Late Neogene *Mar Micropaleontol* **19** 355-367

- Gupta A K, Joseph S and Thomas E (2001) Species diversity of Miocene deep-sea benthic foraminifera and watermass stratification in the northeastern Indian Ocean *Micropaleontology* **47** 111-124
- Gupta A K and Melice J –L (2003) Orbital forcing of the Plio-Pleistocene Indian monsoons: benthic foraminiferal proxies from ODP Site 758 *Curr Sci* **85** 179-184
- Gupta A K and Thomas E (2003) Initiation of Northern Hemisphere glaciation and strengthening of the northeast Indian monsoon: Ocean Drilling Program Site 758, eastern equatorial Indian Ocean *Geology* **31** 47-50
- Guptha M V S, Curry W B, Ittekkot V and Murlinath A S (1997) Seasonal variation in the flux of planktonic foraminifera: sediment trap results from the Bay of Bengal, northern Indian Ocean *J Foraminiferal Res* **27** 5-19
- Haenel P (1987) Paleoceanographic value of *Orbulina universa* d'orbigny (Foraminifera) *Oceanol Acta* **10** 15-25
- Haug G H, Günther D, Peterson L C, Sigman D M, Hughen K A and Aeschlimann B (2003) Climate and the collapse of Maya civilization *Science* **299** 1731-1735
- Hecht A D (1976) An ecologic model for test size variation recent planktonic foraminifera: application to the fossil record *J Foraminiferal Res* **6** 295-311
- Jorissen F J, Wittling I, Peypouquet J P, Rabouille C and Relexans J C (1998) Live benthic foraminiferal faunas off Cape Blanc, NW-Africa: Community structure and microhabitats *Deep Sea Res* **45** 2157-2188
- Jorissen F J, Fontanier C and Thomas E (2007) Paleoceanographical proxies based on deep-sea benthic foraminiferal assemblage characteristics. In: Proxies in Late Cenozoic Paleoceanography: Pt. 2: Biological tracers and biomarkers (Eds: Hillaire-Marcel C and de Vernal A) pp 263-326, Elsevier.
- Kaiho K (1999) Evolution in the test size of deep-sea benthic foraminifera during the past 120 m.y. *Mar Micropaleontol* **37** 53-65
- Kandiano E S and Bauch H A (2002) Implications of planktic foraminiferal size fractions for the glacial-interglacial paleoceanography of the polar north Atlantic *J Foraminiferal Res* **32** 245-251
- Kawahata H, Nishimura A and Gagan M K (2002) Seasonal change in foraminiferal production in the western equatorial Pacific warm pool: evidence from sediment trap experiments *Deep-Sea Res* **49** 2783-2800
- Keany J and Kennet J P (1972) Pliocene-early Pleistocene paleoclimatic history recorded in Antarctic-Subantarctic deep-sea cores *Deep-Sea Res* **19** 529-548
- Kirci-Elmas E (2013) Benthic foraminiferal distribution (living and dead) from a permanently stratified marginal sea (Marmara Sea, Turkey) *J Foraminiferal Res* **43** 340-360
- Ku T L and Oba T (1978) A method for quantitative evaluation of carbonate dissolution in deep-sea sediments and its application to paleoceanographic reconstruction *Quat Res* **10** 112-129
- Le J and Shackleton N J (1992) Carbonate dissolution fluctuations in the western equatorial Pacific during the late Quaternary *Paleoceanography* **7** 21-42
- Lee J J and Anderson O R (1991) *Biology of Foraminifera*. Academic Press.

- Lejzerowicz F, Pawlowski J, Fraissinet-Tachet L and Marmeisse R (2010) Molecular evidence for widespread occurrence of foraminifera in soils *Environ Microbiol* **12** 2518-26
- Lohmann G P (1995) A model for variation in the chemistry of planktonic foraminifera due to secondary calcification and selective dissolution *Paleoceanography* **10** 445-457
- Loubere P and Fariduddin M (1999) Quantitative estimation of global patterns of surface ocean biological productivity and its seasonal variation on time scales from centuries to millennia *Glob Biochem Cycles* **13** 115-133
- Lutze G and Coulbourn W (1984) Recent benthic foraminifera from the continental margin off northwest Africa: community structure and distribution *Mar Micropaleontol* **8** 361-401
- Mackensen A, Grobe H, Hubberten H -W and Kuhn G (1994) Benthic foraminiferal assemblages and the  $\delta^{13}\text{C}$ -signal in the Atlantic sector of the Southern Ocean: glacial-to-interglacial contrasts. In: Carbon cycling in the glacial ocean: constraints on the ocean's role in global change (Eds: Zahn R, Pederson T F, Kaminiski M A and Labeyrie L) **117** pp 105-144, Springer-Verlag, Berlin, Heidelberg, NATO ASI Series
- Mackensen A, Schmiedl G, Harloff J and Giese M (1995) Deep sea foraminifera in the South Atlantic Ocean: ecology and assemblage generation *Micropaleontology* **41** 342-358
- Malmgren B A (1987) Differential dissolution of Upper Cretaceous planktonic foraminifera from a temperate region of the south Atlantic Ocean *Mar Micropaleontol* **11** 251-271
- Murray J W (1973) *Distribution and Ecology of Living Benthic Foraminiferids*. Heinemann, London.
- Murray J W (1991) *Ecology and Paleocology of Benthic Foraminifera*. John Wiley, New York and Longman Scientific and Technical, Harlow, U.K..
- Murray J W (2006) *Ecology and Applications of Benthic Foraminifera*. Cambridge University Press.
- Murray J W and Alve E (1999) Natural dissolution of shallow water benthic foraminifers: taphonomic effects on the paleoecological record *Paleogeogra Paleoclimatol Paleoecol* **146** 195-209
- Nagy J and Alve E (1987) Temporal changes in foraminiferal faunas and impact of pollution in Sandebucta, Oslo Fjord *Mar Micropaleontol* **12** 109-128
- Naidu P D (1993) Distribution patterns of Recent planktonic foraminifera in surface sediments of the western continental margin of India *Mar Geol* **110** 403-418
- Naidu P D and Malmgren B A (1996) Relationship between Late Quaternary upwelling history and coiling properties of *Neogloboquadrina pachyderma* and *Globigerina bulloides* in the Arabian Sea *J Foraminiferal Res* **26** 64-70
- Naik D K, Saraswat R, Khare N, Pandey A C and Nigam R (2014) Hydrographic changes in the Agulhas Recirculation Region during the late Quaternary *Clim Past* **10** 745-758
- Nigam R (1990) Palaeoclimatic implications of size variation in *Orbulina universa* in a core from the North Indian Ocean *Curr Sci* **59** 46-47
- Nigam R and Caron D A (2000) Does temperature affect dimorphic reproduction in benthic foraminifera? A culture experiment on *Rosalina leei* *Curr Sci* **79** 1105-1106

- Nigam R and Henriques P J (1992) Planktonic percentage of foraminiferal fauna in surface sediments of the Arabian sea (Indian Ocean) and a regional model for paleodepth determination *Palaeogeogra Palaeoclimatol Palaeoecol* **91** 89-98
- Nigam R and Khare N (1992) The reciprocity between coiling direction and dimorphic reproduction in benthic foraminifera *J Micropalaeontol* **11** 221-228
- Nigam R and Khare N (1995) Significance of correspondence between river discharge and proloculus size of benthic foraminifera in paleomonsoonal studies *Geo-Mar Lett* **15** 45-50
- Nigam R, Khare N and Borole D V (1992) Can benthic foraminiferal morpho-groups be used as indicator of paleomonsoonal precipitation? *Est Coastal Shelf Sci* **34** 533-542
- Nigam R and Rao A S (1987) Proloculus size variation in Recent benthic foraminifera: Implications for paleoclimatic studies *Est Coastal Shelf Sci* **4** 649-655
- Nigam R and Sarkar A (1993) Mean proloculus size,  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  variations in recent benthic foraminifera from the west coast of India and their climatic implications *Ind J Earth Sci* **20** 1-6
- Nigam R, Saraswat R and Kurtarkar S R (2006a) Laboratory experiment to study effect of salinity variations on benthic foraminiferal species - *Pararotalia nipponica* (Asano) *J Geol Soc India* **67** 41-46
- Nigam R, Saraswat R and Panchang R (2006b) Application of foraminifers in ecotoxicology: Retrospect, prospect and prospect *Environ Int* **32** 273-283
- Nigam R, Linshy V N, Kurtarkar S R and Saraswat R (2009) Effects of sudden stress due to heavy metal mercury on benthic foraminifer *Rosalina leei*: Laboratory culture experiment *Mar Pollut Bull* **59** 362-368
- Panchang R and Nigam R (2012) High resolution climatic records of the past ~489 years from Central Asia as derived from benthic foraminiferal species, *Asterorotalia trispinosa* *Mar Geol* **307-310** 88-104
- Panchang R, Nigam R, Baig N and Nayak G N (2005) A foraminiferal testimony for the reduced adverse effects of mining in Zuari Estuary, Goa *Int J Environ Stud* **62** 579-591
- Peeters F, Ivanova E, Conan S, Brummer G-J, Ganssen G, Troelstra S and van Hinte J (1999) A size analysis of planktic foraminifera from the Arabian Sea *Mar Micropaleontol* **36** 31-63
- Peterson L C and Prell W L (1985) Carbonate dissolution in recent sediments of the eastern equatorial Indian Ocean: preservation patterns and carbonate loss above the lysocline *Mar Geol* **64** 259-290
- Pinxian W, Luejiang W, Yunhua B and Zhimi J (1995) Late Quaternary paleoceanography of the South China Sea: surface circulation and carbonate cycles *Mar Geol* **127** 145-165
- Polyak V J and Asmerom Y (2001) Late Holocene climate and cultural changes in the southwestern United States *Science* **294** 148-151
- Prell W L and Curry W B (1981) Faunal and isotopic indices of monsoonal upwelling: western Arabian Sea *Oceanol Acta* **4** 91-98
- Rao N R, Jayaprakash M and Velmurugan P M (2013) The ecology of *Asterorotalia trispinosa* (Thalman, 1933)- New insights from Muthupet Lagoon, southeast coast of India *J Foraminiferal Res* **43** 14-20

- Reiss Z, Luz B, Almogi-Labin A, Halicz E, Winter A, Wolf M and Ross D A (1980) Late Quaternary paleoceanography of the Gulf of Aqaba (Elat), Red Sea *Quat Res* **14** 294-308
- Robbins S G, Blevins K E, Cox B, Gray K and Mushrif-Tripathy V (2013) Infection, disease, and biosocial processes at the end of the Indus Civilization *PLoS One* **8** e84814. doi:10.1371/journal.pone.0084814
- Saraswat R, Kurtarkar SR, Mazumder A and Nigam R (2004) Foraminifers as indicators of marine pollution: a culture experiment with *Rosalina leei* *Mar Poll Bull* **48** 91-96
- Saraswat R, Nigam R and Barreto L (2005) Palaeoceanographic implications of abundance and mean proloculus diameter of benthic foraminiferal species *Epistominella exigua* in sub-surface sediments from distal Bay of Bengal fan *J Earth Syst Sci* **114** 453-458
- Saraswat R and Khare N (2010) Deciphering the modern calcification depth of *Globigerina bulloides* in the southwestern Indian Ocean from its oxygen isotopic composition *J Foraminiferal Res* **40** 220-230
- Saraswat R, Deopujari A, Nigam R and Henriques P J (2011) Relationship between abundance and morphology of benthic foraminifera *Epistominella exigua*: Palaeoclimatic implications *J Geol Soc India* **77** 190-196
- Schafer C T, Collins E S and Smith J N (1991) Relationship of foraminifera and thecamoebian distributions to sediments contaminated by pulp mill effluent: Saguenay Fiord, Quebec, Canada *Mar Micropaleontol* **17** 255-283
- Schmidt D N, Renaud S and Bollmann J (2003) Response of planktic foraminiferal size to late Quaternary climate change *Paleoceanography* **18** doi:10.1029/2002PA000831
- Schmidt D N, Renaud S, Bollmann J, Schiebel R and Thierstein R (2004) Size distribution of Holocene planktic foraminifer assemblages: biogeography, ecology and adaptation *Mar Micropaleontol* **50** 319-338
- Schröder-Adams C J and Rooyen D V (2011) Response of Recent Benthic Foraminiferal Assemblages to Contrasting Environments in Baffin Bay and the Northern Labrador Sea, Northwest Atlantic *Arctic* **64** 317-341
- Scott D B, Schafer C T, Honig C and Younger D C (1995) Temporal variations of benthic foraminiferal assemblages under or near aquaculture operations: documentation and impact history *J Foraminiferal Res* **25** 224-235
- Scott D B, Medioli F S and Schafer C T (2001) *Monitoring in Coastal Environments using Foraminifera and Thecamoebian Indicators*, Cambridge University Press
- SenGupta B K (1977) Depth distribution of modern benthonic foraminifera on continental shelves of the world Ocean *Indian J Earth Sci* **4** 60-83
- SenGupta B K (1991) *Modern Foraminifera*. Klower Academic Publishers.
- SenGupta B K and Machain-Castillo M L (1993) Benthic foraminifera in oxygen-poor habitats *Mar Micropaleontol* **20** 183-201
- Sharma V and Takayanagi Y (1982) Paleobathymetric history of late Neogene foraminiferal assemblages of the Kakugawa area, Central Japan *Sci Rep Tohoku Imp Univ* **52** 77-90
- Stehli F G and Creath W B (1964) Foraminiferal ratios and regional environments *Bull Am Assoc Pet Geol* **48** 1810-1827

- Stouff V, Geslin E, Debenay J-P and Lesourd M (1999) Origin of morphological abnormalities in *Ammonia* (foraminifera): studies in laboratory and natural environments *J Foraminiferal Res* **29** 152-70
- Stuut J-B W, Prins M A and Jansen J H F (2002) Fast reconnaissance of carbonate dissolution based on the size distribution of calcareous ooze on Walvis Ridge, SE Atlantic Ocean *Mar Geol* **190** 581-589
- Thiede J (1971) Variations in coiling ratios of Holocene planktonic foraminifera *Deep-Sea Res* **18** 823-831
- Thompson P R, Bé A W H, Duplessy J-C and Shackleton N J (1979) Disappearance of pink-pigmented *Globigerinoides ruber* at 120,000 yr BP in the Indian and Pacific Oceans *Nature* **280** 554-557
- Thunell R C and Honjo S (1987) Seasonal and interannual changes in planktonic foraminiferal production in the North Pacific *Nature* **328** 335-337
- Urey H C, Lowenstam H A, Epstein S and McKinney C R (1951) Measurement of paleotemperatures and temperatures of the Upper Cretaceous of England, Denmark, and the southern United States *Bull Geol Soc Am* **62** 399-416
- Weber M E, Niessen F, Kuhn G and Wiedicke M (1997) Calibration and application of marine sedimentary physical properties using a multi-sensor core logger *Mar Geol* **136** 151-172
- West S, Jansen J H F and Stuut J-B (2004) Surface water conditions in the Northern Benguela region (SE Atlantic) during the last 450 ky reconstructed from assemblages of planktonic foraminifera *Mar Micropaleontol* **51** 321-344
- Williams D F (1976) Late Quaternary fluctuations of the polar front and subtropical convergence in the southeast Indian Ocean *Mar Micropaleontol* **1** 363-375
- Williams D F, Healy-Williams N and Leschak P (1985) Dissolution and water-mass patterns in the Southeast Indian Ocean. Part I. Evidence from Recent to late Holocene foraminiferal assemblages *Geol Soc Am Bull* **96** 176-189
- Wright R G (1977) Planktonic-benthonic ratio in foraminifera as paleobathymetric tool. Quantitative evaluation. *Ann Am Assoc Pet Geol and Soc Econ Paleontol Mineral Conv, Washington, D C*, 842
- Yanko V, Ahmad M and Kaminski M (1998) Morphological deformities of benthic foraminiferal tests in response to pollution by heavy metals: implications for pollution monitoring *J Foraminiferal Res* **28** 177-200

## **Legend to Figures**

Figure 1: Calcareous benthic foraminifera from the north-eastern Indian Ocean. The white scale bar at the base of each figure equals 100  $\mu\text{m}$ .

Figure 2: Planktic foraminifera from the north-eastern Indian Ocean. A majority of planktic foraminiferal tests have globular chambers and are usually perforate. The white scale bar at the base of each figure equals 100  $\mu\text{m}$ .

Figure 3: Agglutinated benthic foraminifera from the north-eastern Indian Ocean. The white scale bar at the base of each figure equals 100  $\mu\text{m}$ .

Figure 4: A schematic of measuring the test size as well as the size of the proloculus. 'P' indicates the maximum diameter of the proloculus or the first formed chamber. The foraminiferal test should be kept in spiral view to measure the maximum diameter of the proloculus.

Figure 5: Dextral (A) and sinistral (B) coiling in a trochospirally coiled foraminifera.

Figure 6: A schematic of counting the total number of chambers in a test. The foraminiferal test should be kept in spiral view to count the total number of chambers.



**Figure 1**



**Figure 2**





Figure 3




Figure 4




Figure 5




Figure 6