

Evaluating changes in the elemental composition of micrometeorites during entry into the earth's atmosphere

N. G. Rudraswami^{1*}, M. Shyam Prasad¹, S. Dey^{1,2}, J. M. C. Plane³, W. Feng³, S. Taylor⁴

¹National Institute of Oceanography (Council of Scientific and Industrial Research), Dona Paula, Goa 403004, India

²Indian Institute of Technology, Roorkee, Uttarakhand 247667, India

³School of Chemistry, University of Leeds, Leeds LS2 9JT, UK

⁴Cold Regions Research and Engineering Laboratory, 72 Lyme Road, Hanover, New Hampshire 03755-1290, USA

Corresponding Author: rudra@nio.org

ABSTRACT

We evaluate the heating of extraterrestrial particles entering the atmosphere using the comprehensive Chemical ablation model (CABMOD). This model predicts the ablation rates of individual elements in a particle with a defined size, composition, entry velocity and entry angle with respect to the zenith (ZA). In the present study, bulk chemical analyses of 1133 Antarctica micrometeorites (collected from South Pole water well) are interpreted using CABMOD. The marked spread in Fe/Si values in unmelted, partially-melted and melted micrometeorites is explained by the loss of relatively volatile Fe during atmospheric entry. The combined theoretical modelling and elemental composition of the micrometeorites (Mg/Si ratios) suggest that ~85% of particles have a provenance of carbonaceous chondrites, the remaining ~15% are either ordinary or enstatite chondrites. About 65% of the micrometeorites have undergone <20% ablation, while a further 20% have lost between 20 and 60% of their original mass. This has implications for understanding the micrometeorite flux that reaches the Earth's surface, as well as estimating the pre-atmospheric size of the particles. Our work shows that the unmelted particles which contribute ~50% to the total micrometeorite collection on Earth's surface have a small entry zone: $ZA = 60-90^{\circ}$ if the entry velocity is ~11 km/s, and $ZA = 80-90^{\circ}$ for >11 - 21 km/s.

1. INTRODUCTION

The extraterrestrial dust entering the Earth's atmosphere ($\sim 30,000 - 40,000 \text{ t y}^{-1}$) is produced from comets and asteroids, and is a unique source of information on the formation processes of the early Solar System (Love and Brownlee 1993; Peucker-Ehrenbrink and Ravizza 2000; Plane 2012). The size of most of the dust ranges from a few μm to a few mm. A significant aspect of micrometeorite studies is that some of these particles may have sampled unfamiliar asteroids that are not represented in our meteorite collection (Yada et al. 2005; Gounelle et al. 2009). The dust particles are anticipated to originate pre-dominantly in the asteroid belt, further drift into the inner solar system with a large probability of getting trapped in orbital resonance with Earth due to their smaller orbital eccentricities compared to cometary particles (Dermott et al. 1994; Reach et al. 1997; Brownlee 2001). There is some noteworthy contribution from the cometary bodies based on Infrared Astronomical Satellite (IRAS) data, although the IRAS evidence is not very clear to arrive at accurate estimates (Dermott et al. 1994; 1996). Similarly, cosmic background explorer (COBE) observation suggested trapping of asteroidal dust dominantly, making cometary dust insignificant (Kortenkamp et al. 2001). The numerical model by Kortenkamp and Dermott (1998) estimated $\sim 25\%$ of dust reaching Earth is contributed by comets. Nesvorny et al. (2010, 2011) suggested comets of Jupiter family are major source of dust particles and contributes $>80\%$ of dust flux arriving to the Earth in contradiction to above estimates. In any circumstance, estimating the exact contribution of the asteroidal and cometary reservoirs to the total flux is challenging. The low orbital eccentricities of asteroidal dust compared to that of higher eccentricities of cometary dust gives an advantage of greater gravitational capture cross section to asteroidal particles making them as the largest contributor of extraterrestrial dust, probably followed by cometary dust (Flynn 1989b; Kortenkamp and Dermott 1998). The cometary particles travel at higher velocities encounter a small amount gravitational capture than their counterpart asteroidal particles (Flynn 1989b; Brownlee 2001). However, it is worthwhile to note that the cometary particles trapped in resonance with Jupiter will have similar behaviour to that of asteroid dust particles (Kortenkamp et al. 2001).

The bulk chemical composition of a meteoroid can change because of the rapid heating it experiences during atmospheric entry. Besides a relatively small loss of matter through sputtering with air molecules, the particle's temperature may reach its melting point and then lose a much larger fraction of mass through evaporation. Both of these processes are described by the term ablation (Vondrak et al. 2008). The degree of elemental change depends on the range of the heating event it undergoes, which in turn depends on the density, mass, chemical composition, entry velocity and

entry zenith angle (ZA) of the particle. Understanding the nature and extent of the changes in physical and chemical properties may provide insights into the source of the particle. For many decades, studies of the chemical composition did not make a quantitative correction for ablation in order to infer the original composition of the particle. Indeed, although the effects of heating due to variations in size, ZA and entry velocity were discussed (e.g. Fraundorf 1980; Flynn 1989a; Love and Brownlee 1991), the problem of understanding changes in the chemical composition during entry has not been a subject of extensive investigation.

The purpose of this work is to understand the changes in chemical composition during atmospheric heating by using the Chemical Ablation Model (CABMOD) developed by Vondrak et al. (2008). In particular, we will focus on changes to elemental ratios as a function of entry parameters. The chemical and mineralogical compositions of 1133 Antarctic micrometeorites will then be examined in order to infer the degree of mass loss and compositional alteration experienced during atmospheric entry, and hence the nature of the particles pre-entry.

2. SAMPLE COLLECTIONS

The samples were collected during 2000 from the South Pole Water Well (SPWW) at a depth of ~100 m below the snow surface. This melt-water well had a diameter of ~24 m, depth of ~15 m and a water volume of ~5000 m³ (Taylor et al. 1998, 2000). The micrometeorites collected from this well were dated to between 1100 and 1500 AD (Taylor et al. 1998). The polar ice offers a pristine environment for micrometeorite collection, where the melt water is filtered and the filtrates are examined for micrometeorites. Contaminants are very few in this environment and the compressed ice also provides a pre-concentration of the extraterrestrial particles from the large volumes of ice melted. The polar samples are better preserved and therefore less weathered compared to deep sea micrometeorite collections. The samples collected in different size ranges above 50 µm were sieved into various size fractions (Taylor et al. 1998). Using a binocular microscope ~1200 particles were sorted, mounted in epoxy and polished for further study using a scanning electron microscope and electron microprobe. There are standard parameters for identification of extraterrestrial particulate matter identified over a period of time (Blanchard et al., 1980; Brownlee et al., 1997; Taylor et al. 2000 and Genge et al. 2008) : spherical morphology or even non-spherical and irregular shapes for unmelted particles, presence of magnetite rims due to entry heating, Fe-Ni bead, platinum group nuggets, Mg-rich relict olivines, formation of vesicles, chemical composition, elemental ratios, mineralogical properties and absence of terrestrially abundant elements. The classification of the

micrometeorites is based on the textures and mineralogical composition as suggested in Genge et al. (2008).

3. ANALYTICAL TECHNIQUES

Back Scattered electron (BSE) images of the polished epoxy-mounted and carbon-coated micrometeorites were obtained using a JEOL JSM 5800LV SEM with an OXFORD INCA Energy Dispersive Spectrometer detector (EDS, ISIS-300 at the National Institute of Oceanography, Goa. The image is useful for classifying and identifying the proper phases for chemical analyses using a Cameca SX5 Electron Micro Probe Analyzer (EPMA, at National Institute of Oceanography, Goa) equipped with four spectrometers. The elemental compositions determined in the micrometeorites were Na, Mg, Si, Al, P, K, Ca, Ti, Cr, Mn, Fe and Ni. The operating conditions used were as follows: accelerating voltage: ~15 kV, beam current ~12 nA, beam diameter of ~1–2 μm . The data are corrected using the PAP model (Pouchou and Pichoir 1991). The analytical totals are less than 100% in some micrometeorites, this can be due to multiple reasons. The presence of the hydrous phases or porous contents, sub-micron void spaces, and cracks result in analytical deficiencies. However, care is taken by carrying out multiple analyses on the same spot. In any case, a small deviation from 100% can be considered to be within the permissible limits of the instrument analytical capabilities.

4. ABLATION MODEL DESCRIPTION

CABMOD (Vondrak et al. 2008) uses the classical physics equations to describe the time dependence of the meteoroid energy and velocity. The model includes sputtering by collision with air molecules before the particle melts, followed by Langmuir evaporative loss of metal atoms and oxides from the molten particles (treated by the thermodynamic model MAGMA) as a function of altitude (Schaefer and Fegley 2005). Particles are considered to be a single phase with a density of 2 g cm^{-3} , with a CI composition taken from multiple sources (Mason 1971; Sears and Dodd 1988; Lodders and Fegley 1998). The fundamental properties of density for the majority of unmelted extraterrestrial particles has been suggested to be 2 g cm^{-3} which is similar to that of the CI chondrites suggests that the particle are primitive (Love et al. 1994). In addition, the major element compositions of the micrometeorites studied by various groups are close to CI composition (e.g. Brownlee et al. 1997; Taylor et al. 2000; Prasad et al. 2013), therefore, the CI chondrites with density of 2 g cm^{-3} is an appropriate choice for the model calculation based on observations on large population of micrometeorites by many investigators.

In the present study we use CABMOD to quantify the elemental composition changes that particles undergo as a function of particle diameter, velocity and ZA. Differing combinations of these variables can produce similar compositional changes. The maximum entry velocity for a majority of the particles to survive atmospheric entry is ~ 16 km/s is for those particles that have size range from 100 to few hundred μm in size. The choice of this size is based on the population of the micrometeorites studied (e.g. Taylor et al. 2000; Prasad et al. 2013). If the velocity of the particle goes beyond 16 km/s the ablation can surpass $\sim 70\%$ (Table 1). However, for particles less than 100 μm in size, there is greater chance of survival even at higher velocities. For the present study, we have used the following boundary conditions: entry velocity of ≤ 16 km/sec, peak temperature the particle can attain during entry to < 2000 K, sizes ranging from ~ 100 μm to few hundred μm across and zenith angles of $\sim 45\text{--}90^\circ$. Love and Brownlee (1991) suggested that the increase in mass loss with increasing temperature reaches a zenith at 2000 K, which is suggested as the practical upper limit that any particle can reach during atmospheric entry. Even if a particle survives beyond this temperature, its chemical composition deviates due to large scale vaporization and tends to lose its characteristic features such as elemental ratios which are crucial for identification of precursors. Most of the micrometeorites recovered have size in the range of 100 to few hundred μm . However, model calculation suggest zenith angle $\sim 0\text{--}45^\circ$ will experience large heating and ablation depending on the velocity (discussed in section 6.1 and Table 1). Hence, we restrict ourselves to velocities ≤ 16 km/s and temperatures < 2000 K, as temperature above this will lead to elemental vaporization that can potentially alters the chemical composition, losing our understanding on the precursors composition (Table 1). It should be noted that the present work does not consider the atmospheric fragmentation of a particle during entry (Lal and Jull 2002).

5. MICROMETEORITE CHEMICAL COMPOSITION

We have classified 1133 Antarctica micrometeorites using BSE images. Although the mount was dominated by melted spherules, there were 11 particles which can be classified as unmelted micrometeorites. The remaining micrometeorites are classified as follows: 42 are relict-grain bearing, 57 are scoriaceous, 355 are porphyritic, 137 are barred, 119 are cryptocrystalline 371 are glass, 18 are I-type spherules and 23 are G-type spherules. Among the porphyritic, a large number of spherules had relict grains with sizes > 50 μm , indicative of preservation during atmospheric entry. Olivine is the most abundant relict mineral, dominated by a fosteritic composition; however, some minor minerals such as pyroxene, spinel, and plagioclase are also found. Fig.1 shows representative BSE images of the unmelted and relict-bearing micrometeorites. Figs. 1a-d are unmelted single

mineral grains with uniform chemical composition. Fig. 1e is an example of a multiple mineral micrometeorite. Figs. 1f-h are micrometeorites that experienced partial heating during atmospheric entry, but largely preserved their relict mineral compositions. P160 shows the presence of a large number of Mg-rich fosterites along with pyroxenes (Fig. 1i). Large relict-grain bearing micrometeorites are seen in Figs. 1j-l. Table 2 lists the bulk chemical composition for the unmelted and relict-bearing micrometeorites. The unmelted micrometeorites should preserve chemical compositions similar to those of their parent bodies. Spinel is one of the refractory minerals which is rarely found in micrometeorites, though commonly found in calcium Aluminum-rich Inclusions (CAIs). Previously, some evidence of spinel earlier in micrometeorites/cosmic spherules has been reported (Taylor et al. 2008, 2012). In the present study, spinels were observed in four spherules, whose BSE images are shown in Fig. 2 and chemical composition listed in Table 3.

Multiple analyses by electron microprobe on different phases and compositions are helpful in finding the bulk chemical composition of micrometeorites. The Lispix software program (<http://www.nist.gov/lispix/>) produces phase images where modal proportion of the volume is calculated (Berlin 2009). This method to determine the bulk chemical composition is more accurate than a simple average of electron microprobe data, or the defocussed beam technique where the broad beam frequently hits a void space or vesicle, leading to low total counts of chemical composition. The bulk chemical compositions of the cosmic spherules are similar to those found in the deep sea sediments and Polar Regions (Brownlee et al. 1997; Taylor et al. 2000; Rudraswami et al. 2012, 2014; 2015; Prasad et al. 2013). The depletion of volatile elements such as Na and S, followed by Fe, is evident during atmospheric heating. A plot of Ca/Si versus Al/Si follows along the solar composition line (Fig. 3). The glass, barred olivine and cryptocrystalline spherules are essentially close to the solar line; however, there is some spread in other micrometeorites (e.g. scoriaceous and porphyritic). The Mg/Si ratios exhibit a spread from 0.5–1.5 (Fig. 4a), although dominated by the CI composition ratio irrespective of the type of micrometeorite. The same behaviour of Mg/Si is also seen in the case of the deep sea spherules (Brownlee et al. 1997; Prasad et al. 2013). Fe is more volatile than Mg (which is bound in the melt as a stable oxide), and its faster evaporation causes a much greater spread in the Fe/Si ratio (Fig. 4b), in contrast to Mg/Si. The Fe/Si is spread much lower than CI value. The peak at Fe/Si >2 is due to the Fe-rich composition in Scoriaceous, porphyritic, cryptocrystalline and barred micrometeorites. The behaviour of Fe/Si is slightly different in the deep sea spherules, probably because of a magnetic bias in the collection which causes the collected spherules to be richer in Fe (Brownlee et al. 1997).

The Al/Si (Fig. 4c) and Ca/Si (Fig. 4d) ratios are in good agreement with their CI values, unlike Fe/Si. Si is more volatile than Ca and Al, leading to depletion of Si during entry (Hashimoto 1983; Vondrak et al. 2008) and a second peak of Al/Si and Ca/Si. The Mg-Fe-Si ternary diagram (Fig. 5) for 1110 micrometeorites (excluding G-type) along with the trend of extraterrestrial particles and terrestrial particles suggests that different types of collections from deep sea, Antarctica or Greenland have similar chemical composition ranges (Brownlee et al. 1997; Taylor et al. 2000; Rochette et al. 2008).

6. DISCUSSION

6.1. Chemical variation of particles during atmospheric entry

The present study focuses on understanding the changes in the elemental composition of the incoming particles as a function of size, ZA and velocity. Here we examine the major elements such as Si, Mg, Fe, Ca and Al which contribute more than 90% of the particle mass (along with O). For the present the parameters considered will be discussed in section 6.3. For. e.g., considering the above constraints, for the minimum entry velocity of ~ 11 km/s, ZA $\sim 0^\circ$ and particle size ~ 100 μm , the peak temperature achieved is ~ 1700 K where the percent of ablation is close to ~ 1 %. This percent of mass ablation and the peak temperatures achieved by the particles increases rapidly, primarily with entry velocity and ZA but also as a function of the mass of the incoming particle. For two ZAs of 45° and 60° with different velocities and sizes, it can be seen in Fig. 6 that the peak temperature of the particles exceeds 2000 K even for 200 μm particles, at velocities greater than 16 km/s. The particles that have crossed 2000 K have greater chances to undergo large scale changes in chemical composition by vaporization of elements, thereby losing distinctive features of the precursors (Love and Brownlee 1991).

At an entry velocity of 11 km/s, a large mass of the particles is preserved irrespective of ZA with maximum ~ 30 % ablation (Table 1). However, as the entry velocity increases, ablation increases for all ZA apart from near-horizontal entry where ZA = 80 – 90° (Fig. 7 and Table 1). The chances of survival of the particles without any disturbance are highest for those that enter in the narrow window of ZA = 80 – 90° and entry velocity < 21 km/s. This is true for ZA 60 – 90° and an entry velocity 11 km/s (Fig. 6 and Table 1). Almost 50% of the micrometeorites collected from the Greenland ice cap are unmelted, especially those in the size range of 50–100 μm (Maurette et al. 1987). Yada et al. (2004) found that nearly 60% of the particles collected around the Yamato Mountains in Antarctica are unmelted, while the remaining are cosmic spherules. The SPWW

micrometeorite collection also found that roughly half of the particles collected were unmelted in the size range of 50–100 μm (Taylor et al. 2007a). The Antarctic and Greenland Ice collections therefore suggest that unmelted micrometeorites and melted spherules are approximately equal in number, unlike deep sea spherules (Taylor and Brownlee 1991; Brownlee et al. 1997; Rudraswami et al. 2012; Prasad et al. 2013). There is a narrow zone of ZA and velocity which will allow a meteoroid with a diameter greater than 100 μm to enter the atmosphere without melting and altering the mineralogical features or texture: $\text{ZA} = 80\text{--}90^\circ$ and entry velocity $< 21 \text{ km/s}$. The $\text{ZA } 80\text{--}90^\circ$ is the zone where nearly 50% of the total recovered particles in the size greater than 100 μm reaches earth surface survive ablation and are well preserved. However, if the velocity of the entering particle is $\sim 11 \text{ km/s}$, then $\text{ZA } 60\text{--}90^\circ$ is the minimum ablation zone for any size less than few hundred microns across as sputtering will always occur in unmelted particles also.

The elemental changes that the particles undergo are observed largely in Mg, Fe, Si which contribute more than 90% of the particle mass in the oxide form as the major silicate minerals, followed by the refractory elements Ca and Al. Fe is the most volatile of these elements (after Na and K) and after particle melting contributes most to the ablative loss, followed by Si and Mg while there is no major ablation of Ca and Al (Fig. 8). For example, for a $\text{ZA} = 45^\circ$, diameter $\sim 200 \mu\text{m}$ and entry velocity $\sim 16 \text{ km/s}$, Fe decreases from 28 wt% to $< 8 \text{ wt\%}$ by the time at the particle descends to 70 km altitude, while the loss in Fe results in enrichment of Mg and Si from $\sim 15\%$ and 16% to 25% and 19% , respectively (Fig. 9). The jogs in Fig. 9 are the regions where the ablation begins to occur at the altitude of 95-101 km. The first element to be ablated is Na followed by K. As a result, the particle loses mass, but not of Mg, Si or Fe, so there seems to be a slight enrichment of these elements compared to their initial composition. For any given size, a smaller ZA and larger velocity give rise to a greater fraction of mass loss through elemental ablation (Table 1). As the ablative fraction increases, the Fe/Si ratio decreases: 20% ablation results in a 33% decrease of the Fe/Si ratio, and once ablation is higher than 70% the particle has lost more than 90% of its Fe (Fig. 10b). The Mg/Si ratios (Fig. 10a) exhibit a reverse trend compared to Fe/Si. The Mg/Si ratios increase with mass ablation until 80 % of the mass is gone, at which point the ratio decreases. There is little change in the Mg/Si ratio up to $\sim 20\%$ ablation. A higher degree of ablation will potentially alter the mineralogical composition due to loss of Fe followed by Si and Mg. In contrast, the refractory elements such as Ca and Al do not evaporate significantly even after 50% ablation, at which point Si evaporation causes a substantial increase in the Ca/Si ratio (Fig.10c). The Al/Si ratio behaves in a similar way (Fig.10d). The slow vaporization of Ca and Al can be useful for identifying the

precursors from Ca/Si and Al/Si ratios. The disadvantage is the difficulty involved in considering the chemical composition of the particle which does not represent the bulk mineralogical of the precursors. Small fragments of dust generated by disruption during asteroidal collisions may be of individual mineral or group of minerals and far from the bulk mineralogical composition of the parent bodies.

Various methods to determine the mass accretion of extraterrestrial particles has provided a wide range of estimates (Plane 2012). The analysis of micro-impact craters on the space-facing side of the Long duration exposure facility (LDEF) satellite at an altitude of 350–480 km constrains the total accreted mass to be 30,000 t y⁻¹ with a large uncertainty of ±20,000 t y⁻¹; however, this estimate requires an assumed velocity distribution of the particles (Love and Brownlee 1993; Taylor et al. 1998). Os isotopes studies in marine sediments have indicated an accretion rate of 30,000±15,000 tons y⁻¹ which is similar to that of the LDEF satellite estimate (Peucker-Ehrenbrink and Ravizza 2000). The accumulation of Ir, Pt and super-paramagnetic Fe particles in polar ice cores also imply a large dust flux of at least 36,000 t y⁻¹ (Gabielli et al. 2004; Lanci et al. 2007). In contrast, the sampling of cosmic spherules from the SPWW gave an estimate of 2700±1400 t y⁻¹ (Taylor et al. 1998). The accretion rate of micrometeorites in blue ice was estimated to be 16,000±9000 t y⁻¹, based on the collective analyses of handpicked particles and noble gas measurements done of the residual particles (Yada et al. 2004). The Greenland ice micrometeorite collection yields an estimate of ~4000 t y⁻¹ (Maurette et al. 1987). The mass accretion rate estimated by counting cosmic spherules in deep sea sediments yielded a low estimate, as most of the fragile particles seems to get destroyed in the harsh environment (Prasad et al. 2013). Each of these estimation techniques depends on various assumptions and is then scaled up to provide the global input (Plane 2012). In any case, a significant fraction of the meteoroid flux entering the atmosphere ablates. The resulting metal vapours then oxidize and polymerize to form meteoric smoke particles (Plane et al. 2015), which is the major form of extra-terrestrial material sampled in ice cores. For meteoroids with ZA = 0 to 50° and velocity >16 km s⁻¹, at least 70% of the particle mass is vaporized during entry. However, for an entry velocity <21 km s⁻¹ and ZA = 50–70°, a particle will melt and form a cosmic spherule (such as porphyritic, barred, cryptocrystalline, glass). ZA = 70–80° is the zone where relict-grain bearing, scoriaceous spherules are partially heated and transported to the Earth's surface. As discussed earlier, unmelted micrometeorites have a very narrow window of ZA = 80–90° for velocity ~16–21 km s⁻¹, or 60–90° for ~11 km s⁻¹.

6.2. Micrometeorites and their likely precursors

The mineralogy of the micrometeorites is important for gaining information on the precursors they originated from. CABMOD is useful to some extent for extrapolating from the micrometeorite back to the original mass and nature of the meteoroid from which it formed. As shown in the ternary diagram (Fig. 5), the micrometeorites in the present study fall within the limit of the extraterrestrial line, similar to those seen in other studies of Antarctic micrometeorite collections obtained either by melting ice or traps (Taylor et al. 2000; Rochette et al. 2008). The diffusion and evaporation rate of elemental Fe is much faster than that of Mg (Hashimoto 1983), which explains the occurrence of magnetite rims formed on micrometeorites during atmospheric entry. A thin Fe-rich rim is common even on unmelted particles or single minerals, suggesting Fe fractionating in particles takes place even at relatively low temperatures. The spread in the Fe/Si ratio in Fig. 4b suggests that the process of diffusion or evaporation of Fe leads to the rapid depletion of Fe/Si, since there are a comparatively small number of micrometeorites near the CI value. An additional reason may be because of a large precursor that is dominated by chondrules that have low Fe/Si values.

The Mg/Si values are largely close to the CI chondrite value (~0.8-1.0), within the analytical error involved in estimating the bulk chemical composition of the micrometeorites. The number of micrometeorites studied here makes them appropriate for statistical interpretation. Nearly 65% of the micrometeorites from the present study have Mg/Si values ranging from 0.8 to 1.0, irrespective of the texture or type. CABMOD suggests that the range will have $\leq 20\%$ of ablation for an entry velocity less than 16 km/s and ZA greater than 45° marked as region I in Fig. 11. For other lower velocities and similar ZA the ablation is much less than 20% and close to CI values and fall in region I. On the other hand, 20% of micrometeorites have bulk Mg/Si in the range of 1.0–1.4 have ablation from 20–60% suggesting large alteration during the entry (marked as region II in Fig. 11). Above this range the ablation is very high with large scale vaporization of Fe and Si, and any particle that has large dimension may not be able to retrievable as a micrometeorite on the Earth's surface (marked as region III in Fig. 11). For velocity 11 km/s the ablation is less than 20% and will lie in region I. However, ~15% of particles that do not fall in these three regions are those with bulk Mg/Si values that range between ~0.6 to 0.8. It should be noted that the Mg/Si ratio becomes enriched during heating due to Si evaporation which is much faster than Mg. A Mg/Si ratio between ~0.6 and 0.8 therefore suggests a different precursor from carbonaceous chondrites, such as ordinary chondrites which have Mg/Si values of ~0.8, EH ~0.6 and EL~0.7 (Lodders and Fegley 1998). Thus it appears that ~15% of the total particle that survive entry have a provenance to ordinary or enstatite

chondrites, while ~85% are related to carbonaceous chondrites. Among all global collections of cosmic dust (Taylor et al. 2000, 2007b; Prasad et al. 2013) it is seen that a majority of the particles are sourced either from CI or CM chondrites, which can be distinguished from ordinary chondrites (Brownlee et al. 1997). Engrand and Maurette (1998) suggested that up to 99% of the particulate flux of cosmic matter is dominated by either carbonaceous or cometary bodies, this is especially true for unmelted particles. The CI and the CM chondrites being more porous and hydrated, fragment easily during asteroidal collisions and produce greater quantities of dust sized materials (Tomeoka et al. 2003) which stream into the inner solar system. Whereas the anhydrous meteorites (ordinary chondrites and others) have greater tensile strengths and do not breakdown that easily into the smallest particles during comminution in the asteroid belt, therefore their presence is not dominant among micrometeorites. On the basis of their elemental ratios and textures, the micrometeorites analysed in the present investigation also suggest a dominant contribution from carbonaceous chondrites.

6.3. Precursors for Refractory minerals

The spinels are refractory minerals with melting temperatures above 2000 K. In the present study only four micrometeorites contain spinels, which are occasionally found in the CAIs of carbonaceous chondrites. The oxygen isotope composition of the spinels in the Antarctic micrometeorites shows they are ^{16}O -rich, similar to those found in CAIs which suggests refractory inclusion (Taylor et al. 2012). P219 is nearly pure anorthitic in composition (An_{90-94}), with few chromite grains enclosed. These types of anorthitic rich spherules could have originated from the CAIs of CV, CO, CH and CR chondrites (Krot et al. 2002). The number of micrometeorites with a CAI component is less than 0.5%, which is far from CM, CO, CV, CK chondrites which have ~1.2, 1.0, 3.0 % modal abundances (Hezel et al. 2008). One of the primary reasons for a poor contribution from refractory minerals towards the flux of micrometeorites is the very fluffy nature: the loosely packed assemblage with a large void space easily disintegrates during collisions either in the asteroidal belt or during atmospheric entry (Lal and Jull 2002). Only fragments that have similar mineral assemblages have a greater chance of survival.

7. CONCLUSIONS

A chemical ablation model (CABMOD) has been used to evaluate the effect of meteoroid size, velocity and ZA on the degree of ablation and chemical variation of particles entering the atmosphere. We describe the analyses of 1133 micrometeorites collected from the South Pole Water

Well. We predict, based on the measured Mg/Si ratios of the micrometeorites and the CABMOD analysis, that ~15% of recovered samples are either ordinary or enstatite chondrites as precursors, while the rest have a carbonaceous chondritic origin. The unmelted micrometeorites that contribute half of the total flux that reaches Earth's surface have a very narrow ZA of 60–90° and 80–90° for entry velocity ~11 km/s and >11–21 km/s. In spite of the fact that the micrometeorites recovered on the surface have often undergone large scale alteration during entry, a significant number of micrometeorites (~65%) have Mg/Si ratios ~0.8–1.0 suggesting modest ablation of less than 20%, while another 20% of the particles have undergone 20–60% ablation. This quantification of ablation is important for estimating the micrometeorite precursor sizes and flux.

Acknowledgements

This work is supported by the CSIR XII Plan funded Project GEOSINKS and the PLANEX project, Physical Research Laboratory, Ahmedabad (NGR and MSP). This is NIO's contribution No. xxxx. The work at the University of Leeds is supported by the European Research Council (project number 291332 - CODITA)

REFERENCES

- Berlin J. 2009, PhD Thesis, University of New Mexico, Albuquerque, New Mexico, USA.
- Blanchard M. B., Brownlee D. E., Bunch T. E., Hodge P. W. & Kyte F. T. 1980, *EPSL* 46, 178
- Brownlee D. E. 2001, *Acretion of Extraterrestrial Matter Throughout Earth's History* (eds. B. Peucker-Ehrenbrink and B. Schmitz), pp. 1–12. Kluwer/Plenum, New York.
- Brownlee D. E., Bates B. & Schramm L. 1997, *M&PS* 32, 157
- Dermott S.F., Jayaraman S, Xu. Y. L., Grogan K. & Gustafson B. A. S. 1996, *AIP conference proceedings*, 348, 25
- Dermott S.F., Jayaraman S, Xu. Y. L., Gustafson B. A. S. & Liou J. C. 1994, *Nature* 369, 719
- Flynn G.J. 1989a, *Proc. LPSC*. 19, 673
- Flynn G.J. 1989b, *Icarus* 77, 287
- Fraundorf P. 1980. *GRL*, 7, 765
- Gabrielli, P., Barbante, C., Plane, J. M. C., Varga, A., Hong, S., Cozzi, G., Gaspari, V., Planchon, F. A. M., Cairns, W., Ferrari, C., Crutzen, P., Cescon P. and Boutron, C. F. 2004, *Nature*, 432, 1011
- Genge M. J., Engrand C., Gounelle M. & Taylor S. 2008, *M&PS*, 43, 497

- Gounelle M., Chaussidon M., Morbidelli A., Barrat J. A., Engrand C., Zolensky M. E. & McKeegan K. D. 2009, PNAS, 106, 6904
- Hashimoto A. 1983, *Geochem. J.* 17, 111
- Hezel D. C., Russell S. S., Ross A. J. & Kearsley A. T. 2008, M&PS, 43, 1879
- Kortenkamp S. J. & Dermott S. F. 1998, *Icarus* 135, 469
- Kortenkamp S. J., Dermott S. F., Fogle D., & Grogan K. 2001, *Acretion of Extraterrestrial Matter Throughout Earth's History* (eds. B. Peucker-Ehrenbrink and B. Schmitz), pp. 13–30. Kluwer/Plenum, New York.
- Krot A. N. & keil K. 2002, M&PS, 37, 91
- Lal D. and Jull A. J. T. 2002, APJ, 576, 1090
- Lanci L., Kent D. V. & Biscaye P. E. 2007, GRL, 34, L10803, doi:10.1029/2007GL029811.
- Lodders K. & Fegley B. Jr. 1998, *The Planetary Scientist's Companion*. Oxford University Press, New York.
- Love S. G. & Brownlee D. E. 1991, *Icarus* 89, 26–43.
- Love S. G. & Brownlee D. E. 1993, *Science* 262, 550
- Mason, B. (1971) *Handbook of Elemental Abundances of the Elements in Meteorites*, Gordon and Breach, Newark, USA.
- Maurette, M., J'ehanno C., Robin E., and Hammer C. 1987, *Nature*, 328, 699
- Nesvorný, D., P. Jenniskens, H. F. Levison, W. F. Bottke, D. Vokrouhlický, & M. Gounelle 2010, APJ, 713(2), 816
- Nesvorný, D., D. Janches, D. Vokrouhlický, P. Pokorný, W. F. Bottke, & P. Jenniskens (2011), APJ, 743 (2), 129
- Nicol E. J., Macearlane J. and Hawkes R. L. 1985, PSS, 33, 315
- Peucker-Ehrenbrink B. and Ravizza G. 2000, GCA, 64, 1965
- Plane J. M. C. 2012, *Chem. Soc. Rev.* 41, 6507
- Plane, J. M. C., W. Feng; Dawkins, E. C. M. 2015, *Chemical Reviews*, 115, 4497
- Pouchou J. L. and Pichoir F. 1991, *In Electron Probe Quantification* (eds. K. F. J. Heinrich and D. E. Newbury). Plenum Press, New York, pp. 31–75.
- Prasad M. S., Rudraswami N. G. & Panda D. K. 2013, JGR, 118, 2381
- Reach W. T., Franz B. A, Weiland J. L. 1997, *Icarus* 127, 461–484.

- Rochette P., Folco L., Suavet C., van Ginneken M., Gattacceca J., Perchiazzi N., Braucher R. & Harvey R. P. 2008 PNAS, 105, 18206
- Rudraswami N. G., Prasad S. M., Plane J. M. C., Berg T., Feng W. & Balgar S. 2014, GCA, 131, 247–266.
- Rudraswami N. G., Shyam Prasad M., Nagashima K., & Jones R. H. 2015, GCA, 164 , 53
- Rudraswami N. G., Shyam Prasad M., Babu E. V. S. S. K., Vijaya Kumar T., Feng W. & Plane J.M.C. 2012, GCA, 99, 110
- Schaefer, L., and B. Fegley (2005), EMP, 95(1-4), 413
- Sears, D. W. and Dodd, R. T. 1988 in: Meteorites and the Early Solar System, edited by: Kerridge, J. F. and Matthews, M. S., University of Arizona Press, Tucson, 3–31
- Suavet, C., Gattacceca J., Rochette P., Perchiazzi N., Folco L., Duprat J., and Harvey R. P. 2009, JGR, 114, B04102, doi:10.1029/2008JB005831
- Taylor S, Alexander C. M. O'D. & Wengert S. 2008, LPSC, abstract #1628
- Taylor S. and Brownlee D. E. 1991, Meteoritics 26, 203
- Taylor S., Lever J. H. & Harvey R. P. 2000, M&PS, 35, 651
- Taylor S., Matrajt G. & Guan Y. 2012, M&PS, 47, 550
- Taylor, S., Lever, J. H. & Harvey, R. P. 1998, *Nature* 392, 899
- Taylor S., Matrajt G., Lever, J. H., Joswiak D. J., & Brownlee D. E. 2007a, Proc. 'Dust in Planetary Systems', Kauai, Hawaii, USA. 26--30 September 2005 (ESA SP-643, January 2007).
- Taylor S., Herzog G. F. & Delaney J. S. (2007), M&PS, 42, 155
- Tomeoka K., Kiriya K., Nakamura K., Yamahana Y. & Sekine T. 2003, *Nature* 423, 60
- Vondrak T., Plane J. M. C., Broadley S. & Janches D. 2008, *Atmos. Chem. Phys.* 8, 7015
- Yada T., Nakamura T., Noguchi T., Matsumoto N., Kusakabe M., Hiyagon H., Ushikubo T., Sugiura N., Kojima H., & Takaoka N. 2005, GCA, 69, 5789
- Yada T., Nakamura T., Takaoka N., Noguchi T., Terada K., Yano H., Nakazawa T., & Kojima H. 2004, EPS, 56, 67

Figures caption

Figure 1.

Back-scattered electron images of representative unmelted and relict-bearing micrometeorites collected from South Pole Water Well (SPWW). P743, P730, P1029 and P339 are unmelted single mineral micrometeorites. Single mineral micrometeorite P1029 has also been reported earlier by Taylor et al. (2012). P372 is a multiple mineral particle with a large portion dominated by pyroxene and some olivine; the dark grey region is Al-rich feldspathic glass. P728, P261, P160 and P44 have mixtures of olivine and pyroxene. P1042 exhibits a large relict grain. P504 and P1028 exhibit large multiple relict minerals. The bulk chemical compositions of the unmelted and relict-bearing micrometeorites are listed in **Table 2**.

Figure 2.

Four spinel-bearing micrometeorites. Micrometeorites P583 and P440 have also been reported earlier by Taylor et al. (2008, 2012). The chemical compositions of the spinel and the relict minerals of the micrometeorites are given in **Table 3**.

Figure 3.

Ca/Si versus Al/Si for different types of micrometeorites collected from the South Pole Water Well (SPWW). The solid line indicates the solar line.

Figure 4.

A histogram plot of elemental ratio for different types of micrometeorites collected from South Pole Water Well (SPWW): a) Mg/Si, b) Fe/Si, c) Ca/Si and d) Al/Si. The dashed vertical line is the CI composition (Lodders and Fegley, 1998). The peaks of Mg/Si, Ca/Si and Al/Si are similar to the CI composition, apart for Fe/Si which exhibits a large spread.

Figure 5.

Ternary diagram representing Mg-Si-Fe (atoms) of different types (unmelted: 11, relict bearing: 42, scoriaceous: 57, porphyritic: 355, cryptocrystalline: 119, barred olivine: 137, glass: 371, I-type: 18) of micrometeorites (total: 1110) collected from the South Pole Water Well (SPWW). The elemental ranges of micrometeorites and terrestrial particles suggested by Suavet et al. (2009) are indicated by the dark grey line and dotted red line, respectively.

Figure 6.

The distribution of temperature for different size of entry particle over a range of entry velocities from ~ 11 to 31 km s^{-1} for $ZA = 45^\circ$ and 60° . It can be seen that the temperature exceeds 2000 K , even for a velocity of only 16 km s^{-1} and a diameter of a few hundred μm .

Figure 7.

Percentage ablation for different particle sizes and ZAs, at entry velocities of 11 and 16 km s^{-1} . Note that unmelted particles have a narrow zone of ZA at 16 km s^{-1} , which broadens at 11 km/s . The number indicated is the percentage ablation (e.g. ablation is 95% for $ZA = 0^\circ$, velocity = 16 km s^{-1} , and size = $400 \mu\text{m}$).

Figure 8.

The contribution of individual elements to the percentage mass ablated. Fe ablates more rapidly than Si and Mg, thereby contributing more to the total mass ablated. The contribution from refractory elements Ca and Al is very low.

Figure 9.

Change in elemental percent versus altitude. At 80 km the ablation of a $200 \mu\text{m}$ particle at $ZA 45^\circ$ and entry velocity 16 km s^{-1} is fast; the resulting loss of Fe leads to an increase in the bulk Mg and Si composition of the particle.

Figure 10.

Change in elemental ratio with respect to the percent of mass ablated, for particles of various diameters, ZAs and entry velocities: a) Mg/Si, b) Fe/Si, c) Ca/Si and d) Al/Si.

Figure 11.

Variation of the Mg/Si ratio with percent of mass ablated, for different ZA at velocity = 11 and 16 km s^{-1} . The variation Mg/Si in the $0.8\text{--}1.0$ range, combined with the histogram plot of Fig. 4, suggests that $\sim 65\%$ of the particles undergo $\sim 20\%$ ablation (region I). Mg/Si $\sim 1.0\text{--}1.4$ suggests that $\sim 20\%$ of particles undergo $20\text{--}60\%$ ablation (region II). Only 3% of particles which undergo more than 60% ablation survive atmospheric entry (region III).

Table 1. The percent of mass ablated for various sizes, zenith angles and entry velocities.

Entry velocity (km/s)	Size (μm)	Percent of mass ablated									
		ZA=0°	10°	20°	30°	40°	50°	60°	70°	80°	90°
11	100	1	1	0	0	0	0	0	0	0	0
	200	3	3	3	2	2	1	1	0	0	0
	300	16	15	13	10	7	4	2	0	0	0
	400	33	32	29	25	18	11	5	0	0	0
16	100	22	21	18	14	9	5	2	1	0	0
	200	76	75	72	67	59	45	26	8	0	0
	300	93	93	92	89	85	77	62	35	0	0
	400	96	95	95	95	94	90	80	58	0	0
21	100	90	89	87	83	76	64	43	14	0	0
	200	97	97	97	96	96	95	91	71	0	0
	300	97	97	97	97	97	97	96	93	0	0
	400	98	98	98	97	97	97	97	96	0	0
26	100	97	97	97	97	96	95	92	72	16	0
	200	97	97	97	97	97	98	97	96	77	0
	300	98	98	98	98	98	97	97	97	95	0
	400	98	98	98	98	98	98	97	97	96	0
31	100	98	98	98	99	98	98	97	95	65	0
	200	98	98	98	98	98	97	97	98	96	0
	300	98	98	98	98	98	98	98	97	97	0
	400	98	98	98	98	98	98	98	98	98	0

Table 2. Bulk chemical composition (wt%) of unmelted and relict bearing micrometeorites.

Micrometeorites	Na ₂ O	MgO	Al ₂ O ₃	SiO ₂	CaO	Cr ₂ O ₃	MnO	FeO	CoO	NiO	Total
a) Unmelted											
P339	–	34.9	–	37.6	0.2	0.3	0.7	24.1	–	–	97.8
P46	0.1	33.8	1.1	39.1	0.7	0.3	0.4	23.9	–	0.3	99.7
P561	0.8	30.9	1.5	47.4	1.2	0.5	0.7	12.9	–	–	96.3
P730	–	41.7	–	38.8	–	–	0.4	18.2	–	–	99.2
P743	–	53.7	0.1	41.7	0.2	0.3	0.1	2.3	–	–	98.6
P1029	0.1	30.3	0.7	37.2	0.7	0.3	0.4	28.3	–	0.1	98.0
P1042	–	26.5	0.4	34.5	0.5	0.3	0.3	34.1	0.1	0.4	97.1
P29	0.8	26.2	5.7	45.7	3.7	0.5	0.8	14.7	–	0.1	98.3
P261	–	43.3	2.4	46.8	1.7	0.5	0.1	3.7	–	0.2	98.8
P372	1.5	25.0	4.5	52.8	0.9	0.1	0.3	9.7	–	–	95.2
P727	0.1	29.6	0.2	36.0	0.4	0.2	0.5	32.1	–	0.1	99.1
b) Relict bearing											
P44	–	32.9	1.5	44.0	1.0	0.5	0.2	18.3	–	0.1	98.6
P59	–	34.1	1.3	36.8	1.7	0.3	0.2	23.6	–	0.6	98.6
P88	–	39.4	1.3	45.0	0.5	0.4	0.2	11.2	–	0.4	98.4
P95	0.2	26.7	1.3	36.8	1.5	0.4	0.3	28.0	–	0.2	95.4
P106	0.1	23.0	0.6	52.1	0.7	0.5	0.6	19.7	–	0.1	97.5
P108	0.1	27.7	1.6	43.6	1.2	0.5	0.4	19.6	–	1.2	96.0
P109	0.1	27.0	1.2	40.5	1.6	0.7	0.2	23.0	–	0.6	95.0
P128	0.9	27.9	4.0	55.1	2.2	0.6	0.4	6.7	–	0.1	98.1
P160	0.3	37.2	2.8	44.4	1.6	0.7	0.2	9.4	–	0.3	96.8
P167	–	29.6	1.2	41.2	1.1	0.6	0.5	23.5	–	0.1	97.9
P220	0.5	22.8	3.6	42.9	3.6	0.3	0.3	23.4	–	0.2	97.7
P251	1.0	27.8	2.0	57.7	1.4	0.5	0.4	6.9	–	–	98.0
P302	–	30.3	1.3	44.4	1.2	0.5	0.5	18.3	–	0.1	96.8
P304	0.3	25.3	1.6	37.1	1.5	0.6	0.4	29.4	–	0.1	96.4
P351	–	33.4	0.8	37.7	0.7	0.3	0.3	24.4	–	–	97.7
P363	0.2	31.1	2.1	46.4	1.7	0.4	0.2	13.4	–	0.5	96.0
P392	–	55.4	0.1	40.1	0.4	0.2	0.1	0.9	–	–	97.4
P408	–	26.2	3.5	35.0	4.0	0.3	0.2	27.1	0.1	0.1	96.5
P415	0.1	23.8	3.8	43.9	3.8	0.4	0.6	20.3	–	0.1	96.8
P419	0.1	28.9	3.7	44.4	2.8	0.3	0.4	17.4	–	0.2	98.3
P504	–	34.3	0.6	36.7	0.7	0.4	0.7	24.9	–	0.1	98.4
P522	–	51.1	0.5	41.6	1.4	0.2	0.2	3.4	–	0.2	98.5
P541	0.2	30.4	1.7	39.4	1.4	0.2	0.5	22.7	0.1	0.3	96.9
P544	–	33.6	1.8	38.8	1.4	0.2	0.2	22.0	–	1.0	99.1
P586	–	24.9	1.4	49.8	1.1	0.5	0.4	18.7	0.1	0.5	97.4
P607	–	34.7	1.8	45.9	1.5	0.4	0.2	12.8	–	0.3	97.6
P627	0.1	30.9	1.7	49.1	3.2	0.4	0.2	11.0	–	0.5	97.2

P645	0.2	34.2	1.0	40.5	3.3	0.0	0.3	15.3	–	0.1	95.2
P668	–	27.2	2.3	38.3	2.3	0.4	0.2	26.8	–	0.4	98.0
P700	0.4	26.1	2.0	37.6	1.5	0.2	0.4	29.3	–	0.3	97.8
P703	0.5	27.1	1.7	36.6	1.3	0.3	0.4	30.0	–	0.1	98.1
P710	–	33.2	1.7	37.9	1.6	0.1	0.2	22.4	–	0.5	97.7
P721	0.4	29.9	2.1	43.6	2.0	0.3	0.5	19.7	–	0.1	98.7
P728	–	44.1	1.1	45.3	0.6	0.5	0.2	6.8	–	0.1	98.7
P791	1.9	22.1	2.1	43.6	1.6	0.2	0.6	25.4	–	–	97.7
P913	0.2	28.0	0.7	50.9	0.9	0.2	0.4	16.8	–	–	98.1
P930	0.1	31.9	0.4	38.0	0.6	0.2	0.3	28.0	–	0.1	99.6
P931	–	42.0	0.2	38.9	0.2	0.3	0.9	16.3	–	0.1	98.9
P939	–	43.3	1.8	46.1	1.2	0.5	0.3	4.5	–	0.2	98.1
P1028	–	30.4	1.5	37.1	1.4	0.3	0.5	24.4	–	–	95.7
P1162	0.5	32.9	1.4	49.2	1.0	0.3	0.3	10.2	–	0.9	96.7
P1209	0.2	24.3	3.3	38.2	2.8	0.3	0.3	27.4	0.1	0.2	97.2

Table 3. Chemical composition (wt%) of spinel found in micrometeorites.

Micrometeorites	Na ₂ O	MgO	Al ₂ O ₃	SiO ₂	CaO	TiO ₂	Cr ₂ O ₃	MnO	FeO	CoO	NiO	Total
<u>P522</u>												
Sp1	–	26.7	66.4	0.1	–	–	1.2	–	0.8	0.1	–	95.3
<u>P583</u>												
Sp1	–	23.1	55.5	6.9	1.5	0.0	0.2	–	11.4	–	0.1	98.9
<u>P440</u>												
Sp1	–	25.1	60.4	5.1	1.9	0.1	0.3	0.1	5.4	–	–	98.3
Sp2	–	27.5	69.3	0.1	0.3	0.1	0.3	–	0.8	–	0.1	98.5
Sp3	–	25.9	62.5	2.7	1.1	0.1	0.5	–	5.0	–	0.1	97.9
<u>P392</u>												
Sp1	–	30.5	57.1	6.2	0.1	0.0	1.4	0.1	1.6	–	–	97.0
Sp2	–	31.0	53.9	7.9	0.2	0.0	1.1	–	2.2	–	0.1	96.5

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11

