
Sea Farming — an Appropriate Technology for Generating Sea Food

by S.Z. Oasim, National Institute of Oceanography, _{Dona Paula}, Goa 401 004, India

h a developing country like India which has a population g about 630 inillion and an annual growth rate of 2.1%, to additional yearly requirement of protein cannot easily a gained from the conventional sources such as agriculwe durying, poultry and cattle farming or traditional Shing. While the production of food grain in India is gaining an all-time record of about 140 million tonnes/ Rul, the production of sea food with an annual yield of phoximately 1.3 million tonnes, has, during the last few gols, shown only a marginal increase. This is also true for world fish production which has remained more or less gationary at about 68-70 million tonnes and the last six or gen years have often been referred to as 'stagnating years f world fisheries'. About 5 million tonnes of food is partition in the world by aquaculture from approximately 20000 hectares of wetland under cultivation, therefore igh shout 7-8% of the total world's fish production comes from aquaculture.

At present the exploitation of natural fish populations from in traditional inshore fishing grounds of India has almost leaked a level from where, unless a gigantic effort is made leaxploit the deep sea areas, no substantial increase appears to possible. Deep sea fishing involves a great deal of tensy in investment and is likely to give a very low rate of them. Therefore, what seems to be the most promising by in which to gain extra food resources is by sea fairning leaguefulture.



Rela Raft made of wooden frame and hambous used for metallization of mussels on hanging ropes in Goa.



field increase in size of mussels on ropes within five months (laid on side).



tele Large bunches of mussels attached to a portion of the rope (laid on side).

Sea farming involves the application of certain principles of animal husbandry to reduce natural predation, improve chances of survival and finally introduce such changes in the environment which can lead to a speeding up of growth from early life to maturity so that the yield is quick and large.

Although the tradition of fish culture in India on a small scale dates back to ancient times, the application of sectarning technology on a commercial scale is a relatively new field. It is now being considered an appropriate technology for producing large quantities of protein-rich food. The reasons for such an approach are, to produce high quality sea food at a reasonable cost; to utilise the vast wetlands which are at present innised and are available at a cheap rate, and finally, to make use of the abundant manpower for the production of the choicest variety of sea foods.

In India, nearly 7.5 million people depend on marine and freshwater fish resources for their livelihood including those who are engaged in industries related to fishing Of the country's available coastal wetlands of about one million hectares, only a small percentage is presently used for fishing while most of it is readily available for farming. The brackish water and nearshore areas offer great potential for farming shrimps, shell-fish, lobsters and exotic lishes.

Studies on the technical feasibility of farming of aquatic organisms have yielded very promising results and some of the production figures are much higher than those reported either from Europe or America. In tropical countries, marketable sizes of captive fish can be obtained in less than half the time needed in colder regions, resulting in annual yields many times greater. Cost-benefit-analysis shows a high rate of return on investment, thus making sea farming an economically viable industry. With new and better technologies a much higher yield is expected.

These technologies include production of fish spawn, for mulation of fish food, development of mechanical food dispensing devices, maintenance of an adequate volume of water, control of disease and genetic upgrading.

One of the difficult problems as yet unsolved in fish farming is the non-availability of quality fish spawn for stocking on a large scale. Attempts have been made to find good sources of spawn from the estuarine and nearshore areas where cultivable species normally breed. The other method used to raise fish spawn is to make the pregnant adults spawn in captivity by injecting hormones. Although both these methods have given encouraging results, the spawn obtained from the two sources are not adequate to meet the total demand. It has been demonstrated that an Indian fish farmer can earn from Rs.7,000 to Rs.10,000 in one season by selling shrimp, lobster and fish spawn of stockable size by raising them in one to two hectares of brackish water.





Fig. 2a Oyster bags hung from nylon ropes in an estuarine farm

As compared to mechanised fishing, aquaculture can be carried out using inexpensive gadgets and simple techniques. The advantages of aquaculture as compared to traditional fishing are:

- 1. dispensing with expensive fishing craft and gear together with the cost of their operation and maintenance
- 2. low capital investment and little recurring expenditure
- 3, prediction of unit yield in advance
- 4. less loss of man-hours due to bad weather or breakdowns
- 5, no risk to consumers as the controlled environment could be made more or less pollution-free
- supplementary income from the supply of fish spawn if fish breeding and hatchery practices are combined with fish farming.

During the last few years very encouraging results have been obtained on composite fish culture of freshwater arps, resulting in a very high annual yield of 8750 kg/ha. Tope culture of mussels from floating rafts (Fig.1a) in stuarine and coastal waters, with a possibility of three rops per year, has shown a phenomenal yield of 480 pmes/ha/year. It has a rate of return of 181% on a low littal investment of Rs.2,500. Similarly, the giant edible yeter, when grown in frames or in bags made of nylon at hung from floating rafts or from ropes in an estuarine (Figs.2a and 2b) has led to a 40% reduction in time in taining commercial size as compared to the period it takes grow in its natural environment.





The success achieved in the production of cultured in India clearly demonstrates that aquaculture is an arriate technology. In recent years, the natural populai the pearl oyster, because of mass killing by humnatural pearls, had declined alarmingly. The development indigenous technique for the culture of pearl oyste floating rafts resulted in the building of a large popular of pearl oysters within a short time (Fig.3). This gave the development of a technology for producing perfectly spherical pearls by nuclei implantation. The nuclei implanted in the tissue of oysters were also development of a technology.

In India, culture of shrimps in low-lying areas and p fields is an age-old practice but the rate of return traditional methods is fairly low. With the application a few simple innovations, such as choosing the right specontrolled stocking and maintenance of an adequate su of natural food, the yield is constantly on the inci (Fig.4). Recently, the polyculture of prawns along mullets and milk fish in brackish water farms has g very promising results. Fish and shrimp production, a cheap technology and without artificial feeding, was ab 2220 kg/ha/year. This included 1960 kg of fish and 26 of shrimps and fetched a price of Rs.15,770. Lobster faing is going to give even better returns once an answer their natural behaviour in captivity—cannibalism—is four

However, in spite of a number of 'plus points' in favour sea farming, the ultimate success of aquaculture in a count



3a Frame-net containing pearl oysters ng lowered in water.



Fig. 3b A cultured pearl show i in the tissue by arrow.




Fig. 3c First few cultured pearls produced in India.

ust 1979

27



fit. 4 Male (larger) and female (smaller) of the giant

the India will largely depend upon the availability of subtantial assistance in the form of inputs including credit subtantial assistance from a technical guidance from since scientists. Besides financial and technical assistance, aboral leusing policy of suitable wetlands is also necessary that fish farmers do not have much at stake at least in semilal phase of the development of aquaculture.

has lish farming on a commercial scale has great potential stoolly in providing work to the rural unemployed but bein inceiling the protein requirement—the lack of which had at the major causes of many diseases in developing series. Though aquaculture cannot replace traditional beins, it would certainly provide an appropriate technology for generating extra sea food. The importance of paralture in the Indian conomy as a small-scale rural clustry should not be overlooked at the present stage of a national development because it is a labour-intensive clusty which would mean an excellent investment of paralters ources.

0305-0920/79/1907-0045/\$01.50/0