

Some Oceanographic Observations in the Polynya and along a section in the Southwest Indian/Antarctic Ocean

S. W. A. Naqvi¹

ABSTRACT

Hydrographical and hydrochemical observations have been made at a shelf station off Princess Astrid Coast, Dronning Maud Land, Antarctica, and at 11 other stations along a transect from Antarctica to about 30°S latitude. (The shelf station, located in a trough, has been occupied at 1-month interval in January and February.) Significant vertical and temporal variations in temperature and salinity are restricted to the upper 200 m; the water is more or less homogeneous below this depth, probably trapped in the trough and renewed by the winter convection. There is some near-surface thermohaline stratification, especially pronounced in January. Substantial warming of surface waters occurs from January to February, resulting in the melting of ice which reduces the surface salinity. A sub-surface oxygen maximum is observed in January associated with a maximum in primary production. Oxygen concentrations at all depths exhibit decreases from January to February in conjunction with increases in nutrients except phosphate, probably resulting from the oxidation of organic matter following high primary production in January.

Vertical sections of temperature, salinity, potential density, dissolved oxygen, apparent oxygen utilization, phosphate, nitrate, silicate, pH and alkalinity through Antarctic, subantarctic and subtropical zones are presented and discussed. (The Antarctic Divergence is observed appreciably to the south of its previously reported positions around 20°E longitude, indicating some variability in its position. No large gradients in the sea surface temperatures are observed in the Polar Front Zone, supporting the view that the transition from Antarctic to subantarctic waters is less sharp in the area around Crozet island.) Most striking changes in all properties are observed in the transition zone between waters of subantarctic and subtropical types. An oxygen minimum occurs within the upper deep water in conjunction with the temperature maximum. The layer is also characterized by high nutrient (phosphate and nitrate) concentrations and low pH. The lower deep water, identified by a salinity maximum, is characterized by low nutrients. The salinity minimum corresponding to the Antarctic Intermediate Water is occasionally associated with an oxygen maximum. An oxygen maximum and a pronounced thermocline corresponding to the Subantarctic Mode Water, however, are not observed at any station, probably due to a weakly-developed subantarctic zone south of the subtropical convergence in the area of study.

The ratio between the changes in nitrate and phosphate is deduced as $\Delta N : \Delta P = 13.6:1$ (by atoms), in conformity with some recent data from the Indian sector of the Southern Ocean, suggesting a lower N:P ratio in the biomass than the corresponding oceanic average value (16:1).

INTRODUCTION

The oceanographic processes that occur in Antarctic waters are essential to : (1) maintenance of aerobic conditions in the intermediate, deep and bottom waters of the world oceans; (2) the removal of heat and addition of fresh water necessary for a steady state character of deep water; (3) the renewal of warm water sphere; and (4) the equalization of water characteristics of the three major oceans (Gordon, 1971). The importance of these processes had been realized by early workers, which lead to a spurt in research activities in the southern ocean in the second quarter of this century. The coverage of the oceanic regime around Antarctica for this purpose, however, has not been uniform, and, perhaps, the Indian sector has been studied least extensively. Some reports on the hydrography and hydro-chemistry of this region have appeared from time to time (Ivanenkov and Gubin, 1960; Jacobs and

¹ National Institute of Oceanography, Dona Paula, Goa

Georgi, 1977; Sen Gupta and Qasim, 1983; Rama Raju and Somayajulu, 1983; Le Jehan and Treguer, 1983). The present study was undertaken to provide additional information on the variations in some hydrographical and chemical parameters based on observations along a transect in the Southwest Indian/Antarctic Ocean as a component of the Third Indian Scientific Expedition to Antarctica.

MATERIALS AND METHODS

11 hydrographic stations were occupied on a transect running from 68°30'S, 19°55'E to 29° 43'S, 54°34'E, during the return voyage of *M.S. Finn polaris* from 3 to 16 March, 1984. Unfortunately due to some mechanical problems with the hydrographic winch, the sampling was restricted to the upper 2,000 m of the water column.

Some oceanographic work was also carried out within a polynya when the ship was alongside or near the ice-shelf. A station was selected in a trough off Lazarev, the abandoned Soviet station (69°59.20'S, 12°42.15'E) where depths exceeding 700 m were encountered on the shelf, otherwise 100–200 m deep (Gupta, 1984 – Fig. 2). The station was occupied twice on 11 January and 7 February, 1984. By late February, just before the vessel sailed on her return voyage, attempts

to occupy the same station again were unsuccessful due to hostile weather.




Fig. 1 Location of sampling stations

Water samples from standard depths at the stations shown in Fig. 1 were collected using PVC Niskin samplers equipped with reversing thermometers. These were analysed onboard the ship shortly after the collection for nutrients (phosphate-phosphorus, nitrate-nitrogen, nitrite-nitrogen and silicate-silicon), dissolved oxygen, alkalinity, pH and salinity. Inorganic micro-nutrients and dissolved oxygen were estimated following standard procedures (Grasshoff, 1976). The pH was measured with the ORION Research Model 701 A/digital IONALYZER, calibrated with NBS buffers. The pH was corrected for *in-situ* temperature according to Gieskes (1969). The pressure effect was incorporated with the data of Culberson and Pytkowicz (1968). Alkalinity was estimated by the pH method of Culberson *et al.* (1970) and salinity was determined with Autosal model 8400 salinometer.

RESULTS AND DISCUSSION

Characteristics of Shelf Waters off Lazarev

Profiles of salinity, potential temperature (θ), potential density (σ_θ), silicate-silicon, dissolved oxygen, apparent oxygen utilization (AOU), phosphate-phosphorus and nitrate-nitrogen at a station along the ice-shelf off Lazarev, occupied in January and repeated in February, are given in Figs. 2a and 2b. Marked variations in all the properties are observed with depth, and there are appreciable differences between the two sets of observations, especially pronounced in the upper 200 m of the water




Fig. 2a Vertical profiles of salinity, potential temperature (θ), potential density (σ_θ) and silicate-silicon at the shelf station occupied on 11th January (circles) and 7th February (triangles)

column. Below this level, the water is almost isothermal, with the two sets of observations yielding remarkably similar temperature data. Variations in salinity are also relatively small below 200 m than those observed in the overlying water column. There is an evidence for shallow thermohaline stratification, more pronounced in January as compared to February, and a thick-well-mixed surface layer does not appear to exist during either month. A pronounced temperature minimum, characteristic of waters south of the Polar Front Zone (PFZ), was not observed at this station. This is not surprising considering that the temperature minimum sinks to depths greater than 200 m near Antarctica, and may even be absent at some locations (Wyrki, 1971). A weakly-developed temperature maximum was observed at 45 m in January (Fig. 2a), possibly due to a decrease in sea surface temperature through contact with pack ice; however, it eroded in February with considerable warming of the surface layer (the mean temperature in the upper 100 m increased from -1.38 in January to -1.19°C in February). The increase in surface temperatures was accompanied by a decrease in salinity from 34.212 to 34.130×10^{-3} , evidently resulting from the melting of ice. Although the entire water column appears to be

fresher in February as compared to January, the difference in the deeper layers (depth > 200 m) are not very large, and could well be due to a slight offset in salinity data. In view of the relatively small variations in thermohaline characteristics of waters below ca. 200 m, it seems reasonable to assume that this layer is renewed by thermal convection only in winter and is topographically isolated during the summer. Consequently, it should reflect the hydrographic conditions prevailing during the winter months. If this is true then the salinity of shelf waters during winter should not exceed 34.4×10^{-3} , and the temperature should range from -1.80 to 1.85°C , corresponding to $\sigma_\theta < 27.80$. Thus, it can be inferred that the ice formation during the winter does not seem to increase the density adequately for this part of the shelf to be a source of bottom water production.




Fig. 2b Vertical profiles of dissolved oxygen, apparent dissolved oxygen utilization, phosphate-phosphorus and nitrate-nitrogen at the shelf station occupied on 11th January (circles) and 7th February (triangles)

The oxygen profiles for January and February are quite similar, but there are two important differences between the two sets of observations (Fig. 2b): (1) a pronounced oxygen maximum observed at 23 m in January was absent in February; and (2) the oxygen concentrations at all levels were consistently lower during the latter observations, the difference ($\sim 0.1 \text{ cm}^3 \text{ dm}^{-3}$) being larger than the estimated analytical error ($< 0.05 \text{ cm}^3 \text{ dm}^{-3}$). The oxygen maximum observed in January was associated with pronounced minima in AOU, nitrate and phosphate (Fig. 2b). These features probably result from the inhibition of photosynthesis at the surface during January which shifts the depth of maximum production a few metres below the surface. Measurements of phytoplankton biomass and the primary production support this view with the values being maximal between 10 and 20 m (Pant, 1985). The phytoplankton biomass was at maximum around mid-January (about the time the first station was worked), after which the values decreased until the end of the month, increased slightly in early February, and decreased again thereafter until the early March (Pant, 1985). The lower oxygen

concentrations observed in February merely reflect the biochemical consumption of oxygen for the oxidation of organic matter following high primary production in the preceding month. Qualitatively consistent with this interpretation are increases in nitrate and silicate at all levels. Surprisingly, the phosphate concentrations apparently decreased during the same period, a discrepancy that is difficult to account for. It is possible that the phosphate values for January are in error, and hence the phosphate data are excluded from the following discussion.

The increases in nutrient concentrations appear to be larger than predicted from the Redfield-Ketchum-Richards (RKR) model (Redfield *et al.* 1963). The increase at six depths between 40 and 400 m ranged between 8.4 and 18.9 (mean 12.9 $\mu\text{g-at. dm}^{-3}$) for AOU; 1.2 and 3.4 (mean 2.18 $\mu\text{M dm}^{-3}$) for nitrate; and 0.6 and 2.4 (mean 1.15 $\mu\text{M dm}^{-3}$) for silicate. If the regeneration ratios $\Delta\text{C} : \Delta\text{N} : \Delta\text{P} : \Delta\text{Si} = 106:16:1:15$ are assumed to hold good (Richards, 1958; Redfield *et al.* 1963). The average biogenic additions of nitrate and silicate between the two sets of observations could be calculated to be 0.70 and 0.75 $\mu\text{M dm}^{-3}$, corresponding to 60 and 34% of the observed increases in silicate and nitrate, respectively. The $\Delta\text{NO}_3 / \Delta\text{SiO}_4$ ratio varied from 0.3 to 1.0, with an average of 0.47, considerably lower than the value reported by Le Jehan and Treguer (1983) from the Indian sector of the southern ocean. The discrepancy could result from different modes of calculation of the regeneration ratios.

Oceanographic section in the southwest Indian/Antarctic Ocean

The distributions of the hydrographical and chemical properties measured along the transect running from 68°30'S, 19°55'E to 29°43'S, 54°34'E are illustrated in Figs. 3–12. The section is neither meridional nor latitudinal; however, latitudinal variations in the properties are likely to be much smaller than the meridional changes, considering that the flow in the southern ocean is mostly zonal (Gordon, 1971).

The most dramatic changes in the properties are observed between Stas. 10 and 11, the transition zone between waters of subantarctic and subtropical types, the so-called subtropical convergence (STC) positioned at about 42°S in this region (Deacon, 1982). South of this discontinuity, the changes in all properties are gradual. Based on observations along a section at 66°30'E, west of Kerguelen, Gambéroni *et al.* (1978) and Gambéroni (1979) suggested that the classical scheme of two convergences, subtropical and Antarctic, did not exist at this longitude. Instead, a single discontinuity was observed between 43°S and 46°S latitudes, leading these authors to conclude that the subantarctic zone did not exist in this region. Deacon (1983), however, doubted such a complete change of the usual scheme arguing that the Antarctic temperature minimum was observed to sink below 200 m, albeit gradually, between latitude 49°S and 48°S along the section worked by Gambéroni *et al.* (1978), and the salinity minimum continued northwards between 3°C and 4°C isotherms into the Antarctic intermediate layer as in other longitudes. He concluded that ".....the area including Marion, Crozet, Kerguelen and Heard islands seems to be one where there is more interchange and less clear gradation between Antarctic and subantarctic waters than in most latitudes" (Deacon, 1983, p.77). The observations of Rama Raju and Somayajulu (1983), along a transect slightly to the west of the present one, appear to be similar to those of Gambéroni *et al.* (1978) and Gambéroni (1979) in that the discontinuities in the sea-surface temperatures at the STC and PFZ are not well-resolved. Although the stations occupied in the present study are not close enough (especially in the critical region between Stas. 7 and 9) to determine the location of the PFZ precisely, it appears that no large gradients in the sea-surface temperatures exist between Stas. 8 and 9 where the PFZ is supposed to lie based on Deacon's (1983) map (assuming that it corresponds to the location where the Antarctic temperature minimum sinks below 200 m, see Deacon, 1934). In the present study, the temperature minimum sank gradually from 97 m at Sta. 7 to 178 m at Sta. 8, and then rapidly to 1087 m at Sta. 9. However, as pointed out by Deacon (1983), in the neighbourhood of shallow soundings and complex bottom topography, such

SOME OCEANOGRAPHIC OBSERVATIONS IN THE POLYNYA... ANTARCTIC OCEAN




Fig. 3 Vertical section of potential temperature ($^{\circ}\text{C}$) between $68^{\circ}30'\text{S}$, $19^{\circ}55'\text{E}$ and $29^{\circ}43'\text{S}$, $54^{\circ}34'\text{E}$




Fig. 4 Vertical section of salinity ($\times 10^{-3}$)




Fig. 5 Vertical section of potential density (σ_{θ})




Fig. 6 Vertical section of dissolved oxygen ($\text{cm}^3 \text{ dm}^{-3}$)




Fig. 7 Vertical section of apparent oxygen utilization ($\mu\text{g-at. dm}^{-3}$)




Fig. 8 Vertical section of phosphate-phosphorus ($\mu\text{M dm}^{-3}$)

as observed around Marion, Crozet, Kerguelen and Heard islands, the sinking of the temperature minimum below 200 m is not as useful an indicator of the surface discontinuity as it is in the deep ocean.

The Antarctic temperature minimum, invariably observed south of Sta. 9, intensified southward. It is shallowest (~ 50 m) at Sta. 4. Occurrence of upwelling at this station is evident from the distribution of all properties (Figs. 3–12). The presence of high salinity and phosphate cells within the upwelling water (Figs. 4 and 8) implies eddies. However, Sta. 4 is located at least $2\frac{1}{2}$ degrees to the south of the previously reported positions of the axis of the Antarctic Divergence in the region. Deacon's (1982) map shows the divergence to occur slightly north of 65°S ; hence, Sta. 5, instead of Sta. 4, should be expected to be located closer to the zone of circumpolar upwelling. Working in the same area, Rama Raju and Somayajulu (1983) found the divergence to occur close to $63^{\circ}30'\text{S}$ latitude. These data suggest considerable variability in the position of Antarctic Divergence around 20°E longitude.

Underlying the Antarctic surface water which comprises of waters within and above the temperature-minimum layer south of the PFZ, is the Warm Deep Water (WDW) flowing southward and upward to compensate for the northward and downward components of flow of the surface and deep waters. The deep water has two components characterized by maxima in temperature and salinity respectively, the temperature maximum occurring considerably shallower (see Gordon, 1971; Jacobs and Georgi, 1977). An oxygen minimum is also observed, generally below the temperature maximum but could be near or above it in some regions (Gordon, 1967). In the present study, the temperature maximum, shoaling up from 1087 m at Sta. 9 to 285 m at Sta. 4, cannot be observed north of Sta. 9. The oxygen minimum invariably occurs at the same depths as the temperature maximum (Figs. 3 and 6). The depth of the oxygen minimum increased northward, and north of the STC, the minimum being at depths exceeding 1400 m. The distributions of phosphate and nitrate (Figs. 8 and 9) follows much the same pattern, with maxima in their concentrations occurring at same levels, located slightly above the temperature maximum and oxygen minimum. The lower deep water, characterized by a salinity maximum, is also characterized by relatively low concentrations of nitrate and phosphate.

A salinity minimum corresponding to the Antarctic Intermediate Water (AIW) is observed at all stations north of the STC at depths ranging between 971 and 1148 m (Fig. 4). Across the STC, the minimum can be traced to the surface layer in the PFZ between 3°C and 4°C isotherms. Within the core of AIW, salinities are lower than 34.5×10^{-3} at all stations in conformity with the results of Warren (1974), Jacobs and Georgi (1977) and Spencer *et al.* (1982) that the 34.5 isohaline associated with AIW extends well north of 37°S, the boundary inferred from the previous data (Deacon, 1937; Wyrki, 1971). A maximum in dissolved oxygen is associated with the AIW salinity minimum at Sta. 11,




Fig. 9 Vertical section of nitrate-nitrogen ($\mu\text{M dm}^{-3}$)

but not at other stations, indicating erosion of the oxygen maximum during the northerly flow (Fig. 6). Surprisingly, the usually more prominent shallower oxygen maximum is not observed at any station in the present study. This maximum, associated with the Sub-Antarctic Mode Water (SAMW) has been observed north of the STC in the Indian Ocean by several workers (Ivanenkov and Gubin, 1960; Wyrki 1971; Jacobs and Georgi, 1977; Warren, 1981; Spencer *et al.*, 1982). It extends as far to the north as the Arabian Sea (Wyrki, 1971; Sen Gupta and Naqvi, 1984; Naqvi and Sen Gupta, 1985). The oxygen maximum occurs in conjunction with a thermostad, a relic of the distant, late-winter mixed layer in latitudes 40–50°S, at progressively lower temperatures from south Atlantic to southeast Pacific; it should be expected at temperatures 12–13°C between longitudes 40–60°E north of the STC in the Indian Ocean (McCartney, 1977). Indeed the thermostad and the associated pycnostad have been observed as far to the north as lat. 18°S in the Indian Ocean at temperatures 9–12°C (Warren, 1981). However, pronounced thermostad or pycnostad are not discernible at temperatures 12–14°C in Figs. 3 and 5, consistent with the absence of an oxygen maximum. It has been suggested that the circulation of SAMW is related to the wind-driven southern hemisphere anti-cyclonic gyres (McCartney, 1977). Hence, the SAMW in the southwest Indian Ocean would be expected to be derived from the eastern region through the anticyclonic gyre. Erosion of oxygen maximum during such flow is possible, though unlikely. Alternatively, if the flow is mostly meridional at these depths, the possible absence of a well-defined subantarctic zone as discussed earlier might explain the absence of the the thermostad and oxygen maximum associated with SAMW.

The distributions of nitrate and phosphate discussed above are in good agreement with the, previously published results (Redfield, 1960; Wyrki, 1971; Jacobs and Georgi, 1977; Spencer *et al.*, 1982; Le Jehan and Treguer, 1983). However, the nitrate and phosphate concentrations observed in the present study are, inexplicably, substantially higher than reported by Sen Gupta and Qasim (1983) from the same general area. Silicate distribution is more or less similar in both studies. Unlike phosphate and nitrate, surface silicate concentrations fall to low levels well south of the STC, and no large gradients in its concentrations exist between Stas. 10 and 11 (Fig. 10). Distribution of pH (Fig. 11) closely follows that of dissolved oxygen with the minimum values ($\text{pH} < 8.00$) associated with the oxygen minimum within the upper deep water characterized by the temperature maximum.




Fig. 10 Vertical section of silicate-silicon ($\mu\text{M dm}^{-3}$)




Fig. 11 Vertical section of pH

The distribution of normalized alkalinity ($\text{Alk}_n = \text{T.A.} \times 35/\text{S}$) is illustrated in Fig. 12. There are few published alkalinity profiles available from the south Indian Ocean with which to compare the present results. The subtropical front could be seen in Fig. 12 as a zone of rapidly changing Alk_n with the isolines being almost vertical down to a depth of about 1 km, separating the zone of decreasing Alk_n , lying north, from that of uniformly high Alk_n , south of the front. A tongue of low alkalinity associated with AIW observed in the alkalinity sections in the atlas of Spencer *et al.* (1982), based on GEOSECS observations, is not seen in Fig. 12, because of the normalization of alkalinity data.

Nitrate-Phosphate Relationship

Owing to the wide meridional coverage in the present study, large horizontal and vertical variations are expected in reserved nutrients. A relationship between oxygen and nutrients, as determined by Sen Gupta and Qasim (1983), suffers from poor correlation due to considerable scatter of data. The correlation between nitrate and phosphate, on the other hand, is quite good with the two-way linear least squares regression leading to the following equation:


$$\text{NO}_3 = 13.62 \text{ PO}_4 - 0.88 \quad (r = 0.95)$$


Fig. 12 Vertical section of normalised alkalinity ($\text{T.A.} \times 35/\text{S}$) meq. dm^{-3}

The $\Delta \text{NO}_3/\Delta \text{PO}_4$ ratio (13.6) is close to the value (14.3) for the Arabian Sea (Naqvi *et al.* 1982), but is slightly lower than the value (16.0) predicted by the RKR model as well as the ratio (19.0) deduced by Sen Gupta and Qasim (1983). Using the nutrient anomalies calculated from the values expected from simple conservative mixing relationships and the nutrient concentrations actually measured, Le Jehan and Treguer (1983) estimated the uptake and regeneration ratio $\Delta \text{N}/\Delta \text{P}$ to range from 12 to 16.8 between latitudes 40 and 60°S along 66°30'E. Most values (mean 14.0) were, however, close to the ratio deduced in the present study. Thus, the $\Delta \text{N}/\Delta \text{P}$ ratio in the water, and, consequently, the N:P ratio in plankton, in the south Indian/Antarctic Ocean appears to be slightly lower than predicted by the RKR model.

ACKNOWLEDGEMENT

The author expresses his sincere thanks to his colleagues Shri Manoharan, Dr. Aditi Pant and Shri M.R. Nayak for their sustained help in both sampling and analysis. He has pleasure in acknowledging the leadership of Dr. Harsh Gupta and the help rendered by all other members of his team. It has been an honour to be selected to participate in the expedition for which the author is grateful to Dr. S.Z. Qasim and Dr. V.V.R. Varadachari. He is also indebted to Dr. R. Sen Gupta for critically reviewing the manuscript, and to the Captain, officers and the crew of *Finnpolaris* for their generous help and co-operation.

REFERENCES

- Culberson, C. and R. M. Pytkowicz 1968. Effect of pressure on carbonic acid, boric acid, and the pH in sea water. *Limnology and Oceanography*, 13, 403-417.
- Culberson, C., R. M. Pytkowicz and J. E. Hawley 1970. Seawater alkalinity determination by the pH method. *Journal of Marine Research*, 28, 15-21.
- Deacon, G. E. R. 1934. Die Nordgrenzen antarktischen und subantarktischen Wrassers in Weltmeer. *Annalen der Hydrographie und Maritimen Meteorologie*, 42, 129-136.
- Deacon, G. E. R. 1937. The hydrology of the Southern Ocean. *Discovery Reports*, 15, 1-124.
- Deacon, G. E. R. 1982. Physical and biological zonation in the Southern Ocean. *Deep-Sea Research*, 29, 1-15.
- Deacon, G. E. R. 1983. Kerguelen, Antarctic and subantarctic. *Deep-Sea Research*, 30, 77-81.
- Gambéróni, L., J. Geronimi, L. Jeannin and J.-F. Murail. 1978. Hydrologie. *Centre National pour l'Exploration des oceans. Resultates des campagnes a la mer*, 16, 35.
- Gambéróni, L. 1979. Quelques éléments sur l'hydrologie des oceans Indien Sud et Antarctique (Secteur Indien) *Memoires Museum National d'Histoire Naturelle, Nouvelle Serie C, Sciences de La Terre*, 43, 21-31.
- Gieskes, J. M. 1969. Effect of temperature on the pH of seawater. *Limnology and Oceanography*, 14, 679-685.
- Gordon, A. L. 1967. Structure of Antarctic waters between 20°W and 170°W. *Antarctic Map Folio Series*. 6, American Geophysical Society, New York, N.Y.
- Gordon, A. L. 1971. Oceanography of Antarctic waters. In: *Antarctic Oceanography*, edited by J. L. Reid, *Antarctic Research Series*, 15, American Geophysical Union, pp. 169-203.
- Grasshoff, K. 1976. *Methods of seawater analysis*, Verlag Chemie, Weinheim, 326 pp.
- Gupta, H. K. 1984. *Third Indian Scientific Expedition to Antarctica 1983-84, Preliminary Scientific Report*, Centre for Earth Science Studies, Trivandrum, 75 pp.
- Ivanenkov, V. N. and F. A. Gubin. 1960. Water masses and hydro-chemistry of the western and southern parts of the Indian Ocean. *Transactions of the Marine Hydrophysical Institute, Academy of Sciences of the U.S.S.R.*, 22, 27-99, American Geophysical Union.

- Jacobs, S. S. and D. T. Georgi. 1977. Observations on the southwest Indian/Antarctic Ocean. In: *A Voyage of Discovery*, edited by M. Angel, Pergamon Press, pp. 43-84.
- Le Jehan, S. and P. Treguer. 1983. Uptake and regeneration $\Delta Si/\Delta N/\Delta P$ ratios in the Indian sector of the Southern Ocean. *Polar Biology*, 2, 127-136.
- McCartney, M. S. 1977. Subantarctic Mode Water. In: *A Voyage of Discovery*, edited by M. Angel, Pergamon Press, pp 103-119.
- Naqvi, S. W. A., R. J. Noronha and C. V. G. Reddy. 1982. Denitrification in the Arabian Sea. *Deep-Sea Research*, 29,459-469.
- Naqvi, S. W. A. and R. Sen Gupta. 1985. 'NO', a useful tool for the estimation of nitrate deficits in the Arabian Sea. *Deep-Sea Research*, 32, In press.
- Pant, A. 1986. Studeis on Antarctic Phytoplankton. Dec. '83 to Mar '84 (in this volume)
- Rama Raju, D. V. and Y. K. Somayajulu. 1983. Some physical characteristics of the Antarctic and western Indian Oceans. *Scientific Report of First Indian Expedition to Antarctica*, Technical Publication No. 1, Department of Ocean Development, New Delhi, pp. 53-61.
- Redfield, A. C. 1960. The distribution of phosphorus in the deep oceans of the world. *Association d'Océanographie physique, Union géodesique et géophysique internationale, Proces-verbaux 7 (G16)*, 189-193.
- Redfield, A. C., B. H. Ketchum and F. A. Richards, 1963. The influence of organisms on the composition of seawater. In : *The Sea, Vol. 2*, edited by M. N. Hill, Interscience, pp. 26-77.
- Richards, F. A. 1958. Dissolved silicate and related properties of some western north Atlantic and Caribbean Waters. *Journal of Marine Research*, 17, 449-465.
- Sen Gupta, R. and S. Z. Qasim. 1983. Chemical oceanographic studies along a section from 32°S to Antarctica in the south western Indian Ocean. *Scientific Report of First Indian Expedition to Antarctica*, Technical Publication No. 1, Department of Ocean Development, New Delhi, pp. 69-77.
- Sen Gupta, R. and S. W. A. Naqvi. 1984. Chemical oceanography of the Indian Ocean, north of the equator. *Deep-Sea Research*, 31, 671-706.
- Spencer, D., W. S. Broecker, H. Craig and R. F. Weiss. 1982. *GEOSECS Indian Ocean Expedition, Vol. 6, Sections and Profiles*, U.S. Government Printing Office, Washington D.C., 140 pp.
- Warren, B. A. 1974. Deep flow in the Madagascar and Mascarene basins. *Deep-Sea Research*, 21, 1-22.
- Warren, B. A. 1981. Transindian hydrographic section at Lat. 18°S. Property distribution and circulation in the South Indian Ocean. *Deep-Sea Research*, 28, 759-788.
- Wyrtki, K. 1971. *Oceanographic Atlas of the International Indian Ocean Expedition*, National Science Foundation, Washington D.C., 531 pp.