

ERRATA

Ramaiah N. Some aspects of microbiological characteristics in the nearshore waters of Bombay, *Indian J Mar. Sci.*, Vol. 23(2), June 1994, pp. 75-81.

In Table 1 and captions of Figs 3 and 4, the character '8' is missing because of *character missing problem* in the computer system. The corrected versions of Table 1 and captions of Figs 3 and 4 are as below:

Table 1—Various microbiological and primary productivity parameters during February, May and September 1986 at different sampling locations

	Stations					
	1	2	3	4	5	6
<i>February 1986</i>						
TVC (no. $\times 10^3$.ml $^{-1}$)	10.6	38.7	13.5	18.1	16.3	11.1
V _{max} (μ g.C.l $^{-1}$.h $^{-1}$)	0.2851	0.2627	0.3204	0.4351	0.5262	0.7351
Respiration (%)	4.62	5.38	2.47	6.88	9.23	13.80
Turnover (T _i) time (h)	134.15	65.72	42.44	217.76	176.59	114.53
K _t + S _n (μ g.C.l $^{-1}$)	37.37	47.48	25.98	27.84	44.61	51.01
Pr. prod. (mgC.m $^{-3}$.d $^{-1}$)	14.98	25.89	43.97	6.01	6.95	7.69
<i>May 1986</i>						
TVC (no. $\times 10^3$.ml $^{-1}$)	46.8	62.5	57.6	84.0	56.0	10.8
TDC (no. $\times 10^6$.ml $^{-1}$)	4.601	—	7.089	—	0.162	—
V _{max} (μ g.C.l $^{-1}$.h $^{-1}$)	0.2515	0.1972	0.2899	0.3901	0.4147	0.7432
Respiration (%)	4.01	5.46	2.02	6.57	8.46	13.52
Turnover (T _i) time (h)	159.19	105.64	70.89	172.78	151.94	166.27
K _t + S _n (μ g.C.l $^{-1}$)	40.04	5.63	20.55	33.03	21.98	13.64
Pr. prod. (mgC.m $^{-3}$.d $^{-1}$)	6.04	9.30	29.89	6.37	5.55	7.38
<i>September 1986</i>						
TVC (no. $\times 10^3$.ml $^{-1}$)	18.0	12.7	23.3	17.6	22.4	88.0
V _{max} (μ g.C.l $^{-1}$.h $^{-1}$)	0.4025	0.4296	0.3720	0.4341	0.3927	0.8511
Respiration (%)	1.76	7.96	4.41	8.05	9.51	14.67
Turnover (T _i) time (h)	79.50	57.89	39.05	94.40	215.21	180.03
K _t + S _n (μ g.C.l $^{-1}$)	32.00	51.01	41.41	54.21	47.43	58.61
Pr. Prod. (mgC.m $^{-3}$.d $^{-1}$)	34.73	6.49	54.02	7.20	5.91	3.58

Fig. 3—Relationship between Pr. prod. and V_{max} during (A) May, (B) September and (C) February 1986 (n=6, observed r is significant at 1% when it is ≥ 0.875 ; at 5% when it is ≥ 0.755 and at 10% when it is ≥ 0.699)

Fig. 4—Relationship between V_{max} and other kinetic parameters during (A) May, (B) September and (C) February 1986 (n=6, observed r is significant at 1% when it is ≥ 0.875 ; at 5% when it is ≥ 0.755 and at 10% when it is ≥ 0.699)