

## **Massive nitrogen loss over the western Indian continental shelf during seasonal anoxia: Evidence from isotope pairing technique**

**Amit Sarkar<sup>\*1,2</sup>, Syed Wajih Ahmed Naqvi<sup>1,3</sup>, Gaute Lavik<sup>4</sup>, Anil Pratihary<sup>1,4</sup>, Hema Naik<sup>1</sup>,  
Gayatri Shirodkar<sup>1</sup> and Marcel Kuypers<sup>4</sup>**

<sup>1</sup>CSIR-National Institute of Oceanography, Dona Paula, Goa 403 004, India

<sup>2</sup>Present address: Kuwait Institute for Scientific Research, P.O. Box 1638, Salmiya 22017, Kuwait

<sup>3</sup>Present address: Council of Scientific & Industrial Research, Rafi Marg, New Delhi 110 001, India

<sup>4</sup>Max Planck Institute for Marine Microbiology, Bremen, Germany

**\*Corresponding Author:** Phone: +91-9764246977/+965-55470577

Email address: [amitsarkar81@gmail.com](mailto:amitsarkar81@gmail.com)

### **Abstract**

The western Indian continental shelf houses the world's largest naturally-formed coastal low-oxygen zone that develops seasonally during the summer monsoon. We investigated multiple reductive nitrogen transformation pathways and quantified their rates in this system through anaerobic incubations with additions of <sup>15</sup>N-labeled substrates during the anoxic period for three consecutive years (2008 to 2010). Addition of <sup>15</sup>N labeled ammonium (<sup>15</sup>NH<sub>4</sub><sup>+</sup>) resulted in low to moderate anaerobic ammonia oxidation (Anammox) rates in about half of our incubations from the oxygen depleted waters. In contrast, incubations with labeled nitrite (<sup>15</sup>NO<sub>2</sub><sup>-</sup>) led to large production of <sup>30</sup>N<sub>2</sub> over <sup>29</sup>N<sub>2</sub> in all incubation experiments, indicating denitrification to be the dominant N-loss pathway. Rates of dissimilatory nitrate/nitrite reduction to ammonia (DNRA) were found to be highly variable and were lower by an order of magnitude than the denitrification rates. Extrapolation of average rates over the sampling periods and volume of anoxic waters showed large nitrogen removal (3.70 to 11.1 Tg yr<sup>-1</sup>) which is about three times as high as the previously reported estimate (1.3 to 3.8 Tg yr<sup>-1</sup>). Despite the small area it occupies, this shallow seasonal anoxic zone may account for as much as 20-60% the of the total annual fixed nitrogen loss in the perennial oxygen minimum zone of the Arabian Sea.

**Keywords:** Seasonal hypoxia/anoxia, denitrification, anammox, DNRA, nitrogen loss

## Introduction

As in other parts of the North Indian Ocean, hydrography and biogeochemistry over the western Indian continental shelf (WICS) are also dominated by seasonal reversals of surface currents associated with the monsoons (Schott and McCreary, 2001; Naqvi et al., 2006a). During the winter or the Northeast Monsoon (NEM) the West India Coastal Current (WICC) anomalously flows poleward, causing downwelling and oligotrophic, well-oxygenated conditions off the Indian west coast. By contrast, the region behaves as a mini eastern boundary upwelling system during the summer or the Southwest Monsoon (SWM) when the WICC flows toward the equator. The equatorward flow induces upwelling, which is weak except off the southwest coast of India and the west coast of Sri Lanka where winds are more favorable. As a result, there is only modest enhancement of primary productivity and consequent oxygen demand in near-bottom waters (Naqvi et al., 2010a). However, despite moderate oxygen demand there are several factors that contribute to development of intense anoxia in near bottom waters: (1) The upwelled water, derived from the upper part of the mesopelagic oxygen minimum zone of the Arabian Sea is oxygen-depleted to begin with, although not anoxic; (2) Due to slow upwelling the upwelled water remains over the shelf for a long time; and (3) The cold, saline upwelled water is capped by a thin (<10 m) warm, fresher lens, unusual for any upwelling system, formed due to intense precipitation in the coastal zone during the SWM, resulting in very strong stratification very close to the surface, thereby greatly restricting oxygen diffusion from the surface. The exact time of formation of this coastal OMZ varies strongly. It seems to gradually evolve from the south to the north. Upwelling along the WICS has been reported to start in the south (8°N, off Cochin) probably during spring (in March) (Stramma et al., 1996; Schott and McCreary 2001), gradually extending northward. In the north (19°N, off Mumbai) the effect of upwelling is felt at least until early December (Naik, 2003; Shankar et al., 2005). This time span far exceeds the duration of the SWM. However, strong water column stratification starts in June due to intense rainfall in coastal areas, which facilitates the formation of the low oxygen zone. The oxygen deficiency intensifies with time resulting in the prevalence of reducing conditions - conversion the oxidized nitrogen to molecular nitrogen (N<sub>2</sub>) followed by reduction of sulphate (SO<sub>4</sub><sup>2-</sup>) to hydrogen sulphide (H<sub>2</sub>S) - by July-August. In the north (e.g. off Mumbai) such conditions develop later in the season and last up to October-November. Thus, overall, reducing conditions over the WICS prevail for at least 3 months (90 days) (Naqvi et al., 2006b, Naik et al., 2003). These processes operate over a large geographical area, making the WICS the single largest low-oxygen system of its kind in the world, occupying an area of ~200,000 km<sup>2</sup> (O<sub>2</sub><20μM) - an order of magnitude bigger than the area of the famous dead zone of the Gulf of Mexico (area with

$O_2 < 62.5 \mu M$ : 22,000 km<sup>2</sup> - Rabalais et al., 2010). The shallow anoxic zone in the eastern Arabian Sea is believed to provide significant feedback to the global climate change through unusually high emissions of N<sub>2</sub>O (nitrous oxide) and DMS (dimethyl sulphide) (Naqvi et al., 2000; Shenoy et al., 2012).

As evident from the chemical data - a decrease in nitrate (NO<sub>3</sub><sup>-</sup>) and a concomitant accumulation of nitrite (NO<sub>2</sub><sup>-</sup>) - reductive nitrogen removal from this system typically begins to occur during the later phase of the SWM in July/August and continues till October /early November (Naqvi et al., 2009). The reducing zones can be categorized into "suboxic" (functionally identified by the presence of secondary NO<sub>2</sub><sup>-</sup>, although it must be pointed out that recent work has shown that the secondary NO<sub>2</sub><sup>-</sup>-bearing waters are also functionally anoxic (Thamdrup et al., 2012), and anoxic (sulfidic); these zones are geographically separated, often located over the mid- and inner parts of the shelf (Naqvi et al., 2009). Direct measurements of N-loss rates and the processes involved are very scarce from WICS. Heterotrophic denitrification (NO<sub>3</sub><sup>-</sup> → NO<sub>2</sub><sup>-</sup> → NO → N<sub>2</sub>O → N<sub>2</sub>), in which bacteria use NO<sub>3</sub><sup>-</sup> as an electron acceptor for organic matter oxidation (Froelich et al., 1979), was long believed to be the only process removing fixed nitrogen from the ocean, until anammox was recognized to be another major N-loss pathway (Dalsgaard et al., 2003, Kuypers et al., 2003). In previous studies undertaken over the WICS, rates reaching up to 837 nmol L<sup>-1</sup> d<sup>-1</sup> were determined based on changes in NO<sub>3</sub><sup>-</sup> + NO<sub>2</sub><sup>-</sup> concentration over time at fixed sampling locations (Naik, 2003). Similar rate was also calculated assuming that the initial NO<sub>3</sub><sup>-</sup> content of 20 μM in the upwelled water was fully utilized over a month (Naqvi et al., 2006b). Overall annual N-loss in the range of 1.2-3.6 Tg was arrived at by scaling up these rates to the entire shelf for the duration of the seasonal anoxia. However, these estimates should be considered as minimum estimates as replenishment of the nitrate pool had not been taken into account. Devol et al. (2006) estimated N-loss at a few coastal stations by incubating samples spiked with <sup>15</sup>NO<sub>3</sub><sup>-</sup> on board ship and measuring the growth of <sup>15</sup>N-labelled N<sub>2</sub>. The average rate was found to be 33.2 nmol N L<sup>-1</sup> d<sup>-1</sup>. The highest rate measured by these authors was 207 nmol N L<sup>-1</sup> d<sup>-1</sup>, only about a quarter of the rate inferred from the overall changes in the nutrient concentrations. However, these measurements suffered from several uncertainties. For example, they did not take into account the possible N-loss through anammox. Moreover, the signal of labeled N<sub>2</sub> produced from <sup>15</sup>NO<sub>3</sub><sup>-</sup> may be significantly diluted due to mixing of <sup>15</sup>NO<sub>2</sub><sup>-</sup>, originated during the first step of denitrification pathway, with high ambient substrate pool of <sup>14</sup>NO<sub>2</sub><sup>-</sup>. This may lead to underestimation (Holtapells et al., 2011). Also, none of the studies undertaken so far have considered the dissimilatory reduction of nitrate/nitrite to ammonium (DNRA) that could have major implications for the estimation of the N-loss by labeling experiments (Song et al., 2016, Jensen et al.,

2011) as well as for the estimates based on temporal changes in nitrate concentration (i.e. while DNRA removes  $\text{NO}_3^-$  or  $\text{NO}_2^-$ , it is not an N-loss process). Moreover, dissolved inorganic nitrogen (DIN) loading in this region is believed to have significantly increased in recent years due to enhanced synthetic fertilizer consumption in South Asia ( $\sim 17 \times 10^{12} \text{ g N a}^{-1}$ ), exhibiting an increase by factor of 50 since 1960s (Naqvi et al., 2009). Seitzinger et al. (2005) estimated that the South Asian rivers could account for as much as  $4.2 \text{ Tg yr}^{-1}$  of the global riverine DIN input of  $20.8 \text{ Tg N yr}^{-1}$  to the ocean in 1990. However, the concentrations of  $\text{NO}_3^-$  reported from some of the rivers draining into the Arabian Sea are quite moderate (e.g.  $8\text{-}9 \mu\text{M}$  from the Zuari estuary of Goa - Bardhan et al., 2015). In addition, Naqvi et al. (2010b) estimated annual deposition of  $\sim 0.16 \text{ Tg}$  of nitrogen from the atmosphere over the WICS which most probably is also increasing due to fossil fuel burning and industrialization (Duce et al., 2008). Consequently, the N-loss estimated from upwelled  $\text{NO}_3^-$  may be considerably lower than actual.

The main aim of the present study is to identify and quantify the multiple potential reductive nitrogen processes operating in the water column over the WICS during the period of anoxia by employing various combinations of  $^{15}\text{N}$ -labelled substrates. This is the first report evaluating relative contributions of denitrification and anammox to N-loss and also the significance of DNRA from this region. As the experiments were conducted for three consecutive years the results also provide an idea of the inter-annual variability.

## **2. Methods**

### **2.1 Sampling locations and physicochemical profiling of water column**

Samples were collected from several stations located along coast-perpendicular transects during three cruises carried out from 2008 to 2010, on the Coastal Research Vessel *Sagar Sukti* (SaSu#183 and SaSu#205) and using a fishing trawler (CaTS#78). The period of observations was restricted to the time - from late August to late September - when severe oxygen deficiency develops over the WICS. In 2008, only the transect off Goa ( $15.52^\circ\text{N}/73.76^\circ\text{E}$  to  $15.423^\circ\text{N}/73.283^\circ\text{E}$ ) was sampled, whereas in 2009 and 2010 observations along two additional southern transects, off Karwar ( $14.478^\circ\text{N}/74.25^\circ\text{E}$  to  $14.325^\circ\text{N}/73.768^\circ\text{E}$ ) and Mangalore ( $13.14^\circ\text{N}/74.64^\circ\text{E}$  to  $12.87^\circ\text{N}/74.108^\circ\text{E}$ ), were also made (Fig. 1, Supplementary Table T1, T2). Physicochemical characteristics of the water column (temperature, salinity, dissolved oxygen, inorganic nutrients, chlorophyll *a* and turbidity) were determined at each sampling site. N-loss rates were measured over the inner shelf (depth  $<30 \text{ m}$ ) and mid shelf (depths  $30\text{-}50 \text{ m}$ ) where strong reducing conditions prevailed.

A Sea-Bird Electronics CTD system (SBE 911) was used for obtaining continuous profiles of temperature and salinity. Niskin samplers (5 L) fitted with reversing thermometers and mounted on synthetic ropes were used for sampling of seawater from selected depths based on the CTD/O<sub>2</sub> profiles. Dissolved O<sub>2</sub> was measured in each sample following the Winkler titrimetric method (Grasshoff et al., 1983, precision < 1.34 μM). Sub-samples for nutrients (unfiltered) were frozen on *Sagar Sukti* cruises and stored in an ice box on the CaTS field trip for analyses in the shore laboratory using a SKALAR autoanalyser (precisions ±0.06, ±0.01, ±0.01, ±0.01 μM for NO<sub>3</sub><sup>-</sup>, NO<sub>2</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup> and PO<sub>4</sub><sup>3-</sup>, respectively). In most cases, NO<sub>2</sub><sup>-</sup> concentrations were also measured onboard immediately after sampling following the procedure of Bendschneider and Robinson (1952). The lower detection limit of the NO<sub>2</sub><sup>-</sup> measurement was 0.05 μM. Deficiency in DIN was estimated from  $N^*(\mu\text{M}) = [(\text{NO}_3^- + \text{NO}_2^- + \text{NH}_4^+) - 16 \times \text{PO}_4^{3-} + 2.9]$  following Gruber and Sarmiento (1997). Chlorophyll-*a* samples were collected in 1-litre amber colored HDPE bottles and were filtered immediately through glass-fibre filters (GF/F, 47mm diameter, 0.7 μm pore size, Whatman) under low vacuum. The filter papers were soaked in 90% acetone and stored at -20°C in the dark to extract the pigments. Quantitative estimation of chlorophyll-*a* was done after 24 hour spectrofluorometrically at 655 nm wavelength (Strickland and Parsons, 1972). Precision of chlorophyll *a* measurement was ±0.001 mg m<sup>-3</sup>.

## 2.2 Isotope pairing experiments

Depending on the depth and thickness of the perceived reducing layer, incubation experiments were restricted to 1-3 depths. In 2008, incubations were conducted at two stations at two depths each off Goa (Fig. 2, O<sub>2</sub> panel marked in diamond symbol). In 2009, three stations each along Goa and Karwar and 1 station off Mangalore were sampled. In 2010, three stations were sampled off Goa, two off Karwar and one off Mangalore (Fig. 2, Table 1, Supplementary Figures S2 and S3). Water samples for incubation experiments were taken immediately after the retrieval of samplers in 250-ml glass serum bottles. Water approximating three times of the volume of the bottle was allowed to overflow before the sample was sealed using butyl rubber septum, taking care not to trap any air bubble, and crimped with aluminum cap. Samples were purged with helium for 20 minutes to reduce the background N<sub>2</sub> concentration as well as to remove oxygen that might have been inadvertently introduced during sampling. After purging with helium for 15 minutes, various substrates (<sup>15</sup>NH<sub>4</sub><sup>+</sup>, <sup>15</sup>NO<sub>2</sub><sup>-</sup> and <sup>15</sup>NO<sub>2</sub><sup>-</sup> + <sup>14</sup>NH<sub>4</sub><sup>+</sup>, Sigma-Aldrich stable isotope labeled products) were added to serum bottles and purging was continued for another 5 minutes to ensure removal of oxygen added along with the substrates and also proper mixing. The final concentration of each substrate was 8 μM. Samples were then quickly transferred into 12-ml Labco Exetainers carefully avoiding any

atmospheric contamination and incubated in dark at near *in situ* temperature for 36-48 hrs. Microbial activities were terminated at approximately equal time intervals by adding 100  $\mu$ l saturated HgCl<sub>2</sub> solution after introducing 2 ml of helium headspace (Holtappels et al., 2011). Exetainers were stored upside down until analysis to prevent gas loss from the headspace through the septa.

### 2.3 Analysis and calculation of denitrification, anammox and DNRA rates

The stable isotopic composition of N<sub>2</sub>, measured as <sup>28</sup>N<sub>2</sub>, <sup>29</sup>N<sub>2</sub> and <sup>30</sup>N<sub>2</sub> in the headspace, was determined within a few weeks of collection using a continuous flow mass spectrometer coupled with a custom build gas chromatograph (Fisons VG Optima, Manchester, UK). 100  $\mu$ l of headspace sample was slowly and carefully drawn from the vial and injected into the GC-IRMS system using gas-tight syringe. Deionized water was used to compensate for the slight under-pressure in the headspace arising from outward He diffusion during storage as well as to make up for the volume of the gas withdrawn. Air was injected as the standard before and after each batch of 8 samples. Concentrations of <sup>29</sup>N<sub>2</sub> and <sup>30</sup>N<sub>2</sub> produced during incubation were calculated as described by Holtappels et al. (2011). After analysis of the labeled N<sub>2</sub>, <sup>15</sup>NO<sub>2</sub><sup>-</sup> amended samples were used for DNRA rate measurements. The isotopic composition of NH<sub>4</sub><sup>+</sup> in the samples was determined after chemically converting it to N<sub>2</sub> with sodium hypobromite (NaOBr) solution using the same mass spectrometer (Warembourg, 1993; Preisler et al., 2007).

The rates were computed as the slope of linear regression of <sup>15</sup>N-production over time. Concentrations of N<sub>2</sub> produced via denitrification and anammox pathways were calculated following Thamdrup and Dalsgaard (2002). N<sub>2</sub> production via denitrification was computed from <sup>15</sup>NO<sub>2</sub><sup>-</sup> incubation as:

$$\text{Denitrification} = {}^{30}\text{N}_2 \times [\text{F}_{\text{NO}_2}]^{-2}$$

Where, F<sub>NO<sub>2</sub></sub> refers to the <sup>15</sup>N-mole fraction in the NO<sub>2</sub><sup>-</sup> pool.

Anammox was quantified from the <sup>15</sup>NH<sub>4</sub><sup>+</sup> incubation considering the mole fractions of <sup>15</sup>N in the substrate pool (F<sub>NH<sub>4</sub></sub>) and excess <sup>29</sup>N<sub>2</sub>:

$$\text{Anammox} = {}^{29}\text{N}_2 \times [\text{F}_{\text{NH}_4}]^{-1}$$

In some cases (at stas. G6 and G7 off Goa, and sta. K2 off Karwar in 2009) where NH<sub>4</sub><sup>+</sup> concentrations were not available, anammox rates were calculated from <sup>15</sup>NO<sub>2</sub><sup>-</sup> incubation as:

$$\text{Anammox} = {}^{29}\text{N}_2 \times [\text{F}_{\text{NO}_2}]^{-2} - [\text{Denitrification} \times 2 \times (1 - \text{F}_{\text{NO}_2})]$$

DNRA rates were calculated from the slope of linear regression of <sup>15</sup>N-labeled NH<sub>4</sub><sup>+</sup> production with time in the NO<sub>2</sub><sup>-</sup> spiked samples (Jensen et al., 2011).

### 3 Results

#### 3.1 Physicochemical characteristics

The physicochemical conditions along various transects visited in the three years were very similar in that water column was strongly stratified in all cases (as exemplified by transect off Goa in Fig.2). In 2008 and 2009, surface water had potential density ( $\sigma_\theta$ ) of  $\sim 20$ - $22.5 \text{ kg m}^{-3}$  that increased to  $23.5$ - $24.5 \text{ kg m}^{-3}$  in the cooler and more saline subsurface upwelled water below the thermocline (Fig.2, Plates A and B). Due to a more intense rainfall in 2010, surface  $\sigma_\theta$  values were much lower (Fig.2, Plate C) than the  $\sigma_\theta$  values observed during 2008 and 2009. In all cases, surface waters were mostly oxygen saturated or slightly supersaturated ( $>200 \mu\text{M}$ ) but  $\text{O}_2$  concentrations declined sharply ( $<10 \mu\text{M}$ , often reaching non-detectable limit) below the thin surface layer. Presence of sulphide was noticed by its strong odour along all transects in 2009. Surface water was mostly nutrient depleted. Over the deeper outer shelf (depth  $>50\text{m}$ )  $\text{NO}_3^-$  levels were higher ( $>15 \mu\text{M}$ ) below the surface, but the concentrations decreased steadily towards the coast. Lower  $\text{NO}_3^-$  levels in oxygen-depleted water over the inner- and mid-shelf regions often coincided with elevated  $\text{NO}_2^-$  (up to  $6 \mu\text{M}$ ) indicating vigorous anaerobic microbial  $\text{NO}_3^-$  reduction (Fig. 2). Dissolved  $\text{NH}_4^+$ , whenever measured, showed some accumulation in  $\text{NO}_3^-$  reducing water, either being produced through organic matter mineralization and/or DNRA or originating from the underlying sediment. Except for the year 2010, off Goa, strongly negative  $\text{N}^*$  values in subsurface waters over the shelf indicated high fixed N-loss under suboxic/anoxic conditions (Fig. 2, Plates A and C). The two southern transects (off Karwar and Mangalore) also showed similar physicochemical characteristics (Supplementary Figs. S2 and S3)

#### 3.2 Nitrogen reduction pathways over the WICS

Denitrification was detected in all samples, as indicated by significant ( $r^2 \geq 0.7$ ) production of  $^{30}\text{N}_2$  in  $^{15}\text{NO}_2^-$ -spiked incubations. In fact, most of the N-loss over the shelf appears to be through this process (Table 1, Supplementary Figure S4, 1 to 33). Production of  $^{30}\text{N}_2$  exceeded that of  $^{29}\text{N}_2$  in a majority of experiments, maintaining the theoretical production ratio of 2 ( $1 - F_{\text{NO}_2}$ )/ $F_{\text{NO}_2}$  through random isotope pairing as proposed by Nielsen (1992). In 2008, denitrification rates along the Goa transect ranged from  $0.91 \pm 0.08$  to  $1.93 \pm 0.24 \mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ . Higher denitrification rates were observed in 2009 at the inner shelf stations. At Sta. G5, the denitrification rate averaged  $6.49 \pm 0.65 \mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ ; even higher rates were measured along the two southern transects,  $9.34 \pm 1.32 \mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$  off Karwar and  $10.03 \pm 1.7 \mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$  off Mangalore (Table 1). In 2010, denitrification rates were considerably lower over the inner shelf as compared to those in 2009. However, at the mid

shelf stations inter-annual variations were less pronounced. Using all the three year's data, the average denitrification rate was calculated as  $2.45 \pm 0.63 \mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ .

Anammox activity was confirmed by results of  $^{15}\text{NH}_4^+$  incubations that led to significant  $^{29}\text{N}_2$  production in over half (60%) of the all experiments (Table 1, Supplementary Fig. S5, 1 to 20). Anammox rates were calculated using  $^{29}\text{N}_2$  production rates from both  $^{15}\text{NH}_4^+$  and  $^{15}\text{NO}_2^-$  incubations; these are presented in Table 1. The rates were significantly different from each other (Independent samples T test,  $p < 0.05$ ) with consistently higher rates from  $^{15}\text{NO}_2^-$  incubations. This is expected due to the differences in  $^{15}\text{N}$  enrichment in the substrate pool (Bulow et al., 2010 and references there in). For consistency, in the present study we considered anammox rates derived only from  $^{15}\text{NH}_4^+$  incubation which is recommended to be more direct and precise. In 2008, anammox rates ranged between 0 and  $0.02 \pm 0.004 \mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ . This process was mostly undetectable in 2009. In 2010, overall anammox rates were higher but highly variable from below detection to  $0.43 \mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ . The average anammox rate derived from all our  $^{15}\text{NH}_4^+$  incubations was  $\sim 0.06 \pm 0.004 \mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ . These numbers are two orders of magnitude lower than the above-mentioned denitrification rate.

DNRA activities were seen only sporadically. In both 2009 and 2010, DNRA activity was found at one depth along the Goa transect, and anammox was not detected in 2008. Off Karwar DNRA was found at 3 depths in 2009, but it was absent in 2010. A completely opposite trend was observed off Mangalore. The average DNRA rate ( $109.25 \pm 22.1 \text{ nmol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ ) was higher than that observed over the Omani shelf (highest  $\sim 40 \text{ nmol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ ).

### **3.3 Spatial & inter annual variability of nitrogen reduction processes**

The spatio-temporal trends in denitrification and anammox rates were tested using non-parametric Mann Whitney t-test (two-tailed). Due to large intra-annual variability, no inter-annual variability was evident in both processes (Supplementary Figure S10 a and b). However, denitrification rates along the Goa coast were significantly lower than that at the Karwar coast at  $P \leq 0.1$  ( $P$ : 0.10), and Mangalore coast at  $P < 0.05$  ( $P$ : 0.04). Anammox rates were different only between Karwar and Mangalore at a significance level of  $P \leq 0.1$  ( $P$ : 0.07). Denitrification was found to be more active at the oxic-suboxic/anoxic interface compared to deeper water (Figs. 3a and 3b). In 2009, low rates of denitrification were measured at the base of the oxycline at stations G6 (Fig. 3b) and K3 where  $\text{O}_2$  concentrations were  $> 30 \mu\text{M}$  (Table 1). These  $\text{O}_2$  levels probably inhibited denitrification at these stations. In fact, with just a few exceptions, low levels of denitrification (only 10-25% of the mean of all measured rates) were consistently found in waters containing measurable  $\text{O}_2$ . Anammox activity



was low and stochastic to infer any trend in its depth distribution (Table 1, Fig. 3a and 3b). In 2010, anammox rates in bottom waters at stations G5 and G7 off Goa, were not only higher than those at the oxycline but also the highest recorded in this study (Table 1, Figs. 3a and 3b). However, at most of the depths where the anammox activity was detected O<sub>2</sub> levels were low but often measurable (<5 μM).

Overall denitrification rates did not show large inter annual variations over the 3 years period of the study. Over the inner shelf, average denitrification rate was higher in 2009 (6.49±0.65 μmol N<sub>2</sub> L<sup>-1</sup> d<sup>-1</sup>) compared to 2010 (1.98±1.1 μmol N<sub>2</sub> L<sup>-1</sup> d<sup>-1</sup>, respectively). In comparison, average anammox average rate was the highest in 2010 (0.07±0.004 μmol N<sub>2</sub> L<sup>-1</sup> d<sup>-1</sup>), 7 times more than the rates measured in 2008 (0.01±0.002 μmol N<sub>2</sub> L<sup>-1</sup> d<sup>-1</sup>). In 2009, near-bottom waters at most inner shelf stations were found to be sulfidic (Supplementary Fig. S1). At these stations denitrification rates were exceedingly high while anammox activity was not detected (Fig. 3a).

### 3.4 N-loss rates over the WICS

Despite the intra-annual variability observed at individual stations/sections, our data show quite similar average denitrification rates for three years. The mean values of our three-year N-loss rates through denitrification and anammox were 2.45±0.6 and 0.06±0.004 μmol N<sub>2</sub> L<sup>-1</sup> d<sup>-1</sup>, respectively. For up-scaling these values over the entire shelf we followed the assumption previously made by Naqvi et al. (2000) and Naik (2003). Naqvi et al. (2000) estimated the area of the hypoxic zone over the WCSI to be 180,000 km<sup>2</sup>. However, O<sub>2</sub> levels over this entire area are not low enough to support reductive N-loss, and as a lower limit about one third of the entire shelf, 60,000 km<sup>2</sup>, was assumed to functionally anoxic. The total volume of reducing water was estimated to range between 1.2 and 3.6 x 10<sup>12</sup> m<sup>3</sup> taking the thickness of the reducing layer to be 20m. Applying our 3 years average N-loss rates to this volume and assuming the reducing condition to last for 90 days, the overall N-loss would range from 3.70±0.091 to 11.1±2.72 Tg yr<sup>-1</sup> via denitrification, and from 0.09±0.01 to 0.27±0.03 Tg yr<sup>-1</sup> via anammox (Table 2).

## 4 Discussion

### 4.1 Oxygen deficiency and associated biogeochemical properties

Consistent with earlier observations (Naqvi et al., 2009; 2010a), suboxic (Winkler O<sub>2</sub>< 10 μM; NO<sub>3</sub><sup>-</sup> & NO<sub>2</sub><sup>-</sup>> 0 μM) and anoxic (O<sub>2</sub> = 0 μM; H<sub>2</sub>S > 0 μM) conditions were found to prevail over the inner-shelf and mid shelf regions in the present study as well. This oxygen deficient environment that evidently supported reductive nitrogen transformations (Hannig et al., 2007; Naqvi et al., 2010a, Lavik et al., 2009) was chosen for <sup>15</sup>N labeling experiments (Fig.2, depths marked by diamond in O<sub>2</sub>

panels). Oxygenated surface water had extremely low  $\text{NO}_3^-$  concentrations ( $<1 \mu\text{M}$  at most of the stations) with very few exceptions. Average surface chlorophyll *a* concentration was  $\sim 2 \text{ mg m}^{-3}$ , with the highest concentration ( $7.01 \text{ mg m}^{-3}$ ) occurring at a station M8. Earlier observations made during the Southwest Monsoon also revealed high surface chlorophyll levels (e.g.  $2.63 \text{ mg m}^{-3}$  at station G5 - Maya et al., 2011). Bhavya et al. (2017) reported average carbon uptake rate of  $0.23 \text{ mg C L}^{-1} \text{ d}^{-1}$  water over the coast of Mangalore; this was significantly higher than production during the pre-monsoon but slightly lower than that during the post-monsoon period. In the coastal Arabian Sea primary production has been reported to be the largest contributor to the carbon stock (Barber et al., 2001). Therefore, the low  $\text{NO}_3^-$  levels in surface waters observed in the present study are probably due to biological uptake which in turn makes organic matter available to support microbial respiration in sub surface layers, increasing the  $\text{O}_2$  demand (Devol et al., 2006). Hypoxic water below the surface over the outer shelf was  $\text{NO}_3^-$  rich ( $\sim 15 \mu\text{M}$ ), but  $\text{NO}_3^-$  was gradually depleted as the water upwelled and moved up the shelf. Accumulation of  $\text{NO}_2^-$  in the suboxic water over the inner- and mid-shelf points to active denitrification that apparently varied in strength spatially. When the  $\text{NO}_3^-$  was fully utilized, sulphate reduction commenced leading to buildup of  $\text{H}_2\text{S}$  and  $\text{NH}_4^+$ . The highest  $\text{NH}_4^+$  concentrations (up to  $9.1 \mu\text{M}$ ) in low-oxygen waters (Table 1) are consistent with earlier measurements by Naqvi et al. (2009). Concentration of  $\text{H}_2\text{S}$  was not measured in the present study; however, its frequent presence over the inner shelf was indicated by strong odour. Soon after our observations off Goa in 2009, station located on Goa transect was visited for time series sampling and a build-up of sulphide in subsurface water was noticed (Supplementary Figure S1, reaching concentration as high as  $\sim 12 \mu\text{M}$  (Shenoy et al., 2012). Overall, our results show the presence of intensely  $\text{O}_2$ -depleted waters underlying a shallow and sharp pycnocline suitable for suboxic/anoxic microbial processes like denitrification, anammox and DNRA.

#### **4.2 Regulation of nitrogen reduction pathways over the WICS**

In the present study, denitrification was found to be the dominant fixed nitrogen removal process. The average denitrification rate ( $2.45 \pm 0.6 \mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ ) was more than double of the previously reported value derived from the observed  $\text{NO}_3^- + \text{NO}_2^-$  loss over time (Naik, 2003) and almost two orders of magnitude higher than the estimates derived from incubation of samples spiked with  $^{15}\text{NO}_3^-$  (Devol et al., 2006). Contrastingly, anammox process, measured and detected for the first time over the WICS, was found to occur to a much lesser extent ( $\sim 0.06 \pm 0.004 \mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ ). In  $^{15}\text{NH}_4^+$  incubations, production of  $^{30}\text{N}_2$  was not detected thereby ruling out significant involvement of micro-aerobic ammonium oxidation as reported from the Black Sea (Lam et al., 2007). Nevertheless, our anammox rate over the Indian shelf in the eastern Arabian Sea is comparable with the rate measured

off Oman in the western Arabian Sea where this process was found to be the dominant nitrogen loss pathway (Jensen et al., 2011). This indicates that anammox is not limited by the substrate availability as both  $\text{NO}_3^-$  and  $\text{NH}_4^+$  were mostly available in near-bottom waters sampled by us. The growth rate of anammox bacteria are reported to be very slow, with doubling times of 10-30 days in bio-reactors under controlled conditions (Strous et al., 1998). Lower doubling times of 5.5-7.5 days were obtained in specialized membrane bioreactor (Van der Star et al., 2008); however, the growth of anammox organisms is believed to be even slower in the OMZs (Dalsgaard et al., 2012). In the eastern Arabian Sea non-sulphidic (suboxic) anaerobic conditions generally persist only for a few weeks and within this relatively short time anammox bacteria may not be able to build the necessary biomass to utilize the relatively abundant ammonia. In this study DNRA was independently found to occur in some samples in the absence of detectable anammox activity. Along the shelves off Peru and Oman, occurrence of DNRA and its close coupling with anammox has been reported to account for the production of  $^{30}\text{N}_2$  in experiments conducted with  $^{15}\text{NO}_2^-$  (Kartal et al., 2007, Jensen et al., 2011). However, our incubations with  $^{15}\text{NO}_2^- + ^{14}\text{NH}_4^+$  did not show any effect on  $^{30}\text{N}_2$  production compared to incubations with  $^{15}\text{NO}_2^-$  alone. If DNRA-anammox were coupled, then addition of  $^{14}\text{NH}_4^+$  would have diluted the  $^{15}\text{NH}_4^+$  pool thereby resulting in a lower  $^{30}\text{N}_2$  yield (Kartal et al., 2007, Song et al 2016). Furthermore, only ~24% of the samples yielded positive results in our direct measurement of  $^{15}\text{NH}_4^+$  production from  $^{15}\text{NO}_2^-$  in 2009 and 2010, and showed no correlation with the anammox rates. The average DNRA rate ( $109.6 \pm 22.1 \text{ nmol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ ) was higher than that observed over the Omani shelf (highest  $\sim 40 \text{ nmol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ ). This indicates that while DNRA could make some contribution to the observed  $\text{NH}_4^+$  accumulation over the Indian shelf, the sedimentary flux of  $\text{NH}_4^+$  is perhaps more important as Pratihary et al., (2014) reported very high benthic  $\text{NH}_4^+$  efflux ( $3.74 \text{ mmol m}^{-2} \text{ d}^{-1}$ ) based on whole core incubations in this region.

The ambient  $\text{O}_2$  concentration did not show any clear control over denitrification ( $r=0.22$ ,  $p>0.05$ ) and anammox ( $r=0.17$ ,  $p>0.05$ ) in present study. However, it must be noted that the Winkler  $\text{O}_2$  measurements at near-zero levels may result in systematic overestimation. There are several reports on the sensitivity of denitrification to ambient  $\text{O}_2$  concentrations ranging from 0.2 to 5  $\mu\text{M}$  in non sulphidic  $\text{O}_2$ -depleted waters (Dalsgaard et al 2014, and 2012, Bristow et al 2017). For the first step of denitrification - nitrate reduction to nitrite - a somewhat high  $\text{O}_2$  tolerance ( $>25 \mu\text{M}$ ) was noticed from the OMZ waters off Namibia and Peru  $\text{O}_2$  (Kalvalege et al., 2011) than what is inferred from the field data (Thambrup et al., 2012). Earlier observations in the Baltic Sea (Ronner et al., 1985) showed that  $\text{O}_2$  concentrations of  $\sim 9\text{-}11 \mu\text{M}$  could inhibit denitrification; this is supported by the results of Jensen et al. (2009) from the Mariager Fjord. Like denitrification, anammox is also

sensitive to the ambient O<sub>2</sub> concentration. In the Black Sea, anammox bacteria were found to be active at low levels of O<sub>2</sub> (4-8 μM) but showed complete inhibition at 13-14 μM (Jensen et al., 2008). Elsewhere in the oceans tolerance of anammox to O<sub>2</sub> has been found to vary from nanomolar levels in the OMZs to as high as ~15 μM in shallow waters (Kalvalage et al., 2011, Dalsgaard et al., 2012). As stated earlier, the Winkler O<sub>2</sub> data used in some of these studies are not very reliable at vanishingly low levels. However, if real, this variability may be explained either by adaptation of anammox organisms to variable O<sub>2</sub> conditions in coastal waters or by the presence of aggregated particles providing anaerobic micro-niches (Kalvalage et al 2011, Klawonn et al., 2015). Time series observations by Shenoy et al., (2012) carried out for 3 days at station G5 showed highly fluctuating O<sub>2</sub> concentrations on very short time scales (hours), indicating that the anammox and possibly also denitrifying bacteria might be better adapted to these varying O<sub>2</sub> concentrations. Thus, due to multiple uncertainties it is not possible to categorically address the issue of O<sub>2</sub> control on rates measured in the present study.

In many of the incubation experiments carried out by us, an initial lag phase of 10-12 hours occurred before linear increase of <sup>15</sup>N labeled N<sub>2</sub> production. Around 70% of experiments with <sup>15</sup>NO<sub>2</sub><sup>-</sup> amendments in which denitrification were detected and ~30% experiments with <sup>15</sup>NH<sub>4</sub><sup>+</sup> which confirmed anammox activity showed such a lag phase. The most likely explanation of this time lag, also observed by us in similar incubations of anoxic reservoir samples (Naqvi et al., 2018), is that the microbes that perform denitrification and anammox were present in the water but were activated only when O<sub>2</sub> level fell below the threshold (functionally anoxic). The possible causes of the presence of O<sub>2</sub> in our samples subjected to incubation are discussed below. The data from the lag phase were not considered for calculation of rates and only the linear portion of the curve was used for this purpose. In some cases, a large (exponential) increase in labelled N<sub>2</sub> production, departing from the linear trend, was also seen toward the end of the experiments. These data points, ostensibly arising from the so-called "bottle effect", were not used for the rate calculation either. For these reasons, the rates computed by us should be considered as conservative. It is worth mentioning that average denitrification rates estimated from all experiments without lag phases (1.78±0.19 μmol N<sub>2</sub> L<sup>-1</sup> d<sup>-1</sup>) were marginally lower than those from experiments with lag phases (2.74±0.8 μmol N<sub>2</sub> L<sup>-1</sup> d<sup>-1</sup>). Even our lower mean values are more than twice the earlier estimates from the WICS (0.83 μmol N<sub>2</sub> L<sup>-1</sup> d<sup>-1</sup> - Naik, 2003). Contrastingly, in case of anammox, rates calculated from <sup>15</sup>NH<sub>4</sub><sup>+</sup> experiments with lag phase in <sup>29</sup>N<sub>2</sub> production were relatively far lower (0.01±0.003 μmol N<sub>2</sub> L<sup>-1</sup> d<sup>-1</sup>) than those from experiments showing linear production from the beginning (0.08±0.004 μmol N<sub>2</sub> L<sup>-1</sup> d<sup>-1</sup>). This suggests that anammox rates could not have been overestimated.

In isotope pairing technique (IPT) a time lag of ~40 hr in labeled N<sub>2</sub> production has been observed previously in <sup>15</sup>NO<sub>3</sub><sup>-</sup> incubations (Dalsgaard et al., 2003). This can be attributed to the time taken for the conversion of NO<sub>3</sub><sup>-</sup> to NO<sub>2</sub><sup>-</sup> pool before N<sub>2</sub> production starts. In modified IPT involving <sup>15</sup>NO<sub>2</sub><sup>-</sup> as the substrate, time lag could arise from the inhibitory effect of traces of O<sub>2</sub> on denitrification/anammox in the incubated samples. The length of time lag has been found to be directly proportional to the O<sub>2</sub> concentration (Baumann et al., 1997). Results of O<sub>2</sub> amendment experiments by Jensen et al. (2008) revealed that the presence of ~13.5 μM of O<sub>2</sub> could produce 30 hours lag before labeled N<sub>2</sub> production commenced. There are several possibilities of O<sub>2</sub> contamination occurring during experiments involving the IPT. Dalsgaard et al. (2014) pointed out that O<sub>2</sub> could leach out from the PVC samplers to anoxic water samples. However, purging with helium for 15 minutes in 250-ml samples collected in serum bottles should take care of such contamination; this has been practiced as standard operating protocol in many previous studies, both in water from oceanic OMZs (Dalsgaard et al., 2003; Kuypers et al., 2005; Jensen et al., 2008; Holtapells et al., 2011) and anoxic freshwaters (Naqvi et al., 2018). However, there still remains a possibility that some O<sub>2</sub> contamination could occur while transferring samples from serum bottles to Labco exetainers. De Brabandere et al. (2014) also demonstrated that Labco exetainers often used for incubations, as was also done in our study, could suffer from significant O<sub>2</sub> diffusion through their butyl septa. Considering O<sub>2</sub> consumption rates of 0.13 μmol l<sup>-1</sup> h<sup>-1</sup> in <sup>15</sup>NO<sub>2</sub><sup>-</sup> incubation (Jensen et al., 2008), ~1.3 μM O<sub>2</sub> might have produced a time lag of 10-12 hours in our experiments. The initial time lag was seen in samples having measurable O<sub>2</sub> levels (often >10 μM). We strongly believe that He purging for 20 minutes was sufficient to completely remove O<sub>2</sub> initially and that the production of lag phase is most likely due to subsequent contamination. Interestingly, no time lag was observed in the case of samples that contained traces of sulphide, as in our 2009 experiments. We hypothesize that traces of O<sub>2</sub> introduced through contamination could have been quickly utilized for the oxidation of sulphide, but as discussed below we do not rule out the possibility of the occurrence of chemolithotrophic denitrification in the presence of sulphide (Lavik et al., 2009).

Over the Peruvian and Namibian shelves, re-suspension of sediments and presence of copious marine snow are expected to be most conducive for creating particle associated micro-anaerobic niches within which substantial N-loss could occur (Kuypers et al., 2005; Woebken et al., 2007; Kalvelage et al., 2011, 2013). Co-variation of turbidity and anammox rates was observed in these areas. The water column over the western Indian shelf is more turbid (1-4 NTU, measured in 2009), than off Peru and Namibia (0.05-0.1 NTU; Kuypers et al. 2005). In the present study, water turbidity (Fig. 4a) showed statistically significant correlation with denitrification ( $r=0.47$ ,  $p<0.05$ ). This suggests N

removal through denitrification may also occur in particle-associated micro-anaerobic niches over the WICS.

The availability of  $\text{NO}_3^-$  and organic matter are expected to regulate heterotrophic denitrification (Ward et al., 2008, Dalsgaard et al., 2012) as well as anammox (Kalvelage et al., 2013, Callbeck et al 2017) in oceanic oxygen-deficient zones. In our study, in some cases, we noticed high denitrification rates to be associated with higher chlorophyll *a* (a proxy of organic matter availability) as well as lower  $\text{NO}_3^-$  levels (reflecting denitrification/anammox activities).  $\text{NO}_3^-$  showed statistically significant inverse relation ( $r = -0.43$ ,  $p < 0.05$ ), whereas chlorophyll-*a* yielded insignificant correlations with the measured denitrification rates ( $r = 0.20$ ;  $p > 0.05$ ) (Fig. 4b). This indicates that the N loss was not limited by organic matter over the Indian shelf.

Sulphide has been reported to inhibit anammox activity (Dalsgaard et al., 2003; Jensen et al., 2008), and this may account for lower anammox rates in our samples in 2009. Recent studies have also shown that sulphide can act as an electron donor for autotrophic denitrification. Various groups of microbes capable of sulphide driven chemolithoautotrophic denitrification have been detected in anoxic waters of the Baltic Sea (Brettar et al., 2006, Hannig et al 2007), in Saanich Inlet (Zaikova et al., 2010), off Namibia (Lavik et al., 2009) and off Peru (Schunk et al 2013). These bacteria are known to be fast growing and capable of forming blooms within days to weeks (Lavik et al., 2009). Therefore, it is not unlikely that the high rate of denitrification measured in 2009 arises at least in part from autotrophic denitrification by sulphide-oxidizing microbes. However, it must be pointed out that based on results of incubations alone one cannot distinguish between heterotrophic and autotrophic denitrification processes. Nevertheless, since all our denitrification rates were determined in waters with measurable nitrate/nitrite concentrations and were calculated from the linear increase in  $\text{N}_2$  production, it is clear that these processes, whether autotrophic or heterotrophic, is mainly responsible for the apparent N-loss over the Indian shelf. The denitrification rate is of the same order of magnitude but more than double as the  $\text{NO}_3^-$  loss rate observed in the region (Naik, 2003). On the other hand, the estimated anammox rates are much lower.

#### **4.3 N-removal versus uptake over the western Indian continental shelf during seasonal anoxia**

The overall nitrogen removal over the WICS is estimated to range between  $3.70 \pm 0.91$  and  $11.1 \pm 2.72$  Tg annually. This corresponds to ~8-24% of the N-loss estimated for the open ocean OMZ of the Arabian Sea by Bulow et al. (2010) and as high as 24-60% of the model-based value (DeVries et al., 2012). In comparison, anammox contributes only  $0.09 \pm 0.01$ - $0.27 \pm 0.03$  Tg to the N loss per year. Our estimate is also almost three times higher than the previous estimate derived from changes in nitrate

concentrations over time (Naik, 2003). The main shortcoming of this estimate that is based on complete removal of an initial  $\text{NO}_3^-$  concentration of  $\sim 24 \mu\text{M}$  in the upwelled water over a month at a fixed sampling location (Naik, 2003), is that it ignores additional inputs of  $\text{NO}_3^-$ . A number of autochthonous and allochthonous sources of fixed nitrogen exist over WICS during the southwest monsoon season. As mentioned earlier, atmospheric deposition and land runoff contribute significantly towards the nitrogen inventory (Naqvi et al., 2010b, Seitzinger et al., 2005, Bardhan et al., 2015) which is believed to have increased in recent times. Moreover, high (millimolar) concentrations of  $\text{NO}_3^-$  are also found in ground water of Goa very close to coast and the submarine groundwater discharge is expected to be a significant but hitherto poorly quantified source of fixed nitrogen to coastal waters.

A considerable amount of fixed nitrogen is removed by the phytoplankton. The chlorophyll *a* (averaging to  $2.74 \text{ mg m}^{-3}$ ) observed in the present study indicates moderately high primary production. Bhavya et al. (2017) reported a carbon uptake rate of  $38.8 \mu\text{mol C l}^{-1} \text{ d}^{-1}$  from coastal waters of the eastern Arabian Sea during the southwest monsoon. Using the Redfield stoichiometry (Redfield et al., 1963) this would correspond to a fixed N demand of  $5.8 \mu\text{mol. L}^{-1} \text{ d}^{-1}$ . However, most of this nitrogen is expected to be regenerated in subsurface waters and thus made available to microbes facilitating the N loss processes. Once the bottom waters of the WICS turn  $\text{SO}_4^{2-}$  reducing, after the oxidized N has been fully utilized, the nitrogen regenerated is in the reduced form ( $\text{NH}_4^+$ ). However, it is eventually expected to be oxidized to  $\text{NO}_3^-$ . Thus, the nitrogen taken up by the autotrophs does not leave the system. The overall N loss from the system through denitrification and anammox with the former dominating far exceed inputs through upwelling.

## 5. Summary and conclusions

The present study provides the first results, based on direct measurements using the isotope pairing method of denitrification, anammox and DNRA rates over the world's largest naturally formed coastal low- $\text{O}_2$  zone that develops seasonally over the western Indian continental shelf. The results reveal that denitrification is the predominant process responsible for N-loss at rates that are among the highest reported from any aquatic system. Although anammox was also commonly detected in low  $\text{O}_2$  waters its rates were much lower. DNRA was less commonly observed, and at still lower rates. Large spatial and inter annual variations in environmental conditions seem to control the magnitude of different pathways. Denitrification rates were higher over the inner shelf and at the interface of the oxic-suboxic/anoxic layers. Higher rates were observed off Karwar and Mangalore than off Goa. Evolution of sulfate reducing condition and occurrence of dissolved hydrogen sulphide might inhibit anammox activity but stimulate denitrification through chemolithoautotrophy.

As in case of the OMZ off Peru, where the measured N-loss rate is the highest over the shelf but the accumulated  $N^*$  signal is maximal in the offshore OMZ (Kalvelage et al., 2013, Thomsen et al., 2016), strong exchange of water between the coastal and the open ocean OMZ may export N loss signals (i.e.  $N^*$  and  $N_2/Ar$  ratios, Devol et al., 2006; DeVries et al., 2012) to the open ocean in the Arabian Sea as well.

**Authors Contribution:** S.W.A.N. and M.M.M.K. designed the experiments. A.S., S.W.A.N, A.P., G.S. and H.N. performed all experiments. A.S. and S.W.A.N analysed all samples. A.S., S.W.A.N and G.L. processed data. A.S. wrote the manuscript with inputs from S.W.A.N and G.L.

**Acknowledgements:**

We thank all the Master and crew of *CRV Sagar Sukti* and H. Dalvi and A. Methar for extending their assistance in sample collections. Financial support for this research was provided by the European Commission through the award of a Marie Curie International Incoming Fellowship to S. W. A. Naqvi to Max-Planck Society, Germany and the SIBER-INDIA program (Ministry of Earth Sciences, India). The Council of Scientific & Industrial Research (India) supported A. Sarkar and G. Shirodkar through the award of Senior Research Fellowships. Sincere thanks to Dr Rakesh Madhusoodhanan, KISR, Kuwait for support with the statistical analysis.

**Data availability:** The data used in the present study will be archived at National Institute of Oceanography Data Centre ([did.nio.org](http://did.nio.org)) and made available on request to the corresponding author.



## References:

- Barber, R. T., Marra, J., Bidigare, R. C., Codispoti, L. A., Halpern, D., Johnson, Z., Latasa, M., Goericke, R. and Smith, S. L. (2001). Primary productivity and its regulation in the Arabian Sea during 1995. *Deep Sea Research Part II: Topical Studies in Oceanography*, 48(6-7), 1127-1172.
- Bardhan, P., Karapurkar, S. G., Shenoy, D. M., Kurian, S., Sarkar, A., Maya, M. V., Naik, H., Varik, S., Naqvi, S. W. A. (2015). Carbon and nitrogen isotopic composition of suspended particulate organic matter in Zuari Estuary, west coast of India. *Journal of Marine Systems* 141, 90-97.
- Bauman, B., Snozzi, M., Vandermeer, J.R., and Zehnder, A. J. B. (1997). Development of stable denitrifying cultures during repeated aerobic-anaerobic transient periods, *Wat.Res.*, 31, 1947–1954.
- Bendschneider, K., Robinson, R. J. (1952). A new spectrophotometric method for the determination of nitrite in sea water. *Journal of Marine Research* 11, 87-96.
- Bhavya, P. S., Kumar, S., Gupta, G. V. M. and Sudheesh, V. (2017). Carbon uptake rates in the Cochin estuary and adjoining coastal Arabian Sea. *Estuaries and Coasts*, 40(2), 447-456.
- Brettar, I., Labrenz, M., Flavier, S., Botel, J., Kuosa, H., Christen, R., Höfle, M. G. (2006). Identification of a *Thiomicrospira denitrificans* like epsilon proteo bacterium as a catalyst for autotrophic denitrification in the central Baltic Sea. *Applied and Environmental Microbiology* 72(2), 1364-1372.
- Bristow, L.A., Callbeck, C.M., Larsen, M., Altabet, M. A., Dekaezemacker, J., Forth, M., Gauns, M., Glud, R.N., Kuypers, M.M., Lavik, G., Milucka, J., Naqvi, S. W. A., Pratihary, A., Revsbech, N. P., Thamdrup, B., A. Treusch, H., Canfield, D. E. (2017). N<sub>2</sub> production rates limited by nitrite availability in the Bay of Bengal oxygen minimum zone. *Nature Geoscience* 10(1), p.24.
- Bulow, S.E., Rich, J.J., Naik, H.S., Pratihary, A. K., Ward, B.B. (2010). Denitrification exceeds anammox as a nitrogen loss pathway in the Arabian Sea oxygen minimum zone. *Deep Sea Research Part I: Oceanographic Research Papers* 57(3), 384-393.
- Callbeck, C.M. (2017). Distribution and Activity of Anammox and Sulphide-oxidizing Nitrate-reducing Bacteria in Oxygen Minimum Zones. Doctoral dissertation, Universität Bremen, Bremen, Germany.
- Dalsgaard, T., Canfield, D.E., Petersen, J., Thamdrup, B., Acuña-González, J. (2003). N<sub>2</sub> production by the anammox reaction in the anoxic water column of Golfo Dulce, Costa Rica. *Nature* 422(6932), 606-608.
- Dalsgaard, T., Thamdrup, B., Farías, L., Revsbech, N. P. (2012). Anammox and denitrification in the oxygen minimum zone of the eastern South Pacific. *Limnology and Oceanography* 57(5), 1331-1346.
- Dalsgaard, T., Stewart, F.J., Thamdrup, B. De Brabandere, L., Revsbech, N.P., Ulloa, O., Canfield, D.E., DeLong E.F. (2014). Oxygen at nanomolar levels reversibly suppresses process rates and gene expression in anammox and denitrification in the oxygen minimum zone off northern Chile. *MBio* 5(6), pp.e01966-14.

- De Brabandere, L., Canfield, D. E., Dalsgaard, T., Friederich, G. E., Revsbech, N. P., Ulloa, O., and Thamdrup, B. (2014): Vertical partitioning of nitrogen-loss processes across the oxic-anoxic interface of an oceanic oxygen minimum zone, *Environ. Microbiol.*,16(10), 3041-3054.
- Devol, A.H., Uhlenhopp, A.G., Naqvi, S.W.A., Brandes, J.A., Jayakumar, D.A., Naik, H., Gaurin, S., Codispoti, L.A., Yoshinari, T. (2006). Denitrification rates and excess nitrogen gas concentrations in the Arabian Sea oxygen deficient zone. *Deep Sea Research Part I: Oceanographic Research Papers* 53(9), 1533-1547.
- DeVries, T, Deutsch, C., Primeau, F., Chang, B., Devol, A. (2012). Global rates of water-column denitrification derived from nitrogen gas measurements. *Nature Geoscience* 5(8), 547-550.
- Duce, R.A., LaRoche, J., Altieri, K., Arrigo, K.R., Baker, A.R., Capone, D.G., Cornell, S., Dentener, F., Galloway, J., Ganeshram, R.S., Geider, R.J., Jickells, T., Kuypers, M. M. M., Langlois, R., Liss, P. S., Liu, S. M., Middelburg, J. J., Moore, C. M., Nickovic, S., Oschlies, A., Pedersen, T., Prospero, J., Schlitzer, R., Seitzinger, S., Sorensen, L. L., Uematsu, M., Ulloa, O., Voss, M., Ward, B.B., Zamora L. (2008). Impacts of atmospheric anthropogenic nitrogen on the open ocean. *Science* 320(5878), pp.893-897.
- Froelich, P.N., Klinkhammer, G.P., Bender, M.L., Luedtke, N.A., Heath, G.R., Cullen, D., Dauphin, P., Hammond, D., Hartman, B., Maynard, V. (1979). Early oxidation of organic matter in pelagic sediments of eastern equatorial Atlantic: suboxic diagenesis. *Geochimica et Cosmochimica Acta* 43(7), 1075-1090.
- Grasshoff, K., Ehrhardt, M., Kremling K. (1983). *Methods of seawater analysis*. Verlag Chemie. Weinheim, Germany 4(1), 9.
- Gruber, N., Sarmiento J.L. (1997). Global patterns of marine nitrogen fixation and denitrification. *Global Biogeochemical Cycles* 11(2), 235-266.
- Hannig, M., Lavik, G., Kuypers, M.M.M., Woebken, D., Martens-Habbena, W., Jürgens, K. (2007). Shift from denitrification to anammox after inflow events in the central Baltic Sea. *Limnology and Oceanography* 52(4), 1336-1345.
- Holtappels, M., Lavik, G., Jensen, M.M., Kuypers, M.M.M. (2011). <sup>15</sup>N-labelling experiments to dissect the contributions of heterotrophic denitrification and anammox to nitrogen removal in the OMZ waters of the ocean. *Methods in Enzymology* 486, 223-251.
- Jensen, M.M., Kuypers, M.M.M., Lavik, G., Thamdrup B. (2008). Rates and regulation of anaerobic ammonium oxidation and denitrification in the Black Sea. *Limnology and Oceanography* 53(1), 23-36.
- Jensen, M. M., Lam, P., Revsbech, N. P., Nagel, B., Gaye, B., Jetten, M. S. M., Kuypers, M. M. M. (2011). Intensive nitrogen loss over the Omani Shelf due to anammox coupled with dissimilatory nitrite reduction to ammonium. *The ISME journal* 5(10), 1660-1670.
- Kalvelage, T., Jensen, M.M., Contreras, S., Revsbech, N.P., Lam, P., Günter, M., LaRoche, J., Lavik, G., Kuypers, M. M. M. (2011). Oxygen sensitivity of anammox and coupled N-cycle processes in oxygen minimum zones. *PLoS One* 6(12), p.e29299.

- Kalvelage, T., Lavik, G., Lam, P., Contreras, S., Arteaga, L., Löscher, C. R., Oeschies, A., Paulmier, A., Stramma, L., Kuypers, M.M. (2013). Nitrogen cycling driven by organic matter export in the South Pacific oxygen minimum zone. *Nature geoscience* 6(3), 228-234.
- Kartal, B., Kuypers, M.M.M., Lavik, G., Schalk, J., Op den Camp, H.J.M., Jetten, M.S.M., Strous, M. (2007). Anammox bacteria disguised as denitrifiers: nitrate reduction to dinitrogen gas via nitrite and ammonium. *Environmental microbiology* 9(3), 635-642.
- Klawonn, I., Bonaglia, S., Brüchert, V., Ploug, H. (2015). Aerobic and anaerobic nitrogen transformation processes in N<sub>2</sub>-fixing cyanobacterial aggregates. *The ISME journal* 9(6), p.1456.
- Kuypers, M.M.M., Sliemers, A.O., Lavik, G., Schmid, M., Jørgensen, B.B., Kuenen, J. G., Damsté, J. S., Strous, S. M., Jetten, M. S. M. (2003). Anaerobic ammonium oxidation by anammox bacteria in the Black Sea. *Nature* 422(6932), 608-611.
- Kuypers, M. M. M., Lavik, G., Woebken, D., Schmid, M., Fuchs, B.M., Amann, R., Jørgensen, B. B., Jetten M. S. M., (2005). Massive nitrogen loss from the Benguela upwelling system through anaerobic ammonium oxidation. *Proceedings of the National Academy of Sciences of the United States of America* 102(18), 6478-6483.
- Lam, P., M. Jensen, M., Lavik, G., McGinnis, D.F., Müller, B., Schubert, C.J., Amann, R., Thamdrup, B., Kuypers, M. M. M. (2007). Linking crenarchaeal and bacterial nitrification to anammox in the Black Sea. *Proceedings of the National Academy of Sciences of the United States of America* 104(17), 7104-7109.
- Lavik, G., Stührmann, T., Brüchert, V., Van der Plas, A., Mohrholz, V., Lam, P., Mußmann, M., Fuchs, B. M., Amann, R., Lass, U., Kuypers, M. M. M. (2009). Detoxification of sulphidic African shelf waters by blooming chemolithotrophs. *Nature*, 457(7229), 581-584.
- Maya, M. V., Karapurkar, S. G., Naik, H., Roy, R., Shenoy, D. M., and Naqvi, S. W. A. (2011). Intraannual variability of carbon and nitrogen stable isotopes in suspended organic matter in waters of the western continental shelf of India, *Biogeosciences*, 8, 3441–3456.
- Naik, H. (2003). Benthic nitrogen cycling with special reference to nitrous oxide in the coastal and continental shelf environments of the eastern Arabian Sea. Dissertation thesis, Marine Science Department, Goa University, Taleigao Plateau, Goa, India.
- Naqvi, S.W. A., Jayakumar, D. A., Narvekar, P. V., Naik, H., Sarma, V. V. S. S., D'Souza, W., Joseph, S., George, M. D. (2000). Increased marine production of N<sub>2</sub>O due to intensifying anoxia on the Indian continental shelf. *Nature* 408(6810), 346-349.
- Naqvi, S. W. A., Naik, H., Jayakumar, D. A., Shailaja, M. S., Narvekar P. V. (2006a). Seasonal oxygen deficiency over the western continental shelf of India. In: Neretin, L. (Eds.), *Past and Present Water Column Anoxia*, NATO Science Series, IV. Earth and Environmental Sciences 64, Springer, Dordrecht, Netherlands, pp. 195–224.
- Naqvi, S.W.A., Naik, H., Pratihary, A.K., DeSouza, W., Narvekar, P.V., Jayakumar, D.A., Devol, A.H., Yoshinari, T. and Saino, T. (2006b). Coastal versus open-ocean denitrification in the Arabian Sea. *Biogeosciences*, 3, 621–633.
- Naqvi, S.W.A., Naik, H., Jayakumar, A., Pratihary, A.K., Narvenkar, G., Kurian, S., Agnihotri, R., Shailaja, M.S., Narvekar, P.V. (2009). Seasonal anoxia over the western Indian continental shelf. In:

Wiggert, J. D., Hood, R.R., Naqvi, S.W.A., Brink, K.H., Smith, S. L. (Eds.), Indian Ocean Biogeochemical processes and ecological variability 1, pp. 333-345.

Naqvi, S. W. A., Bange, H. W., Farías, L., Monteiro, P. M. S., Scranton, M. I., Zhang, J. (2010a). Marine hypoxia/anoxia as a source of CH<sub>4</sub> and N<sub>2</sub>O. *Biogeosciences* 7(7), 2159-2190.

Naqvi, S. W. A., Naik, H., D' Souza, W., Narvekar, P. V., Paropkari, A. L., and Bange, H. W. (2010b). Carbon and nitrogen fluxes in the North Indian Ocean, in: *Carbon and Nutrient Fluxes in Continental Margins: A Global Synthesis*, edited by: Liu, K. K., Atkinson, L., Quinones, R., and Talaue McManus, L., Springer- Verlag, Berlin.

Naqvi, S. W. A., Lam, P., Narvenkar, G., Sarkar, A., Naik, H., Pratihary, A., Shenoy, D. M., Gauns, M., Kurian, S., Damare, S., Duret, M., Lavik, G., Kuypers, M. M. M. (2018). Methane stimulates massive nitrogen loss from freshwater reservoirs in India. *Nature communications*, 9(1), 1-10.

Nielsen, L. P. (1992). Denitrification in sediment determined from nitrogen isotope pairing. *FEMS Microbiology Letters* 86(4), 357-362.

Pratihary, A.K., Naqvi, S.W.A., Narvenkar, G., Kurian, S., Naik, H., Naik R., Manjunatha, B.R. (2014). Benthic mineralization and nutrient exchange over the inner continental shelf of western India. *Biogeosciences* 11(10), 2771-2791.

Preisler, A., De Beer, D., Lichtschlag, A., Lavik, G., Boetius A., Jørgensen B.B. (2007). Biological and chemical sulphide oxidation in a *Beggiatoa* inhabited marine sediment. *The ISME journal* 1(4), 341-353.

Rabalais, N.N., Diaz, R.J., Levin, L.A., Turner, R.E., Gilbert, D. and Zhang, J. (2010). Dynamics and distribution of natural and human-caused hypoxia. *Biogeosciences*, 7(2), p.585.

Redfield, A. C., Ketchum, B. H. and Richards, F. A. (1963). The influence of organisms on the composition of seawater, in *Thesea*, eds by, Interscience, New York, M. N. Hill, 2, 26-77.

Schlitzer, R. (2018). Ocean Data View, [odv.awi.de](http://odv.awi.de). Schott, F., McCreary, J. P. (2001). The monsoon circulation of the Indian Ocean. *Progress in Oceanography* 51(1), 1-123.

Schunck, H., Lavik, G., Desai, D.K., Großkopf, T., Kalvelage, T., Löscher, C.R., Paulmier, A., Contreras, S., Siegel, Holtappels H., M., Rosenstiel, P. (2013). Giant hydrogen sulphide plume in the oxygen minimum zone off Peru supports chemolithoautotrophy. *PLoS One* 8(8), p.e68661.

Seitzinger, S. P., Harrison, J. A., Dumont, E., Beusen, A. H. W., Bouwman A. F. (2005). Sources and delivery of carbon, nitrogen, and phosphorus to the coastal zone: An overview of Global Nutrient Export from Watersheds (NEWS) models and their application. *Global Biogeochemical Cycles* 19 (4).

Shankar, D., Shenoi, S.S.C., Nayak, R.K., Vinayachandran, P.N., Nampoothiri, G., Almeida, A.M., Michael, G.S., Ramesh Kumar, M.R., Sundar, D. and Sreejith, O.P. (2005). Hydrography of the eastern Arabian Sea during summer monsoon 2002, *J. Earth Syst. Sci.*, 114, 459-474.

Shenoy, D.M., Sujith, K.B., Gauns, M., Patil, S., Sarkar, A., Naik, H., Narvekar, P.V., Naqvi, S.W.A. (2012). Production of dimethylsulphide during the seasonal anoxia off Goa. *Biogeochemistry* 110(1-3), 47-55.

- Song, G.D., Liu, S.M., Kuypers, M.M.M., Lavik G. (2016). Application of the isotope pairing technique in sediments where anammox, denitrification, and dissimilatory nitrate reduction to ammonium coexist. *Limnology and Oceanography: Methods* 14(12), pp.801-815.
- Stramma, L., Fischer, J., Schott, F. (1996). The flow field off southwest India at 8N during the southwest monsoon of August 1993. *Journal of Marine Research*, 54(1), 55-72.
- Strickland, J.D.H., Parsons, T.R. (1972). *A Practical Handbook of Seawater Analysis*. Fisheries Research Board of Canada 167, pp 31.
- Strous, M., Heijnen, J.J., Kuenen, J.G., Jetten M.S.M. (1998). The sequencing batch reactor as a powerful tool for the study of slowly growing anaerobic ammonium-oxidizing microorganisms. *Applied microbiology and biotechnology* 50(5), pp.589-596.
- Thamdrup, B., Dalsgaard, T. (2002). Production of N<sub>2</sub> through anaerobic ammonium oxidation coupled to nitrate reduction in marine sediments. *Applied and environmental microbiology* 68(3), 1312-1318.
- Thamdrup, B., Dalsgaard, T. and Revsbech, N.P. (2012). Widespread functional anoxia in the oxygen minimum zone of the Eastern South Pacific. *Deep Sea Research Part I: Oceanographic Research Papers*, 65, pp.36-45.
- Thomsen, S., Kanzow, T., Krahnemann, G., Greatbatch, R.J., Dengler M., Lavik, G. (2016). The formation of a subsurface anticyclonic eddy in the Peru-Chile Undercurrent and its impact on the near-coastal salinity, oxygen, and nutrient distributions. *Journal of Geophysical Research: Oceans* 121(1), 476-501.
- Van der star, W. R. L., Miclea, A. I., Van dongen, U. G. J. M., Muyzer, G., Picioreanu, C., Van Loosdrecht, M. C. M. (2008). The membrane bioreactor: A novel tool to grow anammox bacteria as free cells. *Biotechnology and Bioengineering* 101(2), 286-294.
- Ward, B.B., Tuit, C. B., Jayakumar, A., Rich, J.J., Moffet, J., Naqvi, S.W.A. (2008). Organic carbon, and not copper, controls denitrification in oxygen minimum zones of the ocean. *Deep Sea Research Part I: Oceanographic Research Papers* 55(12), 1672-1683.
- Warembourg, F. R. (1993). Nitrogen fixation in soil and plant system. In: Knowles, R., Blackburn, T. H., (Eds.), *Nitrogen Isotopes Techniques*, pp.127-156.
- Woebken, D., B. Fuchs, M., Kuypers M.M.M., Amann, R. (2007). Potential interactions of particle-associated anammox bacteria with bacterial and archaeal partners in the Namibian upwelling system. *Applied and environmental microbiology* 73(14), 4648-4657.
- Zaikova, E., Walsh, D.A., Stilwell, C.P., Mohn, W. W., Tortell, P.D., Hallam, S.J. (2010). Microbial community dynamics in a seasonally anoxic fjord: Saanich Inlet., British Columbia. *Environmental microbiology* 12(1), 172-191.

### **Table Captions:**

**Table 1:** Calculated rates (mean and stdev values) of different nitrogen transformation process (anammox rates obtained from both  $^{15}\text{NH}_4^+$  and  $^{15}\text{NO}_2^-$  incubations) along with ancillary chemical data.

**Table 2:** Total nitrogen loss (mean and stdev values) from eastern Arabian Sea from present study and comparison with earlier data extrapolated to the OMZ area and thickness adopted from Naqvi et al., (2000).

Year	Transects	Site	Depth (m)	O <sub>2</sub> (μM)	NO <sub>3</sub> <sup>-</sup> (μM)	NO <sub>2</sub> <sup>-</sup> (μM)	NH <sub>4</sub> <sup>+</sup> (μM)	Denitrification (μmol N <sub>2</sub> L <sup>-1</sup> d <sup>-1</sup> )	Anammox (μmol N <sub>2</sub> L <sup>-1</sup> d <sup>-1</sup> ) based on <sup>15</sup> NH <sub>4</sub> <sup>+</sup> incubation	Anammox (μmol N <sub>2</sub> L <sup>-1</sup> d <sup>-1</sup> ) based on <sup>15</sup> NO <sub>2</sub> <sup>-</sup> incubation	DNRA (μmolN <sub>2</sub> L <sup>-1</sup> d <sup>-1</sup> )		
2008	Goa	G6	30	11.1	3.20	2.7	1.58	1.90±0.16	0.02±0.004	0.52±0.1	n/m		
			35	13.9	4.08	2.6	0.06	0.91±0.08	0.01±0.003	0.68±0.001	n/m		
		G7	35	10.4	4.05	4.4	0.89	1.93±0.24	0.01±0.001	0.89±0.1	n/m		
			40	BDL	11.9	3.03	0.24	1.21±0.95	nd	n/m	n/m		
2009	Goa	G5	18	BDL	2.19	2.1	n/m	4.37±0.20	nd	n/m	nd		
			25	BDL	0.98	0.07	n/m	8.62±1.10	nd	n/m	nd		
		G6	18	32.0	12.6	1.9	n/m	0.45±0.01	nd	n/m	nd		
			25	10.5	4.9	0.8	n/m	1.15±0.48	detected	0.91±0.45	nd		
		G7	35	2.88	4.67	1.11	n/m	2.25±0.37	nd	n/m	0.32±0.07		
			25	4.4	12.6	0.8	n/m	0.53±0.13	detected	0.20±0.10	Nd		
		G7	35	BDL	15.9	1.1	n/m	1.31±0.27	detected	0.50±0.21	0.03±0.01		
			45	4.5	12.4	0.6	n/m	0.90±0.04	detected	0.01±0.05	nd		
		2009	Karwar	K1	15	BDL	1.6	3.08	n/m	9.29±4.52	detected	nd	nd
					20	BDL	2.7	2.83	n/m	7.75±2.25	nd	n/m	nd
K2	10			24.7	n/m	6.4	n/m	9.34±1.32	nd	n/m	0.14±0.02		
	20			7.3	n/m	6.8	n/m	0.66±0.10	nd	n/m	nd		
K3	30			3.9	n/m	6.8	n/m	0.25±0.03	detected	0.07±0.02	0.19±0.009		
	18			39.9	18.2	0.5	n/m	0.15±0.03	nd	n/m	nd		
K3	25			5.7	13.6	3.5	n/m	0.33±0.06	nd	n/m	nd		
	35			2.9	9.5	2.7	n/m	0.19±0.03	nd	n/m	nd		
44	3.7	11.4	4.9	n/m	0.21±0.01	nd	n/m	0.02±0.006					
2009	Mangalore	M1	24	BDL	n/m	2.5	n/m	10.03±1.7	nd		0.1±0.02		
2010	Goa	G5	15	17.9	1.1	0.5	8.3	3.54±2.3	0.02±0.001	0.08±0.29	nd		
			25	4.7	1.9	0.1	9.1	0.43±0.08	0.43±0.02	0.79±0.12	0.15±0.006		
		G6	30	3.8	4.4	5.5	3.7	0.61±0.02	0.003±0.001	0.35±0.15	nd		
			35	3.8	2.6	2.7	4.0	0.24±0.1	0.01±0.003	0.33±0.21	nd		
		G7	40	3.9	4.6	0.4	2.1	2.83±0.66	0.24±0.008	1.88±0.43	nd		
	Karwar	K2	22	3.8	0.9	1.1	3.3	3.05±1.16	0.014±0.002	1.19±0.68	nd		
			26	2.48	5.7	0.1	4.3	0.12±0.03	0.011±0.001	0.12±0.03	nd		
		K3	35	6.7	15.7	1.2	3.1	0.03±0.006	0.03±0.01	0.05±0.02	nd		
			50	2.0	2.7	0.1	5.2	0.03±0.004	0.01±0.001	0.04±0.02	nd		
	Mangalore	M2	20	n/m	11.8	5.9	1.8	3.66±2.11	0.03±0.001	nd	0.04±0.002		
35			17.6	15.8	8.4	2.2	2.77±0.50	0.05±0.001	nd	nd			

**Table 1:**

n/m= = not measured, nd= not detected; BDL= below detection limit

**Table 2:**

Source	Area of Coastal OMZ (Km <sup>2</sup> )	Thickness of OMZ (m)	Average Denitrification ( $\mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ )	Average Anammox ( $\mu\text{mol N}_2 \text{ L}^{-1} \text{ d}^{-1}$ )	Removal of Fixed Nitrogen ( $\text{Tg N yr}^{-1}$ )	
					Denitrification	Anammox
Present Study	60-180	20	2.45±0.6	0.06±0.004	3.70±0.91-11.11±2.72	0.09±0.01-0.27±0.03
Devol et al.,2006	60-180	20	0.033	n/m	0.05-0.15	n/m
Naik et al., 2003	60-180	20	0.83±0.6	n/m	1.3-3.8	n/m

n/m = not measured



### Figure captions:


**Fig. 1:** Study area in the eastern Arabian Sea along western continental shelf of India. Biogeochemical monitoring locations on each coast perpendicular transects are indicated by closed circle points. The stations marked with 'red arrows' are where  $^{15}\text{N}$ -incubations experiments were performed.

**Fig. 2:** Distribution of physicochemical properties including potential density ( $\sigma_\theta$ ), dissolved oxygen ( $\text{O}_2$ ), nitrate ( $\text{NO}_3^-$ ), nitrite ( $\text{NO}_2^-$ ) and nitrogen deficit ( $\text{N}^*$ ) in the water column along coast perpendicular transect off Goa based on observations carried out during three consecutive years Plate A. 2008, Plate B. 2009 and Plate C. 2010. Depths marked as 'diamond- $\diamond$ ' on  $\text{O}_2$  panels in all three years represents the depths where  $^{15}\text{N}$ -incubation experiments were conducted.


**Fig. 3:** Depth wise variations in concentrations of dissolved oxygen, nitrate, nitrite and ammonium (connected vertical profiles) and  $\text{N}_2$  production rates measured by isotope pairing experiments (horizontal bars): (a) G5 inner shelf station off Goa( i)2009 and (ii) 2010 and (b) G6 mid shelf station off Goa during (i)2008, (ii) 2009 and (iii) 2010

**Fig. 4:** Correlations between (a) denitrification versus turbidity and (b) denitrification versus nitrate (blue circles) and chlorophyll *a* (green circle)

Figure 1:



**Figure 2:**




**Figure 3:**

**a. G5 (Inner shelf )**

(i) 2009

(ii) 2010



**b. G6 (Mid shelf)**

(i) 2008

(ii) 2009

(iii) 2010




Figure 4:

