

Appendix 1 Average ( $\pm$  SD) values of abiotic parameters of the three estuaries at each salinity zone (Eu; Euhaline, Poly-meso; Poly-mesohaline, Oligo; Oligohaline) across three seasons (Pr; Premonsoon, M; Monsoon, Po; Postmonsoon)

		Ulhas			Amba			Savitri		
		Pr	M	Po	Pr	M	Po	Pr	M	Po
Salinity	Eu	35.1 $\pm$ 1.4	34.2	34.8	35.0 $\pm$ 0.9	32.4 $\pm$ 1.5	32.8 $\pm$ 1.5	32.6 $\pm$ 2.2	34.0 $\pm$ 2.1	32.8 $\pm$ 1.3
	Poly-meso	15.6 $\pm$ 6.5	16.8	17.6 $\pm$ 7.7	19.4 $\pm$ 8.4	17.2 $\pm$ 9.8	22.3 $\pm$ 5.2	24.0 $\pm$ 4.4	19.4	16.3 $\pm$ 7.3
	Oligo	2.5 $\pm$ 2.2	0.2 $\pm$ 0.6	1.3 $\pm$ 1.6	-	0.2 $\pm$ 0.2	2.7	0.1	0.1 $\pm$ 0.2	0
DO (mg l $^{-1}$ )	Eu	5.3 $\pm$ 0.9	5.3	4.9	5.3 $\pm$ 0.6	4.2 $\pm$ 1.1	5.3 $\pm$ 0.5	5.0 $\pm$ 0.4	4.5 $\pm$ 0.9	6.2 $\pm$ 0.2
	Poly-meso	2.9 $\pm$ 0.5	4.4	3.0 $\pm$ 1.3	5.2 $\pm$ 0.0	5.3 $\pm$ 0.7	4.9 $\pm$ 1.1	4.8 $\pm$ 2.2	4.8	6.5 $\pm$ 0.8
	Oligo	5.8 $\pm$ 2.8	5.5 $\pm$ 1.1	4.0 $\pm$ 4.0	-	7.2 $\pm$ 0.6	6.9	2.9	7.2 $\pm$ 0.8	8.1
NO <sub>2</sub> <sup>-</sup> -N ( $\mu$ mol l $^{-1}$ )	Eu	0.2 $\pm$ 0.2	1.2	0.3	2.7 $\pm$ 3.7	1.2 $\pm$ 0.2	2.1 $\pm$ 1.7	0.9 $\pm$ 1.1	0.2 $\pm$ 0.1	1.2 $\pm$ 0.8
	Poly-meso	16.9 $\pm$ 12.8	2.1	16.8 $\pm$ 6.1	5.5 $\pm$ 5.0	0.8 $\pm$ 0.3	0.5 $\pm$ 0.3	6.9 $\pm$ 6.4	0.1	4.6 $\pm$ 4.1
	Oligo	3.0 $\pm$ 2.8	0.9 $\pm$ 0.7	3.9 $\pm$ 3.7	-	0.7 $\pm$ 0.2	2.4	0.01	0.2 $\pm$ 0.0	0.1
NO <sub>3</sub> <sup>-</sup> -N ( $\mu$ mol l $^{-1}$ )	Eu	46.4 $\pm$ 41.7	12.9	16.7	22.4 $\pm$ 10.4	9.0 $\pm$ 3.1	17.1 $\pm$ 6.7	6.2 $\pm$ 4.6	1.2 $\pm$ 0.0	6.0 $\pm$ 2.7
	Poly-meso	32.9 $\pm$ 14.6	15.4	46.4 $\pm$ 43.8	10.4 $\pm$ 10.8	9.8 $\pm$ 2.9	12.2 $\pm$ 3.1	26.8 $\pm$ 6.7	3.3	17.0 $\pm$ 7.7
	Oligo	16.7 $\pm$ 10.3	4.5 $\pm$ 1.3	36.2 $\pm$ 25.1	-	4.7 $\pm$ 2.4	11.9	3.3	2.4 $\pm$ 1.0	1.2
NH <sub>4</sub> <sup>+</sup> -N ( $\mu$ mol l $^{-1}$ )	Eu	1.6 $\pm$ 1.0	1.5	1.6	1.1 $\pm$ 0.8	2.9 $\pm$ 0.4	3.7 $\pm$ 1.4	1.1 $\pm$ 0.3	2.8 $\pm$ 0.7	1.9 $\pm$ 0.7
	Poly-meso	11.2 $\pm$ 11.1	4.8	34.0 $\pm$ 27.3	0.8 $\pm$ 0.0	3.8 $\pm$ 2.3	4.3 $\pm$ 1.7	3.8 $\pm$ 3.1	2.7	4.2 $\pm$ 1.5
	Oligo	6.9 $\pm$ 5.9	8.0 $\pm$ 6.7	43.4 $\pm$ 38.9	-	4.8 $\pm$ 3.6	10.8	0.5	4.5 $\pm$ 2.6	4.1
Sand (%)	Eu	30.9 $\pm$ 39.4	11.3	7.2	10.3 $\pm$ 7.2	6.8 $\pm$ 1.1	5.4 $\pm$ 7.8	73.9 $\pm$ 46.0	39.6 $\pm$ 55.0	46.6 $\pm$ 65.7
	Poly-meso	22.6 $\pm$ 33.0	19.8	10.9 $\pm$ 6.3	73.9 $\pm$ 0.6	12.7 $\pm$ 7.3	8.8 $\pm$ 6.8	56.1 $\pm$ 47.4	82	65.9 $\pm$ 35.4
	Oligo	32.9 $\pm$ 42.8	17.8 $\pm$ 19.6	26.2 $\pm$ 26.8	-	7.9 $\pm$ 6.7	5.1	3	57.7 $\pm$ 45.5	28.8
Silt (%)	Eu	63.9 $\pm$ 37.1	82.3	85.2	71.8 $\pm$ 12.0	80.0 $\pm$ 3.5	85.2 $\pm$ 13.7	22.7 $\pm$ 42.8	58.8 $\pm$ 56.3	48.3 $\pm$ 62.5
	Poly-meso	66.5 $\pm$ 30.3	65.2	72.3 $\pm$ 10.7	73.9 $\pm$ 0.6	69.4 $\pm$ 11.7	70.4 $\pm$ 15.4	31.3 $\pm$ 33.1	13.9	21.0 $\pm$ 23.7
	Oligo	57.8 $\pm$ 36.1	70.0 $\pm$ 18.6	60.4 $\pm$ 20.9	-	72.1 $\pm$ 11.2	77.1	85.4	32.5 $\pm$ 35.6	65.7
Clay (%)	Eu	5.2 $\pm$ 2.4	6.4	7.6	18.0 $\pm$ 6.7	13.2 $\pm$ 2.4	9.4 $\pm$ 6.1	3.4 $\pm$ 3.3	1.6 $\pm$ 1.3	5.2 $\pm$ 3.2
	Poly-meso	10.9 $\pm$ 6.9	15	16.8 $\pm$ 6.9	18.2 $\pm$ 1.9	17.9 $\pm$ 6.6	20.9 $\pm$ 10.5	12.6 $\pm$ 14.5	4.1	13.1 $\pm$ 12.3
	Oligo	9.2 $\pm$ 6.8	12.2 $\pm$ 9.7	13.4 $\pm$ 6.1	-	20.0 $\pm$ 4.9	17.8	11.6	9.8 $\pm$ 14.5	5.5
C <sub>org</sub> (%)	Eu	1.1 $\pm$ 0.5	1.8	1.9	2.1 $\pm$ 0.5	1.6 $\pm$ 0.4	2.1 $\pm$ 0.5	0.6 $\pm$ 0.8	1.1 $\pm$ 1.3	1.2 $\pm$ 1.1
	Poly-meso	2.2 $\pm$ 1.2	2.3	2.1 $\pm$ 0.3	2.0 $\pm$ 0.2	2.0 $\pm$ 0.4	2.0 $\pm$ 0.1	1.4 $\pm$ 1.3	0.1	0.9 $\pm$ 0.8
	Oligo	3.4 $\pm$ 2.2	2.5 $\pm$ 1.0	3.0 $\pm$ 1.2	-	2.1 $\pm$ 0.5	2.1	2.5	1.3 $\pm$ 1.3	3.6

## Appendix 2 Ecological group (EG) classification of taxa recorded at three studied estuaries (Ulhas, Amba and Savitri).

<b>Ecological group I</b>	<i>Grandidierella</i> sp.*	<b>Ecological group II</b>	<i>Nephthys hystericis</i>	<i>Neanthes chilkaensis</i> *	<i>Lagis koreni</i>	<b>Not assigned taxa</b>
<i>Abra opalina</i> *	Holothuroidea	<i>Alpheus</i> sp.	<i>Ninoe</i> sp.	<i>Nemertea</i>	<i>Linopherus acarunculatus</i> *	<i>Baharilana richmondi</i>
<i>Acetes indicus</i> *	<i>Idunella chilkensis</i> *	<i>Ampelisca diadema</i>	<i>Ophicaris</i> sp.	<i>Nereis falcaria</i>	<i>Mediomastus capensis</i>	<i>Branchiodrilus</i> sp.
<i>Acetes</i> sp.*	<i>Idunella picta</i>	<i>Amphilochus</i> sp.	<i>Paraehlersia ferruginea</i>	<i>Nereis</i> sp.	<i>Namalyctis indica</i>	<i>Cloridina ichneumon</i>
<i>Aglaophamus dibranchis</i>	<i>Idunella</i> sp.*	<i>Amphiura</i> sp.	<i>Phoronis psammophila</i>	<i>Notomastus</i> sp. *	<i>Namalyctis senegalensis</i>	<i>Corophidae</i>
<i>Ampelisca aequicornis</i>	<i>Iphinoe</i> sp.	Anthozoa	<i>Podocerus spongicolus</i> *	<i>Oncholaimus oxyuris</i> *	<i>Parapriopospio cordifolia</i> *	<i>Dotillidae</i>
<i>Ampelisca scabripes</i> *	<i>Laternula</i> sp.	<i>Aulophorus furcatus</i> *	<i>Pycnogonida</i>	<i>Pagurapseudopsis gymnochobia</i>	<i>Parapriopospio cristata</i> *	<i>Eucopiidae</i>
<i>Ampelisca</i> sp.	<i>Lembos</i> sp.	<i>Bodotria</i> sp.*	<i>Siliqua</i> sp.	<i>Paradoneis lyra capensis</i>	<i>Polycirrus</i> sp.	<i>Exhippolysmata</i> sp.
<i>Ampelisca spinipes</i>	<i>Lembos websteri</i>	<i>Caprella</i> sp.	<i>Sinelobus stanfordi</i>	<i>Paraonis fulgens</i>	<i>Polydora ciliata</i>	<i>Gitanopsis gouriae</i>
<i>Ampelisca typica</i>	<i>Leucothoe richiardii</i>	<i>Cirolana aldabrensis</i>	<i>Stenothoe gallensis</i> *	<i>Podocerus brasiliensis</i>	<i>Prionospio cirrifera</i>	<i>Gitanopsis</i> sp.
<i>Amphipholis misera</i> *	<i>Lygdamis indicus</i>	<i>Clibanarius</i> sp.	<i>Stenothoe marina</i>	<i>Pseudopolydora kempfi</i>	<i>Prionospio elongata</i> *	<i>Heptaceras quinquedens</i>
<i>Amphipholis</i> sp.	<i>Magelona cincta</i>	<i>Diogenes</i> sp.	<i>Stenothoe</i> sp.	<i>Sigambra parva</i>	<i>Prionospio pulchra</i>	<i>Macroclymene</i> sp.
<i>Armandia intermedia</i>	<i>Melita dentata</i> *	<i>Dorippoides facchino</i>	<i>Sthenelais boa</i>	<i>Sphaeroma walkeri</i> *	<i>Prionospio sexoculata</i>	<i>Pilumnidae</i>
<i>Barnea birmanica</i> *	<i>Melita</i> sp.	Echiura	<i>Xenanthura orientalis</i> *	<i>Sternaspis scutata</i>	<i>Schistomerings rudolphi</i>	<i>Platorchestia platensis</i>
<i>Branchiomma serratibranchis</i> *	<i>Meretrix</i> sp.	<i>Eteone siphodonta</i>		<i>Trochochaeta</i> sp.*		<i>Platylicoa</i> sp.
<i>Cerapus</i> sp.	<i>Microdeutopus</i> sp.	<i>Glycera capitata</i>				<i>Podoceridae</i>
<i>Cheiriphotis megacheles</i> *	<i>Microprotopus maculatus</i>	<i>Glycera longipinnis</i>				<i>Portunidae</i>
<i>Cheiriphotis trifurcata</i> *	<i>Microprotopus</i> sp.	<i>Glycinde kameruniana</i> *				<i>Prosthiostomidae</i>
<i>Cyprinotus</i> sp.	<i>Neanthes glandicincta</i>	<i>Goniada emerita</i>				<i>Synidotea harfordi</i>
<i>Diopatra cuprea</i>	<i>Notopoma crassicornis</i> *	<i>Grandidierella bonnieroides</i>				
<i>Diopatra huperiana monroi</i> *	<i>Photis longicaudata</i>	<i>Harmothoe</i> sp.				
<i>Diopatra monroi</i> *	<i>Photis</i> sp.	<i>Lepidonotus carinulatus</i> *				
<i>Ericthonius brasiliensis</i>	<i>Pista</i> sp.*	<i>Lepidonotus</i> sp.				
<i>Ericthonius</i> sp.	<i>Podocoryna</i> sp.*	<i>Lumbrineris hartmani</i> *				
<i>Eriopispa</i> sp.	<i>Protodorvillea egena</i>	<i>Lumbrineris heteropoda difficilis</i>				
<i>Exciliana</i> sp.	<i>Scoloplos</i> sp.	<i>Lumbrineris heteropoda</i>				
<i>Gammaropsis</i> sp.	<i>Sipuncula</i>	<i>Lumbrineris meteorana</i> *				
<i>Gammarus</i> sp.	<i>Urothoe pulchella</i>	<i>Lumbrineris</i> sp.				
<i>Gari</i> sp.	<i>Urothoe</i> sp.	<i>Mesopodopsis orientalis</i> *				
<i>Grandidierella megnae</i> *	<i>Urothoe viswanathi</i> *	<i>Nematopalaemon tenuipes</i> *				
<b>Ecological group III</b>						
<b>Ecological group IV</b>						
<b>Ecological group V</b>						
<b>Ignored taxa</b>						

\* EG assignation based on genus

✗ EG assignation based on expert judgement

Appendix 3 Ecological Quality Status (EcoQS) derived from five biotic indices and index value for each station of Ulhas estuary during the three seasons (Pr=Premonsoon, M=Monsoon, Po=Postmonsoon)

Samples	AMBI			MAMBI			BENTIX			BOPA			BO2A			
	Pr	M	Po	Pr	M	Po	Pr	M	Po	Pr	M	Po	Pr	M	Po	
U1	2.8	3.6	1.1	0.37	0.36	0.92	2.5	3.4	4.5	0.1845	0.1845	0.0290	0.1845	0.1845	0.0290	High
U2	3.1	4.4	4.3	0.48	0.23	0.41	3.4	2.0	2.0	0.1311	0.2983	0.2553	0.1311	0.2983	0.2553	Good
U3	2.5	4.3	4.1	0.49	0.45	0.80	5.5	2.3	2.9	0.0414	0.2846	0.2093	0.0414	0.2846	0.2093	Moderate
U4	4.0	4.5	5.5	0.39	0.18	0.19	2.4	2.0	2.1	0.1047	0.3010	0.2866	0.1047	0.3010	0.2866	Poor
U5	3.5	4.5	4.1	0.61	0.18	0.70	2.3	2.0	2.9	0.2406	0.3010	0.1256	0.2406	0.3010	0.1256	Bad
U6	5.4	4.5	5.9	0.26	0.18	0.07	2.0	2.0	2.0	0.2711	0.3010	0.3012	0.2711	0.3010	0.3012	High
U7	6.0	4.5	4.3	0.04	0.18	0.59	2.0	2.0	3.5	0.3010	0.3010	0.1761	0.3010	0.3010	0.1761	Good
U8	7.0	4.6	5.8	0.00	0.47	0.09	0.0	2.0	2.0	Azoic	0.2978	0.3010	Azoic	0.3010	0.3010	High
U9	7.0	4.5	6.4	0.00	0.40	0.24	0.0	2.0	2.0	Azoic	0.3010	0.3010	Azoic	0.3010	0.3010	High
U10	6.0	5.1	5.4	0.16	0.31	0.18	2.0	2.0	2.0	0.0116	0.3010	0.3010	0.3010	0.3010	0.3010	Good
U11	4.4	4.0	4.2	0.47	0.45	0.50	2.0	2.0	2.0	0.2518	0.1856	0.2089	0.2604	0.1856	0.2089	Moderate
U12	2.3	3.2	3.3	0.89	0.49	0.58	3.4	2.0	2.0	0.0807	0.0000	0.0934	0.0821	0.0290	0.0934	Good
Euhaline	2.8	3.6	1.1	0.45	0.36	0.92	3.9	3.4	4.4	0.1190	0.1845	0.0290	0.1190	0.1845	0.0290	High
Poly-mesohaline	5.5	4.4	4.7	0.22	0.23	0.46	1.4	2.0	2.6	0.2294	0.2983	0.2257	0.2294	0.2983	0.2257	Good
Oligohaline	4.2	4.4	4.8	0.51	0.33	0.32	2.5	2.0	2.0	0.1147	0.2574	0.2411	0.2145	0.2606	0.2411	Moderate

Azoic zones are considered ‘Bad’ as per Dauvin and Ruellet, 2007

Appendix 4 Ecological Quality Status (EcoQS) derived from five biotic indices and index value for each station of Amba estuary during the three seasons (Pr=Premonsoon, M=Monsoon, Po=Postmonsoon)

Samples	AMBI			MAMBI			BENTIX			BOPA			BO2A			
	Pr	M	Po	Pr	M	Po	Pr	M	Po	Pr	M	Po	Pr	M	Po	
A1	2.7	6.3	3.0	0.85	0.11	0.53	3.8	1.3	3.4	0.1455	0.1614	0.1828	0.1455	0.1614	0.1828	High
A2	2.8	4.2	3.2	0.57	0.38	0.44	3.8	5.8	2.9	0.0864	0.0095	0.1836	0.0864	0.0095	0.1836	Good
A3	3.3	4.1	3.0	0.66	0.20	0.21	3.8	2.0	2.0	0.0226	0.2041	0.0000	0.0226	0.2041	0.0000	Moderate
A4	3.3	4.6	3.8	0.52	0.14	0.21	3.0	2.0	2.6	0.2392	0.2788	0.2688	0.2392	0.3010	0.2688	Poor
A5	2.6	1.5	2.7	0.64	0.55	0.42	3.2	5.3	2.9	0.1740	0.0000	0.1603	0.1740	0.0000	0.1603	Bad
A6	3.4	5.5	4.9	0.54	0.76	0.27	4.6	4.6	2.6	0.0908	0.0191	0.2553	0.0908	0.0191	0.2553	High
A7	2.8	3.9	1.5	0.59	0.41	0.88	4.3	2.8	4.8	0.1082	0.1741	0.0116	0.1270	0.1741	0.0116	Good
A8	3.6	3.6	2.6	0.59	0.48	0.50	3.2	3.1	4.4	0.1773	0.2088	0.0986	0.2028	0.2088	0.1249	Moderate
A9	2.3	3.9	2.1	0.79	0.89	0.49	3.9	2.5	5.1	0.0358	0.1988	0.0642	0.0358	0.1988	0.0642	Poor
A10	2.0	6.4	2.2	0.40	0.23	0.47	5.2	1.0	5.2	0.0821	0.0235	0.0484	0.0821	0.0235	0.0484	Bad
A11	1.8	4.1	3.6	0.48	0.38	0.43	5.0	2.5	2.9	0.0090	0.2304	0.1885	0.0090	0.2304	0.1965	High
A12	4.4	4.5	6.7	0.44	0.20	0.09	5.6	2.0	0.8	0.0314	0.3010	0.0000	0.0314	0.3010	0.0000	Good
A13	4.5	4.8	6.4	0.12	0.35	0.24	2.0	2.0	2.0	0.2993	0.2434	0.2983	0.2993	0.3010	0.2983	Moderate
Euhaline	2.8	5.2	3.4	0.60	0.25	0.35	4.0	4.1	2.7	0.1064	0.0854	0.1751	0.1105	0.0854	0.1751	High
Poly-mesohaline	4.4	3.9	3.1	0.25	0.42	0.48	3.8	4.9	3.9	0.1654	0.1475	0.0685	0.1654	0.1512	0.0743	Good
Oligohaline	4.7	6.4		0.41	0.24		2.2	2.0		0.1994	0.2983		0.2110	0.2983		Moderate

Appendix 5 Ecological Quality Status (EcoQS) derived from five biotic indices and index value for each station of Savitri estuary during the three seasons (Pr=Premonsoon, M=Monsoon, Po=Postmonsoon)

Samples	AMBI			MAMBI			BENTIX			BOPA			BO2A			
	Pr	M	Po	Pr	M	Po	Pr	M	Po	Pr	M	Po	Pr	M	Po	
S1	1.1	3.9	4.4	0.79	0.22	0.41	4.6	2.5	2.1	0.0061	0.2742	0.2918	0.0061	0.2742	0.2918	
S2	1.6	0.0	0.8	0.72	0.43	0.77	4.2	6.0	5.7	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
S3	3.9	1.8	2.4	0.38	0.33	0.72	4.0	5.7	3.9	0.0000	0.0348	0.0908	0.0000	0.0348	0.0908	
S4	1.4	3.8	1.5	0.49	0.68	0.37	6.0	2.7	6.0	0.0000	0.2385	0.0000	0.0000	0.2385	0.0000	
S5	2.4	4.1	3.0	0.72	0.74	0.81	3.7	2.4	3.1	0.0474	0.2128	0.1759	0.0474	0.2128	0.1759	
S6	1.4	4.0	3.6	0.58	0.48	0.75	5.8	2.4	2.3	0.0180	0.2745	0.1784	0.0180	0.2745	0.1784	
S7	1.4	6.0	5.6	0.56	0.05	0.27	3.6	2.0	1.7	0.1657	0.0000	0.0000	0.1800	0.3010	0.2891	
S8	1.5	7.0	5.3	0.33	0.00	0.30	6.0	0.0	3.0	0.0000	0.0000	0.0000	0.3010	0.0000	0.0000	
Euhaline	2.0	2.0	2.6	0.60	0.33	0.59	5.3	4.2	3.9	0.0015	0.1371	0.1459	0.0015	0.1371	0.1459	
Poly-mesohaline	1.8	1.8	3.2	0.62	0.33	0.59	4.9	5.7	3.7	0.0770	0.0348	0.0890	0.0818	0.0348	0.1468	
Oligohaline	1.5	5.0	5.3	0.33	0.38	0.30	6.0	2.4	6.0	0.0000	0.1451	0.0000	0.3010	0.2054	0.0000	