

A comparison of bacterial communities from omz sediments in the arabian sea and the bay of bengal reveals major differences in nitrogen turnover and carbon recycling potential

^{1,2}Jovitha Lincy * and ^{1,2}Cathrine Sumathi Manohar

¹Biological Oceanography Division, CSIR-National Institute of Oceanography (NIO),
Goa-403004, India.

²Academy of Scientific and Innovative Research (AcSIR), Ghaziabad-201002, India.

Correspondence:

JL: jovithalincy@gmail.com; +91-9847871900 (*corresponding author).

CSM: cathrine@nio.org; +91-832-245-0441.

Abstract:

The Northern Indian Ocean hosts two Oxygen Minimum Zones (OMZ), one in the Arabian Sea and the other in the Bay of Bengal. 16S rRNA gene amplicon sequencing was used to understand the total bacterial diversity in, the surface sediment off Goa within the OMZ of the Arabian Sea, and off Paradeep within the OMZ of the Bay of Bengal. Functional profiling was carried out to pinpoint the occurrence of specific bacterial operational taxonomic units (OTUs) which have been previously described to harbor certain genes/enzymes relevant to biogeochemical cycling of carbon, nitrogen, and sulfur compounds. The dominant phyla identified included Firmicutes (33.08%) and Proteobacteria (32.59%) from the Arabian Sea, and Proteobacteria (52.65%) and Planctomycetes (9.36%) from the Bay of Bengal. Only 30% of OTUs were shared between the sites which make up three-fourth of the Bay of Bengal OMZ bacterial community, but only one-fourth of the Arabian Sea OMZ sediment bacterial community. Statistical analysis indicated the bacterial diversity from sediments of the Bay of Bengal OMZ is ~48% higher than the Arabian Sea OMZ. The community analysis combined with a predictive functional profiling of 16S rRNA amplicons revealed some major differences regarding sediment nitrogen fixation and carbon recycling, and identified a distinct bacterial community structure within the two shallow OMZ sites lying in the east coast and west coast of the peninsular India.

Keywords:OMZ, sediment bacteria, 454 pyrosequencing, Arabian Sea, Bay of Bengal, functional ecology.

1. Introduction

The Northern Indian Ocean consists of two major ocean basins: the Arabian Sea (AS) in the west and the Bay of Bengal (BoB) in the east. Even though both these basins are placed in the same latitude, they differ in many aspects. This includes differences in average salinity, primary productivity, nitrogen loss, the intensity of mesoscale eddies, contrasting transport of dissolved oxygen, and organic matter (McCreary Jr et al., 2013). Both these basins experience intense oxygen depletion below the mixed layer of the water column, where dissolved oxygen (DO) is usually below the detection limit of conventional methods. The AS-OMZ between the water depths of ~100/150 – 1000/1200m is the thickest OMZ, and is identified as a primary site of fixed nitrogen loss (Naqvi et al., 2006). In contrast, the BoB-OMZ occurs at shallow depths, more seasonal in nature (Sarma et al., 2013) and has been reported to be less intense than the AS-OMZ (Paulmier, 2009) with DO concentrations still present in the nanomolar range (Bristow et al., 2017). Nitrogen loss has been described as rather insignificant and limited by substrate availability resulting from low organic matter supply by primary production (Bristow et al., 2017; Löscher et al., 2020). The sequence of electron acceptor utilization in such an environment, generally follow the thermodynamic energy yield (Froelich et al., 1979). However, recent studies support the possibility of co-occurrence of metabolisms using different electron acceptors in OMZs, one example would be the existence of a cryptic sulfur cycle, which occurs along with nitrogen cycle processes (Callbeck et al., 2018; Canfield et al., 2010).

Surface sediment underlying OMZs entraps all recent microbial signatures of the water column above (Gerdes et al., 2000) in addition to the sediment microbiome; hence it is interesting to explore and compare such benthic OMZ ecosystems, especially those located in shallow zones. OMZs act as niches for microorganisms that can use alternative pathways of respiration (Diaz & Rosenberg, 2008; Pitcher et al., 2011). In the BoB-OMZ, aerobic communities have been identified to coexist with anaerobic communities (Bristow et al., 2017). In the AS, such coexistence was explained by separate micro-niches in the same environment (Pitcher et al., 2011). Similar studies carried out in the eastern AS-OMZ sediments have identified Proteobacteria (52%) and Planctomycetes (12.7%) as the dominant phyla (Divya et al., 2011). Other integral phyla of soil/sediment habitat are Bacteroidetes, Acidobacteria, Actinobacteria, and Firmicutes (Lv et al., 2014).

It is vital to understand the dominant microbial taxa and also their functional ecology to throw light on the biogeochemistry of these oxygen-depleted zones (Rajpathak et al., 2018). With the advent of molecular techniques over the last decade, a large volume of data has been generated

which helped to elucidate the bacterial community structure (Hodkinson & Grice, 2015). Phylogenetic profiling, using next-generation sequencing (NGS) techniques, offer high-resolution data from complex environments (Claesson et al., 2010). By using algorithms leveraging functional databases, it is also possible to predict putative functional ecology from 16S rRNA amplicon data. The available data on the bacterial community structure of the northern Indian Ocean OMZ using such high throughput sequencing techniques has been limited to the pelagic realm (Fernandes. et al., 2019; Rajpathak et al., 2018), or restricted to some functionally significant groups rather than total bacterial community (Fernandes et al., 2018). Descriptions of OMZ sediment bacterial communities are largely underrepresented and need special attention.

The objective of our work was to compare the surface sediment bacterial taxonomic and functional diversity within two major OMZs in the northern Indian Ocean, the Arabian Sea (AS) and the Bay of Bengal (BoB), using NGS on the v1-v3 hypervariable region of the 16S rRNA gene. Based on this high throughput sequencing dataset, we predicted the metabolic potential present at both sites, the AS and the BoB with a key focus on genes relevant for nitrogen and sulfur turnover, and many fermentative pathways to understand the bacterial role in carbonate precipitation, a possibility least explored.

2. Materials & methods

2.1. Sample collection and site characteristics

Sediment samples were collected in February 2013 off Goa in the AS-OMZ (SSK-046, *RV Sindhu Sankalp*), at the GS1A site located at 15⁰13'N, 72⁰56'E, and in August 2014 off Paradeep in the BoB (SSD-002, *RV Sindhu Sadhana*), at the PS1B site located at 19⁰57'N, 86⁰46'E [Fig.1-A: Plotted in Ocean Data View v.5.2.1.(Schlitzer, 2015)]. A profile of nutrient and DO distribution is presented from the AS and the BoB, respectively (Fig.1-B-C). These profiles represent typical conditions for the two sampling locations but were obtained from other cruises [B: 15⁰1'N, 72⁰26'E, collected in February 2018; and C: 19⁰41'N, 86⁰45'E, collected in August 2010] and show the distribution of DO, nitrate (NO₃⁻), and nitrite (NO₂⁻) from the surface to 1000m water depth in μM. Sampling at both stations covered surface sediments below a ~200m deep water column, underlying OMZ waters. Though both areas experience intense oxygen depletion with the core of the OMZ located between 150 and 500 m, the maximum NO_x values are twice as high in the AS as compared to the BoB, with a prominent secondary nitrite maxima (SNM). A box corer was used to retrieve the sediment samples. The sediment cores were carefully sub-sampled using acrylic core liners (25 mm ID, ~30 cm length), sub-samples were taken from the center of the core to avoid mixing of sediment layers. The 0-5 cm subsections of samples were transferred into sterile screw-cap containers.

Samples were handled sterile and preserved at -20°C until further analysis. The Temperature/Salinity profiling of the water column above the sediment was carried out using a Sea-Bird Electronics (SBE, USA) conductivity-temperature-depth (CTD) sensor (SBE9), equipped with a Niskin bottle rosette sampling system. The dissolved oxygen (DO) sensor used here (SBE43) yields an accuracy: $\pm 2\%$ of saturation, precision: 1 μM , and detection limit: $\leq 1\mu\text{M}$.

2.2. Sediment characterization

The sediment samples were freeze-dried, homogenized, and ground in an agate mortar prior analysis. Total carbon (TC) and total nitrogen (TN) comprising both inorganic and organic forms were analyzed in an elemental carbon/ nitrogen (CN) analyzer (Fisons NA 1500) using the method described in Bhushan et al. (2001). The calibration of the CN analyzer was done using a reference standard (NC-soil), and the obtained recovery rate was 96% for TC and 99% for TN. The precision was monitored by carrying out replicates for both samples and was $\pm 1\%$. The detection limits were two times the blank value. Total organic carbon (TOC) contents were determined with a colorimetric based wet oxidation method (Azam & Sajjad, 2005), which is reported to be highly reproducible. Total inorganic carbon (TIC) was determined as the difference between TC and TOC (Bernard et al., 1995). Organic matter (OM) was calculated by multiplying TOC with the van Bemmelen factor 1.724 (Heaton et al., 2016), based on the assumption that humidified organic matter of soil contains 58% carbon, however, variations of 40-60% have been observed (Nelson & Sommers, 1982). For determining CaCO_3 abundances, TIC was multiplied with a factor of 8.33 to get the percent calcium carbonate as described previously (Bernard et al., 1995). The TOC/TN value was converted into molar ratio by multiplying with a factor 1.167 derived from atomic weights of Carbon and Nitrogen (i.e., 14/12) (Meyers, 1994).

2.3. Genomic DNA extraction and 454 Pyrosequencing

Total genomic DNA was extracted from 400-500 mg of the sediment samples in triplicates, using the Fast DNATM SPIN Kit for Soil (MP Biomedicals, Santa Ana, CA). The purified DNA was quantified using a Nanodrop 2000 spectrophotometer (Thermo Fisher Scientific, USA). DNA was quality checked on an agarose gel (0.8%). The extracted DNA was pooled and amplified using barcoded fusion primers targeting the v1–v3 region of the 16S rRNA gene using the universal primer 27F (GAGTTTGATCMTGGCTCAG) and 518R (ATTACCGCGGCTGCTGG) (Okubo et al., 2009). Mixed amplicons were subjected to emulsion PCR and then deposited on picotiter plates (Agilent, USA). Amplification conditions consisted of an initial denaturation step at 95°C for 5 min, followed by 30 cycles of denaturation at 95°C for 30 sec, annealing at 55°C for 30 sec, and

elongation at 72°C for 30 sec, with a final elongation at 72°C for 5 min. The detailed procedure of pyrosequencing is described elsewhere (Suh et al., 2014). Sequencing was performed by Chunlab Inc. (Seoul, Korea) using a 454 GS FLX Titanium Sequencing system (Roche Branford, CT, USA) per the manufacturer's instructions.

2.4. Sequence data processing

Amplicon pyrosequencing data were processed using the QIIME software package, ver. 1.7. (Caporaso et al., 2010). Chimaeras and primer mismatch sequences were removed from the amplicon dataset using the Amplicon Noise software, version 1.27 (Quince et al., 2011) available from the FLX Titanium sequence data platform, and implemented in QIIME, a using the program CD-HIT (Edgar, 2010). The average read length of PCR amplicons was 378±45 bp. The resulting reads were taxonomically classified based on similarity scores in both the basic local search tool (BLASTN) searches (E-value >10⁻⁵) on the EzTaxon-e 16S rDNA database (2014.07.01) and on the SILVA SSU database, release 132, based on the RDP classifier method (version14) (Im et al., 2012). Relative abundances of taxonomic groups were estimated using the following cut-off values: species ($x \geq 97\%$), genus ($97\% > x \geq 94\%$), family ($94\% > x \geq 90\%$), order ($90\% > x \geq 85\%$), class ($85\% > x \geq 80\%$) and phylum ($80 > x \geq 75\%$). If the similarity was lower than the specific cut-off value, the sequence was characterized unclassified (un) (Chun et al., 2007); sequences that didn't have any cultivable representatives were shortened as 'ucl.' The diversity indices and rarefaction curves were calculated at 97% sequence similarity using the Mothur platform v.1.43.0. (Schloss et al., 2009). The CLCommunityTM software version 3.46 was used for data visualization. Venn diagrams were used to compare sediment bacterial taxonomic composition between sampling sites (Fig.2).

Out of 17,784 reads, 43% were filtered out during the quality processing. Altogether, 5944 reads for the AS sediment, and 4125 reads for the BoB sediment sample were available for further analysis with a mean length of approx. 470-480 bp. In marine sediments, pyrosequencing read numbers varied between 5,000 and 20,000 per sample in previous studies (Choi et al., 2016; Zhu et al., 2013), the output of our sequencing approach is in the same order of magnitude. The taxonomic assignment done using the SILVA platform (Quast et al., 2012) resulted in classifying the bacterial sequences into several specific clades, the EzTaxon-e data analysis (Chun et al., 2007) was used for species-level taxonomic assignment.

2.5. Functional prediction of 16S rRNA amplicons

For the functional prediction of 16S rRNA pyrosequencing amplicons, the OTUs were clustered at 97% sequence similarity. The OTU table and representative sequence fasta files were

submitted to the Piphillin pipeline (<https://piphillin.secondgenome.com/>) (Iwai et al., 2016). The Piphillin algorithm has the advantage to not rely on phylogenetic trees to predict metagenomic contents. It further uses more recent releases of the functional database Kyoto Encyclopedia of Genes and Genomes (KEGG, updated Oct 2018) and BioCyc as compared to alternative pipelines such as PICRUSt or Tax4Fun (Narayan et al., 2020). It utilizes nearest-neighbor matching 16S rRNA amplicons (or genomes) to predict the representative genomes. The normalized 16S rRNA copy number of each genome is inferred using gene content collected in functional databases (Iwai et al., 2016; Langille et al., 2013; Narayan et al., 2020). The KEGG reference database was used at a 90% cutoff level to predict metabolic functions present in the sequenced microbial community. The final output of this workflow was quantified in terms of predicted gene abundances per number of OTUs per sample. The information extracted was based on a small fraction of the population available from the KEGG database. At 90% similarity cut off, around 338 KEGG pathways were identified from 156 OTU representatives from the AS and 354 KEGG pathways for 469 OTUs from the BoB. We focused on the KEGG database pathways for nitrogen (ko00910), sulfur (ko00920), and methane (ko00680) turnover, as well as on carbon metabolism (ko01200) with specific focus on fermentation and bioenergetics pathway related to carbon fixation, as they are reported to be prevalent in the OMZ sediment.

3. Results And Discussion

3.1. Sediment biogeochemistry

In the present study, both sampling sites showed intense oxygen depletion with dissolved oxygen (DO) concentrations of $2 \pm 0.4 \mu\text{M}$, at bottom waters which was $\sim 3\text{m}$ above the sediment sampled. In the shallow zones of the BoB-OMZ and in the AS-OMZ, the DO concentration sometimes falls below the detection limit of conventional methods, especially during the summer monsoon, due to the increased riverine nutrient loading, coastal high primary production and increased respiration (Sarma et al., 2013). Between both sampling sites, bottom water salinity was comparable, but the temperature was 3°C high at the AS sampling site, which may be a seasonal or permanent feature. The sample characteristics of the collected sediment and near-bottom waters are presented in Table-1. In brief, total organic carbon was slightly higher in the AS with 3.47%, and 2.24% in the BoB, TN values were 0.28% and 0.16% in the AS and BoB, respectively. The TOC/TN ratio was 8.37 in the AS, and 9.66 in the BoB, thus conditions in the organic matter pool were rather comparable. Here the values are in the typical range of OMZ sediments and higher than non-OMZ surface sediments where the TOC and TN concentration as low as 0.2 and 0.02 wt. % reported (Pattan et al., 2013). OMZs enhance the preservation of organic matter, explaining the reported

values of TOC ranging from ~1-2 to 6-7% (Cowie et al., 2014) and TOC/TN ratios within 7.3 – 12.3 (van der Weijden et al., 1999). Our data is in line with those OMZ ranges, with somewhat lower concentrations for both, TOC and TN, in sediments of the BoB. This may result from generally assumed lower productivity of the BoB waters, compared to the AS rapid nitrogen burial as described for OMZ sediments (Robinson et al., 2012), or different activities in re-mineralization processes (Bohlen et al., 2011). TIC, which was substantially higher in the AS with 8.11%, compared to the BoB with only 0.29%, while the values falls within the range of OMZ sediments, strikingly higher difference in TIC between the sites could be attributed to the difference in CaCO₃ content caused by increased carbon sequestration (Sarma et al., 2007). Additionally, meiobenthic fauna with shells may contribute to the difference in TIC, as this is found to be abundant in sediments of the AS while not abundant in the BoB (Ramaswamy & Gaye, 2006). Besides, different microbial communities could explain patterns of carbonate precipitation, a possibility which we will explore in the following.

3.2. Benthic bacterial community structure

Between the study sites, the BoB sediments harbor a more diverse bacterial community than the sediments of the AS, which is illustrated not only by the general diversity of taxa and in line with the few available other studies (Dang et al., 2008; Zhu et al., 2013) but also corroborated by various diversity measures as presented in Table-2. Given that our rarefaction analysis (A1) showed that our sequencing approach was able to recover ~70% of bacterial phylotypes from the BoB and 90 % from the AS sediments, the diversity in the BoB is however still rather underestimated and may be even higher due to the recovery of many rare OTUs.

The dominant communities and their relative percentage remained the same for BLASTN searches using the EzTaxon-e 16S database, and pairwise alignment using the SILVA 132 database. This led to a successful in classification of 44 to 48 phyla, 27 of which were common to both sites. Generally, the dominant bacterial phyla consisted of Firmicutes (33.08%), Proteobacteria (32.59%), Bacteroidetes (17.48%), and Chloroflexi (5.52%) in AS sediments and Proteobacteria (52.65%), Planctomycetes (9.36%), Actinobacteria (7.25%), Firmicutes (5.5%), Acidobacteria (6.74%) and Chloroflexi (4.49%) in BoB sediments. Those abundant taxa contributed >85% to the total bacterial community.

The dominance of Proteobacteria is well documented in marine ecosystems (Wang et al., 2012). In the eastern AS-OMZ surface sediment, nearly 14 phyla were identified in a previous study using the Sanger sequencing technique, the majority of which were Proteobacteria (52%), followed

by Planctomycetes (12.7%) and Chloroflexi (8.8%) (Divya et al., 2011). Similarly, in another study carried out utilizing high throughput sequencing confirms Proteobacteria to be the dominant phylum making up 70-75% in all six sites within benthic OMZ of AS followed by Bacteroidetes. Representative sequences affiliated to phyla Chloroflexi and Firmicutes were also recovered in a considerable number (Fernandes et al., 2018). From sediments collected from off Paradeep port, which is roughly 27 nautical miles from our BoB sampling site PS1B, close to 40 bacterial phyla were reported using 454 pyrosequencing approach. The relative contribution of the phylum Proteobacteria was only 17%, which was lesser than Bacteroidetes (23%) and Firmicutes (19%) (Pramanik et al., 2016) indicating a certain patchiness in relative abundance but an overall comparability of the bacterial community composition in the BoB possibly resulting from factors including DO (Stewart et al., 2012), the availability of nutrients or organic carbon (Fierer & Jackson, 2006). The BoB receives surplus river water run-off (McCreary Jr et al., 2013), which might be a reason for the observed differences in diversity, as two stations had very different nutrient chemistry, else comparable depth and oxygen depletion levels.

The candidate phyla GN02, OD1, TM6, TM7, and WS3, were prevalent in ESP (eastern south pacific) pelagic OMZ microbiome as well, implying that they have an essential role in OMZ nutrient cycling (Ganesh et al., 2014; Ulloa et al., 2013). Candidate phyla GN02, OP3, OP8, were unique to both sampled OMZs sediments of the northern Indian Ocean. A total of 13 candidate phyla were obtained in our study. The prevalence of such “bacterial dark matter” highlights the need to decipher their coding potential, as they cannot be subjected to functional predictions due to a lack of cultivable representatives.

A complete list of taxa is presented in the Supplementary information A2 obtained through SILVA portal. A3 and A4 represents Krona chart of the AS and the BoB sediment bacterial community structure obtained through EzTaxon-e portal. Interestingly, only 28.48% of the identified OTUs were shared between the AS and the BoB, on the genus level (similarity were between 64.29% on the phylum level), leaving 53.10% of unique OTUs in the BoB and 18.42% in the AS (Fig. 2). This suggests that the two sediments, while biogeochemically similar, harbor a largely different bacterial community.

The analysis of 58 bacterial classes recovered from our data set showed that there >50% similarities between the phylotypes at the two sites which makes up ~97% of bacteria. The dominant classes in the AS sediment include Bacilli (32.96%), Gammaproteobacteria (18.34%) and Bacteroidia (17.19%). In the BoB sediment, Gamma-, Alpha, and Delta-Proteobacteria (23.68%, 19.01%, 9.26% respectively) were most abundant, followed by Planctomycetacia (6.72%). Those

clades together contribute between 60-70% of the total in the BoB sediment. Dominant bacterial orders recovered exclusively from the AS-OMZ include Bacillales (32.94%), majorly Planococcaceae (26.06%) followed by Flavobacteriales (17.14%), and Oceanospirillales (12.85%). In BoB sediments, Steroidobacterales (7.05%) and Rhizobiales (11.03%) form the most dominant groups. Exploring the taxonomy in more detail, relative abundances for fermenting organisms such as Planococcaceae, Flavobacteriaceae, Bacillaceae, Oceanospirillaceae, Rhodobacteraceae, and Vibrionaceae are strikingly higher in the AS sediment compared to the BoB amongst the abundant clusters (abundant >1%; Fig. 3). Those clades are mainly described as heterotroph degraders, mostly able to ferment (Glöckner et al., 1999; Yakimov et al., 2003). The presence of Alcanivoraceae in the AS sediment, and their absence in the sediment of the BoB, could be an important factor in the precipitation of CaCO₃, because of their metabolic capability to use ammonification and carbonic anhydrase activity to induce rapid calcium carbonate precipitation (Krause et al., 2018). In the BoB, abundant clades consist mostly of Pseudomonadaceae first described in a deep sea sediment from a Japanese trench [clone AB013829, (Yanagibayashi et al., 1999)] and Desulfobacteraceae, both of which are described as denitrifier groups. Desulfobacteraceae often use acetate (Dyksma et al., 2018) but are also known to degrade other organic compounds (Kümmel et al., 2015). Besides those clades, different proteobacterial clades were found, as well as the purple non-sulfur bacteria Rhodobacteraceae and Rhodospirillaceae, the latter of which are able to fix molecular nitrogen (Madigan et al., 1984). The double pie-chart provides an overview of both sequenced bacterial communities at the class and family level (Fig. 4).

In AS sediments, the most abundant bacterial genus was *Paenisporosarcina* sp. (24.06%), followed by *Salegentibacter* sp. (17%), and as per EzTaxon-e database these were further identified as *Paenisporosarcina quisquiliarum* and *Salegentibacter mishustinae*. Those groups were followed by *Amphritea* (9.02%), *Oceanibulbus* (4.27%), *Alcanivorax* (3.82%), *Photobacterium* (2.76%) and *Salipaludibacillus* (2.61%). All of these clades are unique to the AS sediment and not present in the BoB sediments. In the BoB sediments, the most abundant taxa are *Woeseia* (6.98%) and *Gammaproteobacteria_ucl* (6.5%), with the remaining groups being represented with less than 3%.

The clade *Woeseiaceae/JTB255* is recognized as the most abundant clade in marine sediment, having a cosmopolitan distribution. Moreover, analyzed metagenomes of *JTB255* are known to encode the truncated denitrification pathway to nitrous oxide (Mußmann et al., 2017). Since denitrification mediated nitrogen loss is reported to be dominant in Arabian Sea OMZ, we expected to get more hits in the analyzed amplicon dataset (Ward et al., 2009). Though their

occurrence was not detected, few representative sequences of JTB31 and JTB38 were identified that might play a similar role.

3.3. Predicted functional ecology

For a large proportion of the amplicons, functions could not be assigned clearly, which leads to a rather conservative, and qualitative instead of quantitative estimate of the metabolic potential present at the two sites. The predictive functional profiling of 16S rRNA sequences has identified a high proportion of genes involved in methane cycling, as generally typical for sediments underlying OMZ waters (Bertics et al., 2013; Fulweiler et al., 2007; Gier et al., 2016), followed by genes involved in sulfur and nitrogen cycling. (Fig. 5). Methane turnover rates are rather high in anoxic shelf sediments and were reported to be correlated with the availability of labile organic matter, and concurrent with sulfate reduction (Maltby et al., 2016), explaining the predicted abundance of genes involved in methane turnover in our samples. Despite the difference in diversity between the two sampling sites, almost all predicted gene functions were identical suggesting an overall similar metabolic potential in the two different sediments.

Nitrogen cycle: In northern Indian Ocean OMZs, nitrogen cycling is reported to be very active (Naqvi et al., 2006). At both our sampling sites, genes coding for major nitrogen cycle pathways including nitrogen fixation, dissimilatory nitrate reduction to ammonia (DNRA), nitrification, and denitrification were predicted. Interestingly, anammox genes were not predicted for either site despite the presence of planctomycetes in our dataset (Fig.6). This may be either due to the low number of species-level identifiable OTUs or to a true absence of anammox-capable planctomycetes as consistent with OMZ sediments from the seasonally anoxic Eckernförde Bay in the Baltic Sea and sediments underlying the Peruvian OMZ (Bertics et al., 2013; Dale et al., 2011). While generally planctomycetes were in the sequence pool, hits corresponding to anammox planctomycetes were indeed exceptionally low at our sampling sites, accounting for 0.03 and 0.3% in the AS and BoB, respectively. A sequencing related bias could, however, have led to an underestimation of *Scalindua*-anammox bacteria as a systematic underrepresentation by sequencing of 16S rDNA v1-v3 regions has been reported (Penton et al., 2006). Specifically, the functional marker gene coding for the hydrazine oxidoreductase was not predicted from our 16S rDNA data. This suggests that the contribution of anammox to the nitrogen cycle in the Indian Ocean sediments, at least at our sampling sites, is rather low similar to the pelagic OMZ of the AS where denitrification is reported to be dominant over anammox (Ward et al., 2009), and the BoB where anammox as well as denitrification could not be detected (Bristow et al., 2017). As Planctomycetales are known to encode a large number of sulfatase genes, which makes them as a

specialist for the initial breakdown of sulfated hetero-polysaccharides (Wegner et al., 2013), their role in the Indian Ocean sediments could rather be carbon capture in the sediments (Arango et al., 2007; Dale et al., 2011; Jensen et al., 2011; Shao et al., 2010). Here, the predicted gene abundance was 848 and 2901 for AS and BoB microbiome, the predominant form being arylsulfatase, respectively, contributing 65-85% of the sulfatase pool.

The global annual denitrification rate in sediment would be approximately 200 Tg N, and the majority contributed from sediments underlying OMZ, where its reported two to four times higher (Devol, 2015). Therefore, nitrogen loss processes would be expected to take place in both, sediments of the AS and the BoB. Denitrification and sulfite reductase genes were prevalent in our prediction possibly favoring sulfur driven autotrophic denitrification (Shao et al., 2010), and as previous studies suggested heterotrophic denitrification (Arango et al., 2007). Other denitrifiers recovered from our sequence dataset are Oceanospirillales, Chromatiales, Nitrospirales, Syntrophobacteriales, and NB1-j which are known to encode denitrification genes including *nirS*, *norB* and *nosZ* (de Voogd et al., 2015), and contributed 14.05% in the AS sediment, and 4.46% in the BoB sediment, respectively. Similarly, Flavobacteriales are known denitrifiers (Horn et al., 2005), and was are abundant in the AS with 17.14% of all 16S rDNA sequences.

Recent studies have also linked methane oxidation to nitrite-based denitrification in the Candidatus phylum NC10 (Padilla et al., 2016). This was supported by studies carried out in a freshwater reservoir, where methane stimulated massive nitrogen loss (Naqvi et al., 2018). The Steroidobacter clades which make up 7% of the BoB-OMZ bacterial community are also known to perform denitrification coupled with methane oxidation (Liu et al., 2014). Though, our predictive functional profiling confirms denitrification is dominated over anammox in the northern Indian Ocean OMZ as reported (Sarkar et al., 2020; Ward et al., 2009), a clear link with methane oxidation cannot be established, since the BoB-OMZ is not an active denitrification site. Further, isotopic studies carried out by Shirodkar et al. (2018) in the seasonally anoxic Shelf waters of the Arabian Sea did not show any effect of additional CH₄ amendments on N₂ production through denitrification.

DNRA was predicted as a potential remineralization pathway in both basins of our study. In seasonally hypoxic Baltic Sea sediments, DNRA accounted for almost 75% of benthic nitrogen flux (Dale et al., 2011). In contrast to the other nitrogen cycle genes, the *nifHDK* operon coding for the functional unit of the key gene for nitrogen fixation, the nitrogenase, was predicted in higher abundance in the BoB compared to the AS, with BoB-nif being five times as many as AS-nif. This is consistent with the higher proportions of known sedimentary nitrogen fixers, such as Desulfobacteraceae and Rhodospirillaceae. The presence of nitrogen fixers in sediments underlying

OMZs has been documented for several regions, including the upwelling system off Mauretania, the Baltic Sea and the eastern tropical South Pacific shelves, and nitrogen fixing microbes have been shown to be active although at low rates (Bertics et al., 2013; Gier et al., 2017; Gier et al., 2016).

Sulfur cycling: For the sulfur cycle in both, the AS and the BoB, genes for the assimilatory pathway of sulfate reduction were predicted, as well as sulfur oxidation genes of the *sox* operon (Fig.7) in line with a previous study which identified diverse sulfur reducing bacterial and archaeal OTUs in the AS (Fernandes et al., 2018). In our AS dataset, a potential player in the sulfur cycle could be *Sulfitobacter dubius*, which was represented with 4.32% of all OTUs. All known species of the genus *Sulfitobacter* were isolated from marine habitats and are known to perform sulfite oxidation (Long et al., 2011; Sorokin, 1995). *Thermodesulfovibrio* (phylum Nitrospira) accounted for ~1% of sequences at both sites are known sulfate reducers and have been identified from the eastern tropical South Pacific OMZ, before (Schunck et al., 2013). Sequences corresponding to sulfur reducers like Desulfobacterales (AS: 0.87%, BoB: 2.57%) and Syntrophobacterales (AS: 0.67%, BoB: 1.21%) were also recovered from our dataset and were shown to be abundant in sediments of the Black Sea sulfate-methane transition zone as well as in the Arabian Sea OMZ in both pelagic and benthic realms (Fernandes et al., 2018; Fuchs et al., 2005; Leloup et al., 2007).

Carbon fixation: In the BoB sediment, around 1.75% of gene families were predicted to perform photosynthesis, and major contributors would possibly be Chromatiales (0.07%), Rhodospirillales (0.03%), and members of phylum Cyanobacteria (1.65%). Chromatiales, a group of purple sulfur bacteria, can perform anoxygenic photosynthesis (Manske et al., 2005). Similarly, Rhodospirillales primarily chemoorganotroph and photoheterotroph (Luo & Moran, 2015), can also perform anoxygenic photosynthesis (Manske et al., 2005). It is interesting to note that around 68 Cyanobacterial sequences belonging to order Chroococcales of genus *Prochlorococcus* were retrieved from BoB sediment, where water column depth was ~245m, but only one representative from the AS sediment, which was located at ~200m depth. The observed Chroococcales, are assumed to be a low-light adapted group (West et al., 2001) justifying their occurrence in deeper sites, where light penetration is negligible. Since these groups are also known to fix nitrogen (Van Goethem et al., 2017), will justify the observed difference in *nifH* gene abundance. It may also be that this is sunk out debris, with some preserved DNA. The key enzymes responsible for energy metabolism are presented in Table-3. In particular, the higher predicted abundance of dehydrogenase enzymes responsible for oxidation of organic matter in the AS points towards a difference in carbon metabolism in the two regions. This is most likely due to the increased availability of carbon in the AS, however, as proposed earlier (Orsi et al., 2017), more efficient organic carbon recycling in the

AS may over geological timescales contribute to developing a stronger and more persistent functional anoxia.

Carbon remineralization: As indicated by the high TIC concentration in the AS sediment, carbon remineralization was very active due to the increased availability of organic carbon (Yu et al., 2018). About 20% of the identified bacteria were common soil/sediment inhabitants with a prime role in remineralization of diverse organic carbon compounds (Schimel & Schaeffer, 2015). These include Acidobacteria, Actinobacteria, Bacteroidetes, and Gemmatimonadetes (Janssen, 2006). Similarly, Anaerolineales (phylum Chloroflexi), which contributed 2-3% of the total hits in our dataset, have also been identified with a similar role and were specific to areas that show incredibly low or zero oxygen. In addition, genes responsible for N-glycan degradation (ko00511) were predicted to occur almost twelve times more often in the AS than in the BoB sample. These genes play a role in cell adhesion and sequestration (Varki & Gagneux, 2017). The relative distribution of key enzymes and genes specific to gram-positive and gram-negative bacterial fermenters were compared based on previous reports (Eschbach et al., 2004; Ramos et al., 2000) (Fig.8). Their predicted abundance was higher in AS sediment than in BoB sediments. The connected more complete carbon remineralization which could add an explanation to why the AS-OMZ is more anoxic than the BoB-OMZ as previously suggested(Orsi et al., 2017).

4. Conclusion

We compared bacterial communities from two sites in the northern Indian Ocean OMZ, in the BoB off Paradeep and a site off Goa in the AS. Less than one-third of the phylotypes were shared between the two sites, leaving a large individual proportion of the bacteria for each site. A higher diversity has been identified from the BoB, compared to the AS, however, our functional prediction identified high abundances of typical heterotrophic degraders in the AS, that were only represented in low proportions or absent in the BoB. We further identified denitrifiers, DNRA bacteria and sulfur cycle bacteria at both sites and predicted the presence of their functional genes. The higher functional diversity for organic matter degradation with fermentation in addition to denitrification and sulfur-compound dependent remineralization may explain, why the AS OMZ is generally more anoxic. Here, the variability in carbon respiration pathways may allow for a more efficient or complete respiration along the electron tower, thus consuming more oxidized compounds. The abundance of *Alcanivorax*-like bacteria in AS sediments may provide an explanation for high CaCO₃ precipitation, as this organism has been described to perform this process rapidly when organic nitrogen is available as it is at our sampling site in the AS. A notable finding was the absence of anammox bacteria at both sites. Notably, we predicted nitrogen fixation genes from BoB sediments

but not from AS sediments, possibly resulting from higher nitrogen inputs from the water column in the AS.

Despite the limitation of this study with regard to our sample number, we could contribute a first assessment of bacterial diversity and functionality in coastal sediments of the two Indian Ocean basins, as such, we hope to contribute to the general understanding of how these basins function and why they are so different in their biogeochemistry.

Data Availability

All pyrosequencing reads were submitted to the NCBI GenBank database under accession number KU821783 - KU831324 and MG860544 - MG860851. The supporting information is available as supplementary information.

Appendices

A1: Rarefaction curve of bacterial OTUs (operational taxonomic units) associated with sediments underlying oxygen-depleted waters in the northern Indian Ocean OMZ.

A2: Taxonomic composition of bacterial OTUs analyzed through the SILVA database.

A3: Krona chart showing bacterial community structure of AS-OMZ analyzed through EzTaxon-e database.

A4: Krona chart showing bacterial community structure of BoB-OMZ analyzed through EzTaxon-e database.

Author Contribution

JL prepared the manuscript and performed the experiments and bioinformatics analysis. CSM supervised the research.

Competing Interests

The authors declare that they have no conflict of interest.

Funding

The first author is grateful to the Council of Scientific and Industrial Research (CSIR), India, for fellowship grant 31/026(0245)/2012-EMR-I for doing a Ph.D. This work is supported by CSIR grant: PSC0108. NIO's contribution no: xxxx.

Acknowledgment

We thank the Director of CSIR-NIO for providing the facilities. We acknowledge Chun Lab, Seoul, South Korea, for carrying out pyrosequencing. Special thanks to Dr. Carolin Löscher for the critical comments and valuable suggestions which have significantly helped in improvising the content. We express our gratitude to Dr. Amal Jayakumar, Dr. N. Ramaiah and Dr. Peter Burkill for guidance in manuscript preparation. We acknowledge the contributions of anonymous reviewers of this journal. We also thank crew members of SSK-046 and SSD-002, especially Ms. Larissa Menezes, for GS1A sample collection.

Legends

Table-1: Sediment and bottom water characteristics for the samples collected from the northern Indian Ocean OMZ.

Table-2: Summary of pyrosequencing results and statistical analysis of bacterial sequences retrieved from the northern Indian Ocean OMZ surface sediment samples.

*OTUs (operational taxonomic unit) were calculated using Mothur (3% distance).

§Good's coverage is proportional to non-singleton phylotypes.

Table-3: Distribution of key enzymes relevant in energy metabolism in the northern Indian Ocean surface sediments predicted from 16S rRNA genes.

Fig. 1: A) Station map, blue dots indicate sampling stations, red dots are the regions where OMZ water-column characteristics were obtained from. B) Representative water column profiles of biogeochemical parameters from the AS-OMZ (Unpublished data, personal communication from Dr. G.V.M. Gupa) and C) the BoB- OMZ (Sarma et al., 2013).

Fig. 2: Venn diagram showing OTU number wise comparison of the phylotypes at different taxonomic level assigned through the SILVA database.

Fig. 3: Dominant bacterial taxa retrieved at 1% cut-off based on pairwise alignment in the SILVA SSU database release 132.

Fig. 4: Double Pie chart showing bacterial community composition at the class and family level from the sampling locations based on the EzTaxon-e database.

Fig. 5: Relative distribution of redox metabolic KEGG pathways identified from our 16S rRNA amplicon pyrosequencing dataset utilizing the Piphillin algorithm.

Fig. 6: Proposed pathway for OMZ Nitrogen cycling in sediments of northern Indian Ocean OMZ, and the abundance of enzymes with coding genes are indicated in box, where 'blue' and 'red' denotes AS and BoB gene count. Expansion of abbreviation are as follows:- **amoA**: Ammonia monooxygenase subunit A; **gdh**: Glutamate dehydrogenase; **hao**: Hydroxylamine oxidoreductase; **hzo**: Hydrazine oxidoreductase; **hzs**: Hydrazine synthase; **napA**: Nitrate reductase (cytochrome); **nasA**: Nitrate reductase; **narG**: Nitrate reductase, alpha subunit; **nifH**: Nitrogenase iron protein; **nirK/nirS**: Nitrite reductase subunit K/S; **norB/norC**: Nitric oxide reductase subunit B/C; **nrfA**: nitrite reductase (cytochrome c-552); **nosZ**: Nitrous oxide reductase; **nxrA**: nitrite oxidoreductase, alpha subunit; **ureC**: Urease subunit alpha.

Fig. 7: Proposed pathway for OMZ Sulfur cycling in sediments of northern Indian Ocean OMZ, and the abundance of enzymes with coding genes are indicated in box, where 'blue' and 'red' denotes AS and BoB gene count. Expansion of abbreviation are as follows:- **aprA**: Adenylylsulfate reductase, subunit A; **dsrA/dsrB**: Dissimilatory sulfite reductase subunit-alpha/beta; **dsrC**: Dissimilatory sulfite reductase related protein; **cysC**: Adenylylsulfate kinase; **cysH**: Phosphoadenosine phosphosulfate reductase; **psrA**: Thiosulfite reductase; **rDsr**: Reverse dissimilatory sulfite reductase; **sat**: Sulfate adenylyltransferase; **soeA**: Sulfite:quinone oxidoreductase; **sir**: Sulfite reductase (ferredoxin); **soxB**: S-sulfosulfanyl-L-cysteine sulfohydrolase; **soxC**: Sulfane dehydrogenase; **soxD**: S-disulfanyl-L-cysteine oxidoreductase; **soxX/A**: L-cysteine S-thiosulfotransferase; **soxY/Z**: Sulfur-oxidizing protein; **sseA**: Thiosulfate/3-mercaptopyruvate sulfurtransferase; **sqr**: Sulfide:quinone oxidoreductase; .

Fig. 8: Percentage distribution of key enzymes with coding genes identified in bacterial fermentation (Eschbach et al., 2004; Ramos et al., 2000).

Tables

Table-1

Sampling Details			Sediment Characteristics						Near-bottom water profile (CTD)		
Station code	Date	Sampling depth (m)	TOC	TIC	TN	CaCO ₃	OM	TOC/TN	DO	Temp	Salinity
			%						μM	°C	PSU
GS1A	Feb-2013	197.3	2.01	8.11	0.28	67.56	3.47	8.37	2.31	15.58	35.35
PS1B	Aug-2014	234.4	1.30	0.29	0.16	2.41	2.24	9.66	1.67	12.33	35.019

Table-2

Sample name	Optimized reads	OTU richness*				OTU diversity*		Good's coverage§
		Observed	Chao1	ACE	Jackknife	Shannon	Simpson	
GS1A	5,944	955	2,506	4,305	3,450	4.37	0.934	89.3%
PS1B	4,125	1,889	4,447	7,616	6,242	6.97	0.998	69.5%

Table-3

ENERGY METABOLISM	KEGG	ENZYMES	GS1A	PS1B
Kreb cycle, NADH+H ⁺	K00161	pyruvate dehydrogenase E1 component alpha subunit [EC:1.2.4.1]	836	454
Kreb cycle, NADH+H ⁺	K00627	pyruvate dehydrogenase E2 component (dihydrolipoamide acetyltransferase) [EC:2.3.1.12]	1279	1108
Kreb cycle, NADH+H ⁺	K00382	dihydrolipoamide dehydrogenase [EC:1.8.1.4]	2476	1889
Kreb cycle, NADH+H ⁺	K00031	isocitrate dehydrogenase [EC:1.1.1.42]	993	1345
Kreb cycle, NADH+H ⁺	K00164	2-oxoglutarate dehydrogenase E1 component [EC:1.2.4.2]	895	850
Kreb cycle, GTP	K01902	succinyl-CoA synthetase alpha subunit [EC:6.2.1.5]	1064	952
Kreb cycle, FADH ₂	K00239	succinate dehydrogenase / fumarate reductase, flavoprotein subunit [EC:1.3.5.1 1.3.5.4]	902	906
Kreb cycle, NADH+H ⁺	K00024	malate dehydrogenase [EC:1.1.1.37]	1051	881
Glycolysis, ATP	K00927	phosphoglycerate kinase [EC:2.7.2.3]	898	953
Glycolysis, ATP	K00873	pyruvate kinase [EC:2.7.1.40]	955	1012
Glycolysis, NADH+H ⁺	K00134	glyceraldehyde 3-phosphate dehydrogenase [EC:1.2.1.12]	2125	1176
Glyoxylate cycle, NADH+H ⁺	K00024	malate dehydrogenase [EC:1.1.1.37]	1051	881
Glyoxylate cycle, FADH ₂	K00240	succinate dehydrogenase / fumarate reductase, iron-sulfur subunit [EC:1.3.5.1 1.3.5.4]	916	926
Pentose phosphate pathway, NADPH	K00036	glucose-6-phosphate 1-dehydrogenase [EC:1.1.1.49 1.1.1.363]	678	856
Pentose phosphate pathway, NADPH	K00033	6-phosphogluconate dehydrogenase [EC:1.1.1.44 1.1.1.343]	823	807
Ethylmalonyl pathway, NADP ⁺	K14446	crotonyl-CoA carboxylase/reductase [EC:1.3.1.85]	230	276
Ethylmalonyl pathway, GTPase activity	K01847	methylmalonyl-CoA mutase [EC:5.4.99.2]	1386	669
Ethylmalonyl pathway, NADP ⁺	K00023	acetoacetyl-CoA reductase [EC:1.1.1.36]	309	345
Ethylmalonyl pathway, ADP +Pi	K01965	propionyl-CoA carboxylase alpha chain [EC:6.4.1.3]	575	329
Ethylmalonyl pathway, ADP +Pi	K01966	propionyl-CoA carboxylase beta chain [EC:6.4.1.3 2.1.3.15]	746	351
Malonate semialdehyde pathway, ADP+Pi	K01961	acetyl-CoA carboxylase, biotin carboxylase subunit [EC:6.4.1.2 6.3.4.14]	1212	938
Propanoyl-CoA metabolism, ADP+Pi	K01965	propionyl-CoA carboxylase alpha chain [EC:6.4.1.3]	575	329

Figures




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8

References:

- Arango, C. P., Tank, J. L., Schaller, J. L., Royer, T. V., Bernot, M. J., & David, M. B. (2007). Benthic organic carbon influences denitrification in streams with high nitrate concentration. *Freshw. Biol.*, 52(7), 1210-1222. <https://doi.org/10.1111/j.1365-2427.2007.01758.x>
- Azam, F., & Sajjad, M. (2005). Colorimetric determination of organic carbon in soil by dichromate digestion in a microwave oven. *Pak. J. Biol. Sci.*, 8(4), 596-598. <https://doi.org/10.3923/pjbs.2005.596.598>
- Bernard, B. B., Bernard, H., & Brooks, J. M. (1995). Determination of total carbon, total organic carbon and inorganic carbon in sediments *TDI-Brooks International/B&B Laboratories Inc. College Station* (pp. 1-5).
- Bertics, V. J., Löscher, C. R., Salonen, I., Dale, A. W., Gier, J., Schmitz, R. A., & Treude, T. (2013). Occurrence of benthic microbial nitrogen fixation coupled to sulfate reduction in the seasonally hypoxic Eckernförde Bay, Baltic Sea. *Biogeosciences*, 10, 1243-1258. <https://doi.org/10.5194/bg-10-1243-2013>
- Bhushan, R., Dutta, K., & Somayajulu, B. (2001). Concentrations and burial fluxes of organic and inorganic carbon on the eastern margins of the Arabian Sea. *Mar. Geol.*, 178(1), 95-113. [https://doi.org/10.1016/S0025-3227\(01\)00179-7](https://doi.org/10.1016/S0025-3227(01)00179-7)
- Bohlen, L., Dale, A. W., Sommer, S., Mosch, T., Hensen, C., Noffke, A., Scholz, F., & Wallmann, K. (2011). Benthic nitrogen cycling traversing the Peruvian oxygen minimum zone. *Geochim. Cosmochim. Acta.*, 75(20), 6094-6111. <https://doi.org/10.1016/j.gca.2011.08.010>
- Bristow, L. A., Callbeck, C. M., Larsen, M., Altabet, M. A., Dekaezemacker, J., Forth, M., Gauns, M., Glud, R. N., Kuypers, M. M., & Lavik, G. (2017). N₂ production rates limited by nitrite availability in the Bay of Bengal oxygen minimum zone. *Nat. Geosci.*, 10(1), 24-29. <https://doi.org/10.1002/lom3.10126>
- Callbeck, C. M., Lavik, G., Ferdelman, T. G., Fuchs, B., Gruber-Vodicka, H. R., Hach, P. F., Littmann, S., Schoffelen, N. J., Kalvelage, T., & Thomsen, S. (2018). Oxygen minimum zone cryptic sulfur cycling sustained by offshore transport of key sulfur oxidizing bacteria. *Nat. Commun.*, 9(1), 1729-1740. <https://doi.org/10.1038/s41467-018-04041-x>
- Canfield, D. E., Stewart, F. J., Thamdrup, B., De Brabandere, L., Dalsgaard, T., Delong, E. F., Revsbech, N. P., & Ulloa, O. (2010). A cryptic sulfur cycle in oxygen-minimum-zone waters off the Chilean coast. *science*, 330(6009), 1375-1378. <https://doi.org/10.1126/science.1196889>
- Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N., Peña, A. G., Goodrich, J. K., & Gordon, J. I. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods*, 7(5), 335-336. <https://doi.org/10.1038/nmeth.f.303>
- Choi, H., Koh, H.-W., Kim, H., Chae, J.-C., & Park, S.-J. (2016). Microbial community composition in the marine sediments of Jeju Island: next-generation sequencing surveys. *J. Microbiol. Biotechnol.*, 26(5), 883-890. <https://doi.org/10.4014/jmb.1512.12036>

- Chun, J., Lee, J.-H., Jung, Y., Kim, M., Kim, S., Kim, B. K., & Lim, Y.-W. (2007). EzTaxon: a web-based tool for the identification of prokaryotes based on 16S ribosomal RNA gene sequences. *Int. J. Syst. Evol. Microbiol.*, *57*(10), 2259-2261. <https://doi.org/10.1099/ijs.0.64915-0>
- Claesson, M. J., Wang, Q., O'Sullivan, O., Greene-Diniz, R., Cole, J. R., Ross, R. P., & O'Toole, P. W. (2010). Comparison of two next-generation sequencing technologies for resolving highly complex microbiota composition using tandem variable 16S rRNA gene regions. *Nucleic Acids Res.*, *38*(22), e200. <https://doi.org/10.1093/nar/gkq873>
- Cowie, G., Mowbray, S., Kurian, S., Sarkar, A., White, C., Anderson, A., Vergnaud, B., Johnstone, G., Brear, S., & Woulds, C. (2014). Comparative organic geochemistry of Indian margin (Arabian Sea) sediments: estuary to continental slope. *Biogeosciences*, *11*(23), 6683-6696. <https://doi.org/10.5194/bg-11-6683-2014>
- Dale, A. W., Sommer, S., Bohlen, L., Treude, T., Bertics, V. J., Bange, H. W., Pfannkuche, O., Schorp, T., Mattsdotter, M., & Wallmann, K. (2011). Rates and regulation of nitrogen cycling in seasonally hypoxic sediments during winter (Boknis Eck, SW Baltic Sea): Sensitivity to environmental variables. *Estuar. Coast. Shelf Sci.*, *95*(1), 14-28. <https://doi.org/10.1016/j.ecss.2011.05.016>
- Dang, H., Zhang, X., Sun, J., Li, T., Zhang, Z., & Yang, G. (2008). Diversity and spatial distribution of sediment ammonia-oxidizing crenarchaeota in response to estuarine and environmental gradients in the Changjiang Estuary and East China Sea. *Microbiology*, *154*(7), 2084-2095. <https://doi.org/10.1099/mic.0.2007/013581-0>
- de Voogd, N. J., Cleary, D. F., Polónia, A. R., & Gomes, N. C. (2015). Bacterial community composition and predicted functional ecology of sponges, sediment and seawater from the thousand islands reef complex, West Java, Indonesia. *FEMS Microbiol. Ecol.*, *91*(4), fiv019. <https://doi.org/10.1093/femsec/fiv019>
- Devol, A. H. (2015). Denitrification, anammox, and N₂ production in marine sediments. *Annu. Rev. Mar. Sci.*, *7*, 403-423. <https://doi.org/10.1146/annurev-marine-010213-135040>
- Diaz, R. J., & Rosenberg, R. (2008). Spreading dead zones and consequences for marine ecosystems. *science*, *321*(5891), 926-929. <https://doi.org/10.1126/science.1156401>
- Divya, B., Parvathi, A., Bharathi, P. L., & Nair, S. (2011). 16S rRNA-based bacterial diversity in the organic-rich sediments underlying oxygen-deficient waters of the eastern Arabian Sea. *World J. Microbiol. Biotechnol.*, *27*(12), 2821-2833. <https://doi.org/10.1007/s11274-011-0760-0>
- Dyksma, S., Lenk, S., Sawicka, J. E., & Mußmann, M. (2018). Uncultured Gammaproteobacteria and Desulfobacteraceae Account for Major Acetate Assimilation in a Coastal Marine Sediment. *Front. Microbiol.*, *9*, 3124-3124. <https://doi.org/10.3389/fmicb.2018.03124>
- Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*, *26*(19), 2460-2461. <https://doi.org/10.1093/bioinformatics/btq461>
- Eschbach, M., Schreiber, K., Trunk, K., Buer, J., Jahn, D., & Schobert, M. (2004). Long-term anaerobic survival of the opportunistic pathogen *Pseudomonas aeruginosa* via pyruvate fermentation. *J. Bacteriol.*, *186*(14), 4596-4604. <https://doi.org/10.1128/JB.186.14.4596-4604.2004>

- Fernandes, S., Mazumdar, A., Bhattacharya, S., Peketi, A., Mapder, T., Roy, R., Carvalho, M. A., Roy, C., Mahalakshmi, P., & Da Silva, R. (2018). Enhanced carbon-sulfur cycling in the sediments of Arabian Sea oxygen minimum zone center. *Sci. Rep.*, 8(1), 8665-8680. <https://doi.org/10.1038/s41598-018-27002-2>
- Fernandes., G. L., Shenoy, B. D., Menezes, L. D., Meena, R. M., & Damare, S. R. (2019). Prokaryotic Diversity in Oxygen Depleted Waters of the Bay of Bengal Inferred Using Culture-Dependent and-Independent Methods. *Indian J. Microbiol.*, 59(2), 193-199. <https://doi.org/10.1007/s12088-019-00786-1>
- Fierer, N., & Jackson, R. B. (2006). The diversity and biogeography of soil bacterial communities. *Proc. Natl. Acad. Sci. U. S. A.*, 103(3), 626-631. <https://doi.org/10.1016/j.apsoil.2019.06.008>
- Froelich, P. N., Klinkhammer, G., Bender, M. a. a., Luedtke, N., Heath, G. R., Cullen, D., Dauphin, P., Hammond, D., Hartman, B., & Maynard, V. (1979). Early oxidation of organic matter in pelagic sediments of the eastern equatorial Atlantic: suboxic diagenesis. *Geochim. Cosmochim. Acta*, 43(7), 1075-1090. [https://doi.org/10.1016/0016-7037\(79\)90095-4](https://doi.org/10.1016/0016-7037(79)90095-4)
- Fuchs, B. M., Woebken, D., Zubkov, M. V., Burkill, P., & Amann, R. (2005). Molecular identification of picoplankton populations in contrasting waters of the Arabian Sea. *Aquat. Microb. Ecol.*, 39(2), 145-157. <https://doi.org/10.3354/ame039145>
- Fulweiler, R. W., Nixon, S. W., Buckley, B. A., & Granger, S. L. (2007). Reversal of the net dinitrogen gas flux in coastal marine sediments. *Nature*, 448, 180-182. <https://doi.org/10.1038/nature05963>
- Ganesh, S., Parris, D. J., DeLong, E. F., & Stewart, F. J. (2014). Metagenomic analysis of size-fractionated picoplankton in a marine oxygen minimum zone. *ISME J.*, 8(1), 187-211. <https://doi.org/10.1038/ismej.2013.144>
- Gerdes, G., Klenke, T., & Noffke, N. (2000). Microbial signatures in peritidal siliciclastic sediments: a catalogue. *Sedimentology*, 47(2), 279-308. <https://doi.org/10.1046/j.1365-3091.2000.00284.x>
- Gier, J., Löscher, C. R., Dale, A. W., Sommer, S., Lomnitz, U., & Treude, T. (2017). Benthic Dinitrogen Fixation Traversing the Oxygen Minimum Zone Off Mauritania (NW Africa) [Original Research]. *Front. Mar. Sci.*, 4(390), Article 390. <https://doi.org/10.3389/fmars.2017.00390>
- Gier, J., Sommer, S., Löscher, C. R., Dale, A. W., Schmitz, R. A., & Treude, T. (2016). Nitrogen fixation in sediments along a depth transect through the Peruvian oxygen minimum zone. *Biogeosciences*, 13(14), 4065-4080. <https://doi.org/10.5194/bg-13-4065-2016>
- Glöckner, F. O., Fuchs, B. M., & Amann, R. (1999). Bacterioplankton compositions of lakes and oceans: a first comparison based on fluorescence in situ hybridization. *Environ Microbiol.*, 65, 3721-3726. <https://doi.org/10.1128/aem.65.8.3721-3726>
- Heaton, L., Fullen, M. A., & Bhattacharyya, R. (2016). Critical analysis of the van Bemmelen conversion factor used to convert soil organic matter data to soil organic carbon data: Comparative analyses in a UK loamy sand soil. *Espaço Aberto*, 6(1), 35-44. <https://doi.org/10.36403/espacoaberto.2016.5244>

- Hodkinson, B. P., & Grice, E. A. (2015). Next-generation sequencing: a review of technologies and tools for wound microbiome research. *Adv. Wound Care*, 4(1), 50-58. <https://doi.org/10.1089/wound.2014.0542>
- Horn, M. A., Ihssen, J., Matthies, C., Schramm, A., Acker, G., & Drake, H. L. (2005). *Dechloromonas denitrificans* sp. nov., *Flavobacterium denitrificans* sp. nov., *Paenibacillus anaericanus* sp. nov. and *Paenibacillus terrae* strain MH72, N₂O-producing bacteria isolated from the gut of the earthworm *Aporrectodea caliginosa*. *Int. J. Syst. Evol. Microbiol.*, 55(3), 1255-1265. <https://doi.org/10.1099/ijs.0.63484-0>
- Im, W.-T., Hu, Z.-Y., Kim, K.-H., Rhee, S.-K., Meng, H., Lee, S.-T., & Quan, Z.-X. (2012). Description of *Fimbriimonas ginsengisoli* gen. nov., sp. nov. within the Fimbriimonadia class nov., of the phylum Armatimonadetes. *Antonie van Leeuwenhoek*, 102(2), 307-317. <https://doi.org/10.1007/s10482-012-9739-6>
- Iwai, S., Weinmaier, T., Schmidt, B. L., Albertson, D. G., Poloso, N. J., Dabbagh, K., & DeSantis, T. Z. (2016). Piphillin: improved prediction of metagenomic content by direct inference from human microbiomes. *PloS one*, 11(11), e0166104. <https://doi.org/10.1371/journal.pone.0166104>.
- Janssen, P. H. (2006). Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Appl. Environ. Microbiol.*, 72(3), 1719-1728. <https://doi.org/10.1128/AEM.72.3.1719-1728.2006>
- Jensen, M. M., Lam, P., Revsbech, N. P., Nagel, B., Gaye, B., Jetten, M. S., & Kuypers, M. M. (2011). Intensive nitrogen loss over the Omani Shelf due to anammox coupled with dissimilatory nitrite reduction to ammonium. *ISME J.*, 5(10), 1660-1670. <https://doi.org/10.1038/ismej.2011.44>
- Krause, S., Liebetrau, V., Löscher, C. R., Böhm, F., Gorb, S., Eisenhauer, A., & Treude, T. (2018). Marine ammonification and carbonic anhydrase activity induce rapid calcium carbonate precipitation. *Geochim. Cosmochim. Acta*, 243, 116-132. <https://doi.org/10.1016/j.gca.2018.09.018>
- Kümmel, S., Herbst, F.-A., Bahr, A., Duarte, M., Pieper, D. H., Jehmlich, N., Seifert, J., von Bergen, M., Bombach, P., Richnow, H. H., & Vogt, C. (2015). Anaerobic naphthalene degradation by sulfate-reducing Desulfobacteraceae from various anoxic aquifers. *FEMS Microbiol. Ecol.*, 91(3), 13. <https://doi.org/10.1093/femsec/fiv006>
- Langille, M. G., Zaneveld, J., Caporaso, J. G., McDonald, D., Knights, D., Reyes, J. A., Clemente, J. C., Burkepille, D. E., Thurber, R. L. V., & Knight, R. (2013). Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. *Nat. Biotechnol.*, 31(9), 814-821. <https://doi.org/10.1038/nbt.2676>
- Leloup, J., Loy, A., Knab, N. J., Borowski, C., Wagner, M., & Jørgensen, B. B. (2007). Diversity and abundance of sulfate-reducing microorganisms in the sulfate and methane zones of a marine sediment, Black Sea. *Environ. Microbiol.*, 9(1), 131-142. <https://doi.org/10.1111/j.1462-2920.2006.01122.x>
- Liu, J., Sun, F., Wang, L., Ju, X., Wu, W., & Chen, Y. (2014). Molecular characterization of a microbial consortium involved in methane oxidation coupled to denitrification under

- micro-aerobic conditions. *Microb. Biotechnol.*, 7(1), 64-76. <https://doi.org/10.1111/1751-7915.12097>
- Long, C., Lu, X.-L., Gao, Y., Jiao, B.-H., & Liu, X.-Y. (2011). Description of a *Sulfitobacter* strain and its extracellular cyclodipeptides. *Evid.-Based Complementary Altern. Med*, 2011, 24-29. <https://doi.org/10.1155/2011/393752>
- Löscher, C. R., Mohr, W., Bange, H. W., & Canfield, D. E. (2020). No nitrogen fixation in the Bay of Bengal? *Biogeosciences*, 17(4), 851-864. <https://doi.org/10.5194/bg-17-851-2020>
- Luo, H., & Moran, M. A. (2015). How do divergent ecological strategies emerge among marine bacterioplankton lineages? *Trends Microbiol.*, 23(9), 577-584. <https://doi.org/10.1016/j.tim.2015.05.004>
- Lv, X., Yu, J., Fu, Y., Ma, B., Qu, F., Ning, K., & Wu, H. (2014). A meta-analysis of the bacterial and archaeal diversity observed in wetland soils. *Sci. World J.*, 2014, 1-12. <https://doi.org/10.1155/2014/437684>
- Madigan, M., Cox, S. S., & Stegeman, R. A. (1984). Nitrogen fixation and nitrogenase activities in members of the family Rhodospirillaceae. *J. Bacteriol.*, 157(1), 73-78. <https://doi.org/10.1128/JB.157.1.73-78.1984>
- Maltby, J., Sommer, S., Dale, A. W., & Treude, T. (2016). Microbial methanogenesis in the sulfate-reducing zone of surface sediments traversing the Peruvian margin. *Biogeosciences*, 13(1), 283-299. <https://doi.org/10.5194/bg-13-283-2016>
- Manske, A. K., Glaeser, J., Kuypers, M. M., & Overmann, J. (2005). Physiology and phylogeny of green sulfur bacteria forming a monospecific phototrophic assemblage at a depth of 100 meters in the Black Sea. *Appl. Environ. Microbiol.*, 71(12), 8049-8060. <https://doi.org/10.1128/AEM.71.12.8049-8060.2005>
- McCreary Jr, J. P., Yu, Z., Hood, R. R., Vinayachandran, P., Furue, R., Ishida, A., & Richards, K. J. (2013). Dynamics of the Indian-Ocean oxygen minimum zones. *Prog. Oceanogr.*, 112, 15-37. <https://doi.org/10.1016/j.pcean.2013.03.002>
- Meyers, P. A. (1994). Preservation of elemental and isotopic source identification of sedimentary organic matter. *Chem. Geol.*, 114, 289-302. [https://doi.org/10.1016/0009-2541\(94\)90059-0](https://doi.org/10.1016/0009-2541(94)90059-0)
- Mußmann, M., Pjevac, P., Krüger, K., & Dykstra, S. (2017). Genomic repertoire of the Woeseiaceae/JTB255, cosmopolitan and abundant core members of microbial communities in marine sediments. *ISME J.*, 11(5), 1276-1281. <https://doi.org/10.1038/ismej.2016.185>
- Naqvi, S. W. A., Lam, P., Narvenkar, G., Sarkar, A., Naik, H., Pratihary, A., Shenoy, D. M., Gauns, M., Kurian, S., & Damare, S. (2018). Methane stimulates massive nitrogen loss from freshwater reservoirs in India. *Nat. Commun.*, 9(1), 1265-1274. <https://doi.org/10.1038/s41467-018-03607-z>
- Naqvi, S. W. A., Naik, H., Pratihary, A., D'Souza, W., Narvekar, P., Jayakumar, D., Devol, A., Yoshinari, T., & Saino, T. (2006). Coastal versus open-ocean denitrification in the Arabian Sea. *Biogeosciences*, 3(4), 621-633. <https://doi.org/10.5194/bg-3-621-2006>

- Narayan, N. R., Weinmaier, T., Laserna-Mendieta, E. J., Claesson, M. J., Shanahan, F., Dabbagh, K., Iwai, S., & DeSantis, T. Z. (2020). Piphillin predicts metagenomic composition and dynamics from DADA2-corrected 16S rDNA sequences. *BMC genomics*, *21*(1), 1-12. <https://doi.org/10.1186/s12864-020-6537-9>
- Nelson, D., & Sommers, L. (1982). Methods of soil analysis Part 2. Chemical and Microbiological Properties (*Total Carbon, Organic Carbon, and Organic Matter* (Vol. 2, pp. 539-579).
- Okubo, T., Ikeda, S., Yamashita, A., Terasawa, K., & Minamisawa, K. (2009). Pyrosequence read length of 16S rRNA gene affects phylogenetic assignment of plant-associated bacteria. *Microbes. Environ.*, *27*(2), 204-208. <https://doi.org/10.1264/jsme2.ME11258>
- Orsi, W. D., Coolen, M. J. L., Wuchter, C., He, L., More, K. D., Irigoien, X., Chust, G., Johnson, C., Hemingway, J. D., Lee, M., Galy, V., & Giosan, L. (2017). Climate oscillations reflected within the microbiome of Arabian Sea sediments. *Sci. Rep.*, *7*(1), 6040. <https://doi.org/10.1038/s41598-017-05590-9>
- Padilla, C. C., Bristow, L. A., Sarode, N., Garcia-Robledo, E., Ramírez, E. G., Benson, C. R., Bourbonnais, A., Altabet, M. A., Girguis, P. R., & Thamdrup, B. (2016). NC10 bacteria in marine oxygen minimum zones. *ISME J.*, *10*(8), 2067. <https://doi.org/10.1038/ismej.2015.262>
- Pattan, J., Mir, I. A., Parthiban, G., Karapurkar, S. G., Matta, V., Naidu, P., & Naqvi, S. (2013). Coupling between suboxic condition in sediments of the western Bay of Bengal and southwest monsoon intensification: A geochemical study. *Chem. Geol.*, *343*, 55-66. <https://doi.org/10.1016/j.chemgeo.2013.02.011>
- Paulmier, A. D. R.-P. (2009). Oxygen minimum zones (OMZs) in the modern ocean. *Prog. Oceanogr.*, *80*, 113–128. <https://doi.org/10.1016/j.pocean.2008.08.001>
- Penton, C. R., Devol, A. H., & Tiedje, J. M. (2006). Molecular evidence for the broad distribution of anaerobic ammonium-oxidizing bacteria in freshwater and marine sediments. *Appl. Environ. Microbiol.*, *72*(10), 6829-6832. <https://doi.org/10.1128/AEM.01254-06>
- Pitcher, A., Villanueva, L., Hopmans, E. C., Schouten, S., Reichart, G.-J., & Damsté, J. S. S. (2011). Niche segregation of ammonia-oxidizing archaea and anammox bacteria in the Arabian Sea oxygen minimum zone. *ISME J.*, *5*(12), 1896. <https://doi.org/10.1038/ismej.2011.60>
- Pramanik, A., Basak, P., Banerjee, S., Sengupta, S., Chattopadhyay, D., & Bhattacharyya, M. (2016). Metagenomic exploration of the bacterial community structure at Paradip Port, Odisha, India. *Genom. Data*, *7*, 94-96. <https://doi.org/10.1016/j.gdata.2015.12.005>
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O. (2012). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.*, *41*(D1), D590-D596. <https://doi.org/10.1093/nar/gks1219>
- Quince, C., Lanzen, A., Davenport, R. J., & Turnbaugh, P. J. (2011). Removing noise from pyrosequenced amplicons. *BMC Bioinf.*, *12*(1), 38. <https://doi.org/10.1186/1471-2105-12-38>

- Rajpathak, S. N., Banerjee, R., Mishra, P. G., Khedkar, A. M., Patil, Y. M., Joshi, S. R., & Deobagkar, D. D. (2018). An exploration of microbial and associated functional diversity in the OMZ and non-OMZ areas in the Bay of Bengal. *J. Biosci.*, *43*(4), 635-648. <https://doi.org/10.1007/s12038-018-9781-2>
- Ramaswamy, V., & Gaye, B. (2006). Regional variations in the fluxes of foraminifera carbonate, coccolithophorid carbonate and biogenic opal in the northern Indian Ocean. *Deep Sea Res. Part I*, *53*(2), 271-293. <https://doi.org/10.1016/j.dsr.2005.11.003>
- Ramos, H. C., Hoffmann, T., Marino, M., Nedjari, H., Presecan-Siedel, E., Dreesen, O., Glaser, P., & Jahn, D. (2000). Fermentative metabolism of *Bacillus subtilis*: physiology and regulation of gene expression. *J. Bacteriol.*, *182*(11), 3072-3080. <https://doi.org/10.1128/JB.182.11.3072-3080.2000>
- Robinson, R. S., Kienast, M., Luiza Albuquerque, A., Altabet, M., Contreras, S., De Pol Holz, R., Dubois, N., Francois, R., Galbraith, E., & Hsu, T. C. (2012). A review of nitrogen isotopic alteration in marine sediments. *Paleoceanogr.*, *27*(4). <https://doi.org/10.1029/2012PA002321>
- Sarkar, A., Naqvi, S. W. A., Lavik, G., Pratihary, A., Naik, H., Shirodkar, G., & Kuypers, M. M. (2020). Massive Nitrogen Loss Over the Western Indian Continental Shelf During Seasonal Anoxia: Evidence From Isotope Pairing Technique. *Front. Mar. Sci.*, *7*(678), 1-14. <https://doi.org/10.3389/fmars.2020.00678>
- Sarma, V., Krishna, M., Viswanadham, R., Rao, G., Rao, V., Sridevi, B., Kumar, B., Prasad, V., Subbaiah, C. V., & Acharyya, T. (2013). Intensified oxygen minimum zone on the western shelf of Bay of Bengal during summer monsoon: influence of river discharge. *J. Oceanogr.*, *69*(1), 45-55. <https://doi.org/10.1007/s10872-012-0156-2>
- Sarma, V., Kumar, M. D., & Saino, T. (2007). Impact of sinking carbon flux on accumulation of deep-ocean carbon in the Northern Indian Ocean. *Biogeochemistry*, *82*(1), 89-100. <https://doi.org/10.1007/s10533-006-9055-1>
- Schimel, J. P., & Schaeffer, S. M. (2015). Microbial control over carbon cycling in soil. *Front. Microbiol.*, *3*, 348-358. <https://doi.org/10.3389/fmicb.2012.00348>
- Schlitzer, R. (2015). Ocean data view.
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., Lesniewski, R. A., Oakley, B. B., Parks, D. H., & Robinson, C. J. (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.*, *75*(23), 7537-7541. <https://doi.org/10.1128/AEM.01541-09>
- Schunck, H., Lavik, G., Desai, D. K., Großkopf, T., Kalvelage, T., Löscher, C. R., Paulmier, A., Contreras, S., Siegel, H., & Holtappels, M. (2013). Giant hydrogen sulfide plume in the oxygen minimum zone off Peru supports chemolithoautotrophy. *PLoS One*, *8*(8), e68661.
- Shao, M.-F., Zhang, T., & Fang, H. H.-P. (2010). Sulfur-driven autotrophic denitrification: diversity, biochemistry, and engineering applications. *Appl. Microbiol. Biotechnol.*, *88*(5), 1027-1042. <https://doi.org/10.1007/s00253-010-2847-1>

- Shirodkar, G., Naqvi, S. W. A., Naik, H., Pratihary, A. K., Kurian, S., & Shenoy, D. M. (2018). Methane dynamics in the shelf waters of the West coast of India during seasonal anoxia. *Mar. Chem.*, 203, 55-63. <https://doi.org/10.1016/j.marchem.2018.05.001>
- Sorokin, D. Y. (1995). *Sulfitobacter pontiacus* gen. nov., sp. nov.-a new heterotrophic bacterium from the Black Sea, specialized on sulfite oxidation. *Microbiology*, 64(3), 295-295. <https://doi.org/10.1007/s12275-019-9150-3>
- Stewart, F. J., Ulloa, O., & DeLong, E. F. (2012). Microbial metatranscriptomics in a permanent marine oxygen minimum zone. *Environ. Microbiol.*, 14(1), 23-40. <https://doi.org/10.1111/j.1462-2920.2010.02400.x>
- Suh, S.-S., Park, M., Hwang, J., Lee, S., Moh, S. H., Park, K. H., & Lee, T.-K. (2014). Characterization of bacterial communities associated with seasonal water masses from Tongyoung in South Sea of Korea. *Ocean Sci. J.*, 49(3), 193-200. <https://doi.org/10.1007/s12601-014-0019-4>
- Ulloa, O., Wright, J. J., Belmar, L., & Hallam, S. J. (2013). Pelagic oxygen minimum zone microbial communities. In Rosenberg E., DeLong E.F., Lory S., Stackebrandt E., & T. F. (Eds.), *The Prokaryotes* (pp. 113-122). Springer. https://doi.org/10.1007/978-3-642-30123-0_45
- van der Weijden, C. H., Reichart, G. J., & Visser, H. J. (1999). Enhanced preservation of organic matter in sediments deposited within the oxygen minimum zone in the northeastern Arabian Sea. *Deep Sea Res. Part II*, 46(5), 807-830. [https://doi.org/10.1016/S0967-0637\(98\)00093-4](https://doi.org/10.1016/S0967-0637(98)00093-4)
- Van Goethem, M. W., Makhalanyane, T. P., Cowan, D. A., & Valverde, A. (2017). Cyanobacteria and Alphaproteobacteria may facilitate cooperative interactions in niche communities. *Front. Microbiol.*, 8, 2099. <https://doi.org/10.3389/fmicb.2017.02099>
- Varki, A., & Gagneux, P. (2017). Biological functions of glycans (*Essentials of Glycobiology [Internet]. 3rd edition*). Cold Spring Harbor Laboratory Press.
- Wang, Y., Sheng, H.-F., He, Y., Wu, J.-Y., Jiang, Y.-X., Tam, N. F.-Y., & Zhou, H.-W. (2012). Comparison of the levels of bacterial diversity in freshwater, intertidal wetland, and marine sediments by using millions of illumina tags. *Appl. Environ. Microbiol.*, 78(23), 8264-8271. <https://doi.org/10.1128/AEM.01821-12>
- Ward, B., Devol, A., Rich, J., Chang, B., Bulow, S., Naik, H., Pratihary, A., & Jayakumar, A. (2009). Denitrification as the dominant nitrogen loss process in the Arabian Sea. *Nature*, 461(7260), 78-82. <https://doi.org/10.1038/nature08276>
- Wegner, C.-E., Richter-Heitmann, T., Klindworth, A., Klockow, C., Richter, M., Achstetter, T., Glöckner, F. O., & Harder, J. (2013). Expression of sulfatases in *Rhodopirellula baltica* and the diversity of sulfatases in the genus *Rhodopirellula*. *Mar. Genomics*, 9, 51-61. <https://doi.org/10.1016/j.margen.2012.12.001>
- West, N. J., Schönhuber, W. A., Fuller, N. J., Amann, R. I., Rippka, R., Post, A. F., & Scanlan, D. J. (2001). Closely related *Prochlorococcus* genotypes show remarkably different depth distributions in two oceanic regions as revealed by in situ hybridization using 16S rRNA-targeted oligonucleotides. *Microbiology*, 147(7), 1731-1744. <https://doi.org/10.1099/00221287-147-7-1731>

- Yakimov, M. M., Giuliano, L., Gentile, G., Crisafi, E., Chernikova, T. N., Abraham, W.-R., Lünsdorf, H., Timmis, K. N., & Golyshin, P. N. (2003). *Oleispira antarctica* gen. nov., sp. nov., a novel hydrocarbonoclastic marine bacterium isolated from Antarctic coastal sea water. *Int. J. Syst. Evol. Microbiol.*, 53(3), 779-785. <https://doi.org/10.1099/ijs.0.02366-0>
- Yanagibayashi, M., Nogi, Y., Li, L., & Kato, C. (1999). Changes in the microbial community in Japan Trench sediment from a depth of 6292 m during cultivation without decompression. *FEMS Microbiol. Lett.*, 170(1), 271-279. <https://doi.org/10.1111/j.1574-6968.1999.tb13384.x>
- Yu, Z., Wang, X., Han, G., Liu, X., & Zhang, E. (2018). Organic and inorganic carbon and their stable isotopes in surface sediments of the Yellow River Estuary. *Sci. Rep.*, 8(1), 1-10. <https://doi.org/10.1038/s41598-018-29200-4>
- Zhu, D., Tanabe, S.-H., Yang, C., Zhang, W., & Sun, J. (2013). Bacterial community composition of South China Sea sediments through pyrosequencing-based analysis of 16S rRNA genes. *PloS one*, 8(10), e78501. <https://doi.org/10.1371/journal.pone.0078501>