

Isolation of a Novel Cyanophage Infecting Potentially Bloom-forming *Limnothrix* sp.

Judith M. Noronha¹, Manguesh U. Gauns², Sanjeev C. Ghadi^{1*}

1: Department of Biotechnology, Goa University, Taleigao Plateau, Goa, India

2: Biological Oceanography Division, CSIR-National Institute of Oceanography – Goa, Dona Paula, Goa, India

Corresponding Author: Sanjeev C. Ghadi , Email: saga@unigoa.ac.in

ORCID no. : Judith M. Noronha 0000-0003-3376-6013

Abstract:

A filamentous cyanobacterium, identified as *Limnothrix* sp. VL01, and a novel lytic cyanophage ‘Φ L-VL01’ that infects it, were isolated from a shallow freshwater lake located in Goa, India. Cyanophage-infected *Limnothrix* sp. VL01 demonstrated reduced growth and increased fragmentation of filaments, compared to uninfected control. *Limnothrix* sp. is known to form blooms in freshwater lakes. Φ L-VL01 could be a significant agent in controlling such blooms.

Keywords: *Limnothrix*, cyanophage, bloom-forming cyanobacteria.

Cyanobacterial blooms are on the increase worldwide, due to eutrophication of water bodies and rising global temperatures [1]. Filamentous cyanobacteria are frequently responsible for blooms, and lytic cyanophages are important agents of bloom control [2]. *Limnothrix* sp. is a common bloom-forming filamentous cyanobacterium, with a worldwide distribution [3, 4]. It can dominate the cyanobacterial community in shallow, eutrophic lakes, often along with other filamentous forms such as *Planktothrix agardhii*. Further, *Limnothrix* sp. has been identified as a water contaminant, with strains proposed to show toxicity [5, 6]. The present study reports the isolation and characterization of *Limnothrix* sp. VL01 and its corresponding lytic cyanophage ‘L-VL01’, from a freshwater lake located in Goa, India.

Water samples were collected from Verna Lake (15.347689, 3.945478) situated in the state of Goa, India. Verna Lake is the source of the river Sal, the only river which originates in Goa, and is a site favoured by migratory birds. With an average depth of 4 meters, the lake lies proximate to human habitation, is utilized in local agriculture and fishery activities, and is on the verge of eutrophication. From lake water samples, cyanobacterial cultures were isolated in BG-11 medium [7], purified by streaking on BG-11 + 1% agar plates, and made axenic by treatment with Tienam [8]. Cultures were maintained at 25°C, under a 16:8 light:dark cycle at irradiance between 10 and 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$. The strain used in the present study was morphologically examined under a compound microscope. Molecular-level identification was carried out by amplification of the 16S rDNA gene fragment [9], followed by nucleotide BLAST analysis and generation of a Neighbour-Joining phylogenetic tree.

Water samples collected from Verna Lake were further screened for the presence of phages, using the isolated cyanobacteria as host. 0.22 μm -filtered water samples were added to host culture in the standard agar overlay technique [10]. Cyanophage plaques obtained on the agar plates were subjected to several rounds of purification, and thereafter, the pure phage stock was propagated in liquid and solid medium. Lysis of the host cyanobacteria in liquid medium was monitored by measuring the chlorophyll *a* concentration in test and control flasks over a period of 16 days. Chlorophyll *a* was extracted with methanol, measured spectrophotometrically and its concentration calculated by the following formula [11]:

$$\text{Chl}_a [\mu\text{g/ml}] = 12.9447 (A_{665} - A_{720})$$

The phage lysate was concentrated with 10% w/v PEG 8000 and purified by cesium chloride density gradient ultracentrifugation. 10 μl of purified phage was adsorbed onto a formvar-coated copper grid and stained with 2% (w/v) uranyl acetate, followed by viewing with a Jeol JEM 2100 200kV transmission electron microscope [12]. Genomic DNA was extracted from the purified phage lysate,

by the standard phenol-chloroform method [13]. PCR-amplification of the *g20* gene, which encodes the portal protein of viruses belonging to the family *Myoviridae* was carried out with the primers CPS1 and CPS 8 [14].

Cyanophage-infected and uninfected cyanobacterial filaments were observed under 100X and 1000X magnification on a Lynx optical microscope. Additionally, the filaments collected on 0.2- μ m polycarbonate filters, were dehydrated in an ethanol series, mounted on an aluminium stub, sputter-coated with gold particles and observed under a Zeiss Evo 18 scanning electron microscope.

The filamentous cyanobacterium obtained in axenic culture was designated strain VL01. Strain VL01 comprised non-heterocystous, solitary, unsheathed filaments consisting of narrow, long, cylindrical, cells with gas vacuoles (**Fig. S1**). These morphological features were in concurrence with the description of the genus *Limnothrix* [15]. Nucleotide BLAST analysis of the 16S rDNA sequence of strain VL01 (Accession number MN808643.1) demonstrated 99.80% identity with *Limnothrix* sp. Phylogenetic analysis with comparative homologous sequences from GenBank, using the Neighbor-Joining method, indicated that strain VL01 clustered with *Limnothrix redekei* (**Fig. S2**). Thus, the strain VL01 was designated *Limnothrix* sp. VL01.

The genus *Limnothrix* is primarily a freshwater form, which has been isolated from rivers, lakes and ponds in various locations of the world [16–18]. The *Limnothrix* strain used in the present study was isolated from lake water. The serial dilution method used for establishing a unialgal culture, inherently selects for the fastest-growing or most robust form within the specific aquatic niche. This indicates *Limnothrix* sp. VL01 to be a potential bloom-former. Total phosphorus concentrations of 0.15 mg/l have been observed in Verna Lake, which is six-fold above the threshold of 0.024 mg/l, above which increased proliferation of cyanobacterial blooms has been shown to be likely [19, 20].

Subsequent screening of lake water samples against *Limnothrix* sp. VL01 by the agar overlay method, resulted in clear plaques, indicating the presence of a lytic phage (**Fig 1a**). The plaques were purified and the isolated phage was designated ‘ Φ L-VL01’. Φ L-VL01 demonstrated lysis of host in liquid medium (**Fig. 1b**), with an infective pattern similar to phages of other filamentous cyanobacteria [21–23]. Phage-infected host filaments turned from dark green to yellow, indicating their degradation. Further, phage-induced lysis of *Limnothrix* sp. VL01 was incomplete (**Fig. 1b**), and a residual growth of host filaments was observed. Incomplete lysis of filamentous host has been reported previously, in phage-infection of *Planktothrix agardhii* [21] and *Aphanizomenon flos-aquae* [23]. In comparison to the uninfected culture of *Limnothrix* sp. VL01, phage-infected culture demonstrated an 82 percent reduction in growth over 16 days, as indicated by chlorophyll *a*

concentration (**Fig. 1c**). The difference in growth rate between control and infected culture was visible from the second day post-infection. The lysis time reported for phages of filamentous cyanobacteria varies from twenty four hours [24] to eight days [21].

TEM analysis of Φ L-VL01 revealed phage particles having a head-and-tail structure. The icosahedral head measured approximately 50 nm in diameter (**Fig. 2a**). Cyanophages of filamentous cyanobacteria reported in literature have depicted a variety of head and tail morphologies (**Table S1**). The *g20* gene coding for the portal protein of myoviruses was amplified from the lysate of cyanophage L-VL01. **Fig. 2b** shows the PCR product of expected size around 600 bp. Presence of the *g20* marker indicates that L-VL01 belongs to the family *Myoviridae*.

Under a scanning electron microscope, control filaments of strain VL01 were intact, with cells joined end-to-end (**Fig. 3a**), whereas in the case of virus-infected filaments, cells were detached (**Fig. 3b**). A similar pattern had been observed under an optical microscope as well (**Fig. S3**). Fragmentation of host filaments following phage infection has been reported in several cyanobacterial species, notably, *Planktothrix* sp. [21], *Cylindrospermopsis raciborskii* [25] and *Aphanizomenon flos-aquae* [23]. Fragmentation of cyanobacterial filaments (trichomes) into shorter “hormogonia” is a common mode of reproduction in filamentous cyanobacteria [26]. In addition, the formation of hormogonia facilitates dispersal. Hence, the phenomenon of fragmentation following viral infection, could be a survival strategy of the host cyanobacterium.

The incidence of cyanobacterial blooms worldwide has increased, due to increased eutrophication and global warming. Although *Limnothrix* sp. is a known bloom former and toxin producer, there have been limited attempts to isolate phages infecting members of this genus. The only such report was of Φ L-DHS1, infecting *Limnothrix planktonica* strain DH5, which was isolated from Lake Donghu, China [27]. In the context of the present study, the occurrence of *Limnothrix* blooms would be deleterious to the existing aquatic community including fish that thrive in Verna Lake. From this perspective, the isolation of a lytic cyanophage which could regulate uncontrolled proliferation of *Limnothrix* sp., assumes significance. The first reported *Limnothrix* phage, Φ L-DHS1, demonstrated structural uniqueness among tailed phages [27]. The present study has highlighted certain functional aspects of the isolated phage, Φ L-VL01, such as its infectivity characteristics and potential to suppress the growth of its bloom-forming host. Attempts to increase the titre of the phage are in progress, which would facilitate its genome analysis, thus giving further insight into the characteristics of Φ L-VL01.

Although freshwater systems were critical in the discovery of cyanophages [28], a disproportionate number of cyanophage studies today have come from marine environments [29]. The study of freshwater cyanophages from a functional perspective, is important, as they are distinct from their marine counterparts, even though they share the basic viral morphologies [30]. In the Indian context, cyanobacterial blooms, though common, have not been extensively researched. Previous studies have largely concentrated on the production of microcystins [31, 32].

The isolation and characterization of novel viruses is a critical area in aquatic virus research. Laboratory studies on individual virus-host systems enable a better understanding of the contribution of viruses to the aquatic ecology [33]. At present, in the field of aquatic virology, the primary focus is on metagenomic investigations at the ecosystem level. Metagenomic studies are constantly uncovering an array of novel viruses. However, one crucial question that is not answered by metagenomic discovery is “Who is the host”? [34] The basic information required about a new virus is which host it infects. This question is answered by traditional culture-based methods. Thus, laboratory isolation of viruses and their hosts would remain relevant, and continue to complement metagenomic studies.




Fig. 1

Lysis of *Limnothrix* sp. VL01 by Φ L-VL01 in solid (a) and liquid (b) medium
 (c) Growth of control versus virus-infected *Limnothrix* sp. VL01, over 16 days, measured as chlorophyll *a* concentration




Fig. 2

(a) Transmission electron micrograph of Φ L-VL01

(b) Amplification of *g20* gene fragment from genomic DNA of Φ L-VL01




Fig. 3

Uninfected (a) and virus-infected (b) filaments of *Limnothrix* sp. VL01 under a scanning electron microscope;

Declarations

Funding: This work was supported by the Council of Scientific and Industrial Research (CSIR-SRF Sanction no. 09/409(0025)/2013–EMR–I) and Goa University.

Conflicts of interest/Competing interests: Not applicable

Ethics approval: Not applicable

Consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and material: Not applicable

Code availability: Not applicable

CRedit author contribution statement:

Judith M. Noronha: Conceptualization, Investigation, Writing – original draft, Writing – review and editing; **Mangesh U. Gauns:** Writing – review and editing; **Sanjeev C. Ghadi:** Conceptualization, Writing – review and editing.

References

1. Glibert PM (2020) Harmful algae at the complex nexus of eutrophication and climate change. *Harmful Algae* 91:101583. <https://doi.org/10.1016/j.hal.2019.03.001>
2. Gerphagnon M, Macarthur DJ, Latour D, et al (2015) Minireview Microbial players involved in the decline of filamentous and colonial cyanobacterial blooms with a focus on fungal parasitism. <https://doi.org/10.1111/1462-2920.12860>
3. Humpage A, Falconer I, Bernard C, et al (2012) Toxicity of the cyanobacterium *Limnothrix* AC0243 to male Balb/c mice. *Water Res* 46:1576–1583. <https://doi.org/10.1016/j.watres.2011.11.019>
4. Yang Y, Zheng X, Tang Q, et al (2020) Species diversity and seasonal dynamics of filamentous cyanobacteria in urban reservoirs for drinking water supply in tropical China. *Ecotoxicology* 29:780–789. <https://doi.org/10.1007/s10646-020-02189-1>
5. Daniels O, Fabbro L, Makiela S (2014) The effects of the toxic cyanobacterium *Limnothrix* (Strain AC0243) on *Bufo marinus* larvae. *Toxins (Basel)* 6:1021–1035. <https://doi.org/10.3390/toxins6031021>
6. Rose AK, Fabbro L, Kinnear S (2018) Cyanobacteria breakthrough: Effects of *Limnothrix redekei* contamination in an artificial bank filtration on a regional water supply. *Harmful Algae* 76:1–10. <https://doi.org/10.1016/j.hal.2018.04.010>
7. Rippka R, Deruelles J, Waterbury JB, et al (1979) Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *J Gen Microbiol* 111:1–61. <https://doi.org/10.1099/00221287-111-1-1>
8. Sarchizian I, Ardelean II (2010) Improved Lysozyme Method To Obtain Cyanobacteria in Axenic Cultures. *Rom J Biol Plant Biol* 55:143–150. <https://doi.org/https://www.ibiol.ro/plant/volume%2055/art209.pdf>
9. Nubel U, Garcia-pichel F, Muyzer G (1997) PCR Primers To Amplify 16S rRNA Genes from Cyanobacteria. *Appl Environ Microbiol* 63:3327–3332. <https://doi.org/https://doi.org/10.1128/aem.63.8.3327-3332.1997>
10. Bubeck JA, Pfitzner AJP (2005) Isolation and characterization of a new type of chlorovirus that infects an endosymbiotic *Chlorella* strain of the heliozoon *Acanthocystis turfacea*. *J Gen Virol* 86:2871–2877. <https://doi.org/10.1099/vir.0.81068-0>
11. Zavřel T, Červený J, Sinetova MA (2015) Measurement of Chlorophyll a and Carotenoids Concentration in Cyanobacteria. *Bio-protocol* 5(9) e1467 5:1–5. <https://doi.org/10.21769/BioProtoc.1467>
12. Ackermann H-W (2009) Phage classification and characterization. *Methods Mol Biol* 501:127–140. https://doi.org/10.1007/978-1-60327-164-6_13
13. Sambrook J, Russell DW (2006) Extraction of Bacteriophage λ DNA from Large-scale Cultures Using Proteinase K and SDS. *Cold Spring Harb Protoc* 2006:pdb-prot3972. <https://doi.org/10.1101/pdb.prot3972>

14. Wang G, Murase J, Asakawa S, Kimura M (2010) Unique viral capsid assembly protein gene (g20) of cyanophages in the floodwater of a Japanese paddy field. *Biol Fertil Soils* 46:93–102. <https://doi.org/10.1007/s00374-009-0410-y>
15. Meffert ME (1988) *Limnothrix* Meffert nov. gen. The unsheathed planktic cyanophycean filaments with polar and central gas-vacuoles. *Arch für Hydrobiol* , Suppl 80:269–276
16. Cunha de Oliveira ED, Da Cunha AC, Da Silva NB, et al (2019) Morphological and molecular characterization of cyanobacterial isolates from the mouth of the amazon river. *Phytotaxa* 387:269–288. <https://doi.org/10.11646/phytotaxa.387.4.1>
17. Zhu M, Yu G, Li X, et al (2012) Taxonomic and phylogenetic evaluation of *Limnothrix* strains (Oscillatoriales, Cyanobacteria) by adding *Limnothrix planktonica* strains isolated from central China. *Hydrobiologia* 698:367–374. <https://doi.org/10.1007/s10750-012-1127-8>
18. Ojit SK, Thadoi DA, Indrama T, et al (2015) Fatty acid profiling of filamentous non-heterocystous cyanobacteria from Loktak Lake , the largest freshwater lake in North-Eastern region of India. *Int Res J Biol Sci* 4:69–74
19. Oram B (2019) Phosphates in the Environment. In: *Water Res. Cent. - Phosphates*. <https://water-research.net/index.php/phosphates%0Ahttps://www.water-research.net/index.php/phosphates>
20. Vuorio K, Järvinen M, Kotamäki N (2020) Phosphorus thresholds for bloom-forming cyanobacterial taxa in boreal lakes. *Hydrobiologia* 847:4389–4400. <https://doi.org/10.1007/s10750-019-04161-5>
21. Gao E Bin, Yuan XP, Li RH, Zhang QY (2009) Isolation of a novel cyanophage infectious to the filamentous cyanobacterium *Planktothrix agardhii* (Cyanophyceae) from Lake Donghu, China. *Aquat Microb Ecol* 54:163–170. <https://doi.org/10.3354/ame01266>
22. Pollard PC, Young LM (2010) *Acta Oecologica* Lake viruses lyse cyanobacteria , *Cylindrospermopsis raciborskii* , enhances filamentous-host dispersal in Australia. *Acta Oecologica* 36:114–119. <https://doi.org/10.1016/j.actao.2009.10.007>
23. Šulčius S, Šimoliunas E, Staniulis J, et al (2015) Characterization of a lytic cyanophage that infects the bloom-forming cyanobacterium *Aphanizomenon flos-aquae*. *FEMS Microbiol Ecol* 91:1–7. <https://doi.org/10.1093/femsec/fiu012>
24. Zhou Y, Lin J, Li N, et al (2013) Characterization and Genomic Analysis of a Plaque Purified Strain of Cyanophage PP. 28:272–279. <https://doi.org/10.1007/s12250-013-3363-0>
25. Steenhauer LM, Wierenga J, Carreira C, et al (2016) Isolation of cyanophage CrV infecting *Cylindrospermopsis raciborskii* and the influence of temperature and irradiance on CrV proliferation. *Aquat Microb Ecol* 78:11–23. <https://doi.org/10.3354/ame01800>
26. Komárek J, Johansen JR (2015) Filamentous Cyanobacteria. In: *Freshwater Algae of North America: Ecology and Classification*. Elsevier, pp 135–235

27. Xiangling L, Ying H, Yijun Z, Kai C (2015) A novel freshwater cyanophage with a complex collar infects *Limnothrix planktonica* . *Phycologia* 54:578–582. <https://doi.org/10.2216/15-82.1>
28. Safferman RS, Morris M-E (1963) Algal virus: Isolation. *Science* (80-) 140:679–680. <https://doi.org/10.1126/science.140.3567.679>
29. Šulčius S, Šimoliūnas E, Alzbutas G, et al (2019) Genomic Characterization of Cyanophage vB_AphaS-CL131 Infecting Filamentous Diazotrophic Cyanobacterium *Aphanizomenon flos-aquae* Reveals Novel Insights into Virus-Bacterium Interactions. *Appl Environ Microbiol* 85:e01311-18. <https://doi.org/10.1128/AEM.01311-18>
30. Morimoto D, Šulčius S, Yoshida T (2020) Viruses of freshwater bloom-forming cyanobacteria: genomic features, infection strategies and coexistence with the host. *Environ Microbiol Rep* 12:486–502. <https://doi.org/10.1111/1758-2229.12872>
31. Chaturvedi P, Kumar Agrawal M, Nath Bagchi S (2015) Microcystin-producing and non-producing cyanobacterial blooms collected from the Central India harbor potentially pathogenic *Vibrio cholerae*. *Ecotoxicol Environ Saf* 115:67–74. <https://doi.org/10.1016/j.ecoenv.2015.02.001>
32. Sangolkar LN, Maske SS, Muthal PL, et al (2009) Isolation and characterization of microcystin producing *Microcystis* from a Central Indian water bloom. *Harmful Algae* 8:674–684. <https://doi.org/10.1016/j.hal.2008.12.003>
33. Weitz JS, Poisot T, Meyer JR, et al (2013) Phage-bacteria infection networks. *Trends Microbiol* 21:82–91. <https://doi.org/10.1016/j.tim.2012.11.003>
34. Edwards RA, McNair K, Faust K, et al (2016) Computational approaches to predict bacteriophage-host relationships. *FEMS Microbiol Rev* 40:258–272. <https://doi.org/10.1093/femsre/fuv048>




Fig. S3 Uninfected (**a**) and virus-infected (**b**) filaments of *Limnothrix* sp. VL01 under an optical microscope

Supplementary Table 1:

Comparative dimensions of phages of filamentous cyanobacteria reported in literature

Cyanophages	Host	Head diameter (nm)	Observed tails	References
Various	<i>Nodularia</i> sp.	53-137	Long	[1]
Pf-WMP4	<i>Phormidium foveolarum</i>	55	Short	[2]
Various	<i>Anabaena</i> sp.	Not provided	Mostly short	[3]
PaV-LD	<i>Planktothrix agardhii</i>	76	Tailless	[4]
Cr-LS	<i>Cylindrospermopsis raciborskii</i>	70	Long	[5]
Various	<i>Anabaena circinalis</i> , <i>A. cylindrica</i>	123-170	Mostly short	[6]
L-DHS1	<i>Limnothrix</i> sp.	72	Long	[7]
CL131	<i>Aphanizomenon flos-aquae</i>	97	Long	[8]
CrV	<i>Cylindrospermopsis raciborskii</i>	65	Long	[9]
PA-SR01	<i>Pseudoanabaena</i>	91	Tailless	[10]
Aa-TR020	<i>Arthronema africanum</i>	50	Short	[11]
L-VL01	<i>Limnothrix</i> sp.	49	Short	Present study

Supplementary References:

1. Jenkins CA, Hayes PK (2006) Diversity of cyanophages infecting the heterocystous filamentous cyanobacterium *Nodularia* isolated from the brackish Baltic Sea
2. Liu X, Shi M, Kong S, et al (2007) Cyanophage Pf-WMP4, a T7-like phage infecting the freshwater cyanobacterium *Phormidium foveolarum*: Complete genome sequence and DNA translocation. *Virology* 366:28–39. <https://doi.org/10.1016/j.virol.2007.04.019>
3. Deng L, Hayes PK (2008) Evidence for cyanophages active against bloom-forming freshwater cyanobacteria. *Freshw Biol* 53:1240–1252. <https://doi.org/10.1111/j.1365-2427.2007.01947.x>
4. Gao E Bin, Yuan XP, Li RH, Zhang QY (2009) Isolation of a novel cyanophage infectious to the filamentous cyanobacterium *Planktothrix agardhii* (Cyanophyceae) from Lake Donghu,

- China. *Aquat Microb Ecol* 54:163–170. <https://doi.org/10.3354/ame01266>
5. Pollard PC, Young LM (2010) Acta Oecologica Lake viruses lyse cyanobacteria , *Cylindrospermopsis raciborskii* , enhances filamentous-host dispersal in Australia. *Acta Oecologica* 36:114–119. <https://doi.org/10.1016/j.actao.2009.10.007>
 6. Yeo BH, Gin KY (2013) Cyanophages infecting *Anabaena circinalis* and *Anabaena cylindrica* in a tropical reservoir. *Bacteriophage* 3:. <https://doi.org/10.4161/bact.25571>
 7. Xiangling L, Ying H, Yijun Z, Kai C (2015) A novel freshwater cyanophage with a complex collar infects *Limnothrix planktonica* . *Phycologia* 54:578–582. <https://doi.org/10.2216/15-82.1>
 8. Šulčius S, Šimoliunas E, Staniulis J, et al (2015) Characterization of a lytic cyanophage that infects the bloom-forming cyanobacterium *Aphanizomenon flos-aquae*. *FEMS Microbiol Ecol* 91:1–7. <https://doi.org/10.1093/femsec/fiu012>
 9. Steenhauer LM, Wierenga J, Carreira C, et al (2016) Isolation of cyanophage CrV infecting *Cylindrospermopsis raciborskii* and the influence of temperature and irradiance on CrV proliferation. *Aquat Microb Ecol* 78:11–23. <https://doi.org/10.3354/ame01800>
 10. Lin W, Li D, Sun Z, et al (2020) A novel freshwater cyanophage vB_MelS-Me-ZS1 infecting bloom-forming cyanobacterium *Microcystis elabens*. *Mol Biol Rep* 47:7979–7989. <https://doi.org/10.1007/s11033-020-05876-8>
 11. Petrzik K, Lukavský J, Koloniuk I (2021) Novel Virus on Filamentous *Arthrocnemum africanum* Cyanobacterium. *Microb Ecol* 81:454–459. <https://doi.org/10.1007/s00248-020-01599-2>