

## **Beneficial properties of mucus in coral adaptations and ecological interactions**

**Phartade Vilas Bhagwat<sup>a,b</sup>, Chinnarajan Ravindran<sup>a,c\*</sup>, Lawrance Irudayarajan<sup>a,c</sup>,**

<sup>a</sup>Biological Oceanography Division, CSIR-National Institute of Oceanography,  
Dona Paula, Goa 403004, India.

<sup>b</sup>Department of Biotechnology, Goa University, Taleigao Plateau, Goa 403206, India.

<sup>c</sup>Academy of Scientific and Innovative Research (AcSIR), Ghaziabad, India.

\*Corresponding author's email: cravindran@nio.org

### **Abstract**

Coral mucus secreted by mucocytes provides a protective physicochemical and physiological barrier between coral tissue and external environmental threats. The biomolecules and nutrients of the secreted mucus are derived from endosymbionts and coral polyps and support coral functions such as feeding, sediment clearing and protection against numerous biotic and abiotic stressors. The surface mucus layer also houses a diverse community of beneficial microorganisms that defend against pathogens. Enzymes including peptidases, esterases, and glycosidases were observed and described in mucus. Most importantly, the presence of phenoloxidase, an intracellular enzyme in secretory mucus, generally triggers melanin synthesis, providing fast-acting components of invertebrate immunity in disease resistance. However, the purpose and mechanism of mucus release, effects of mucus components on defense properties, and functional roles in intra, and interspecific interactions are not well described. Thus, the present review aims to understand the mucus components and the complex roles played by mucus in microbial associations, feeding, and resilience that influence coral health.

**Keywords-** coral, mucus, nutrients, environment, pathogens, innate immunity

## Introduction

According to recent studies, 20 % of coral reefs have already been irreparably damaged, and additionally 24 % of the remaining reefs are facing serious threats due to anthropogenic stresses such as overfishing and eutrophication (Sharma and Ravindran 2020; Galli et al. 2021). Biotic and abiotic stressor can cause coral bleaching, a process in which the algal symbionts are expelled from the host coral (Hafezi et al. 2020; Ravindran et al., 2022). The absence of the endosymbiont will weaken coral health, and mucus secretion will be affected (Ravindran et al., 2022). Environmental impacts, including temperature rise, on the secretion of mucus have not been studied in detail (Vaughan et al. 2021). For example, mucus release is enhanced during high sedimentation (Reynaud and Ferrier-Pages 2019), and when corals are exposed to air at low tide (Wild et al. 2004; Wild et al. 2005). In addition, mucus aggregations in the water column increase sedimentation and nutrient recycling (Wild et al. 2005; Plass-Johnson et al. 2015), which control microbial activities (Bythell and Wild 2011). Therefore, a better understanding of the changes in mucus caused by biological and physical stressors may help to monitor coral health under different environmental conditions.

Mucus is a slimy substance produced by epithelial cells called mucocytes by utilizing the organic carbon synthesized by photosynthetic endosymbiotic algae (Brown and Bythell 2005; Bakshani et al. 2018). Mucus production varies between and within coral species (Coffroth 1990; Hadaidi et al. 2019). It is mainly comprised of proteolytic enzymes, microbes, mucins, proteins, polysaccharides, lipids and their substructures, such as monosaccharides, amino acids and mycosporines (Brown and Bythell 2005; Stabili 2019; Vilas Bhagwat et al. 2023). The beneficial microbes within the mucus form a barrier against potential coral pathogens, competing for nutrients and space or producing antibiotics (Ritchie 2006; Krediet et al. 2013; Raina et al. 2016; Irudayarajan et al. 2023). Coral mucus has several important functions, including protection against pathogens and UV radiation, capturing food, acting as an energy carrier, producing antibiotics, and serving as a nutrient source for coral-associated microbial communities (Bakshani et al. 2018; Bythell and Wild 2011; Vilas Bhagwat et al. 2023). Coral mucus contains up to 1000 times more bacteria than seawater (Nakajima et al. 2015). *Vibrio shiloi*, the causative agent of bleaching in the Mediterranean coral, *Oculina patagonica*, bind to the receptors of the surface mucus layer during temperature rise (Thompson et al. 2015). However, it has been demonstrated that beneficial probiotic bacteria enable *O. patagonica* to become resistant to *V. shiloi* infection (Reshef et al. 2006). Thus, studying the components of coral mucus is of great importance for a better understanding of the entire mucosal system, its interactions with other microorganisms (commensals and pathogens), and their involvement in coral immunity and health. Further, although the composition of coral mucus has been described in

some studies (Coddeville et al. 2011; Hadaidi et al. 2019), that is limited to a few coral species and the sources of mucus components in mucus are not clearly understood. Therefore, the present study aims to describe the significant role of coral mucus in maintaining the health and functionality of coral reef ecosystems, especially the synthesis and constituents of coral mucus and their critical roles in protecting coral health.

### **Coral mucus synthesis and secretion**

The mucocyte cells of the coral polyp synthesize mucus, with the major component being carbon-based carbohydrates, and is then secreted out as a coral surface layer (Brown and Bythell 2005). The coral-associated endosymbiotic algae are believed to provide the maximum carbon required for mucus synthesis, although the source of the entire carbon requirement is not clear (Crossland 1987; Brown and Bythell 2005; Fang 2015) (Fig.1). The other compounds required for mucus, such as amino acids, fatty acids, lipids, glycerol and nicotinamide adenine dinucleotide phosphate (NADPH) come from the coral diet (Hadaidi 2018). Only mucocytes contribute to mucus synthesis in corals, unlike in humans where multiple cell types are involved (Hadaidi et al. 2019; McShane et al. 2021). Some other gland cells are observed in coral tissues, but their roles are still unclear (Bruckner 2015). However, in vertebrates, the transformation of similar gland cells to mucus cells has been observed, which may be because these cells are either specialized secretory cells or developing mucocytes (Que 2015). The distribution of mucocytes varies in both abundance and location, not just within individual colonies but also between coral species (Baker et al. 2022). The production of coral mucus can be triggered by various factors, such as changes in light intensity, water flow, and the presence of predators or food particles (Zetsche et al. 2016). For example, an increase in the sea surface temperature increases the mucocyte density in host tissue, however, it decreases after bleaching (Piggot et al. 2009). In addition, more organic components are found in mucus from stressed than in healthy corals (Lee et al. 2016).

The synthesis of coral mucus is unique because it is derived from a significant portion of the photosynthetically fixed carbon (Hadaidi et al. 2019; Brown and Bythell 2005), and contains a high concentration of arabinose and xylose, which is not typically found in animal cells (Campana et al. 2022). Different coral species have varied rates of mucus production (Crossland et al. 1980; Kurihara et al. 2018), depending on the endosymbiont contribution of energy-rich glycerol, alanine, and glucose to the host (Burriesci et al. 2012; Gabay 2018) (Fig.1). Excess photosynthetic carbon is eliminated by the coral through mucus release (Bakshani et al. 2018) along with lipids, such as wax esters and triglycerides for mucus synthesis (Krueger et al. 2018). *Acropora* spp. release up to 4.8 liters of mucus from a 1m<sup>2</sup> patch in one day and 56-80 % of the mucus is dissolved in water (Wild et al. 2004), of which 20-40 % of mucus forms strings and filaments (Stabili 2019). This mucus net filters seawater

up to 100 ml cm<sup>2</sup> of tissue per hour (Lewis 1976). Subsequently, heterotrophic coral reef organisms utilize the energy from the materials trapped in the coral mucus and the energy derived from zooxanthellae (Wild et al. 2004). Invertebrate mucus protects, lubricates, and helps in locomotion, mating, homing, and foraging, performing much more diverse functions than in vertebrates (Denny 1989). In addition, mucus released on the epithelial surface aims to attract particulate matter.

## **Mucus composition**

### **Mucins and the importance of mucus rheology**

The glycoprotein core, mucin, is made up of 20 % polymers and 80 % sugars and contains cysteine, which undergoes cleavage during mucus dispersion (Bakshani et al. 2018). O-glycosylation of mucin occurs on oligosaccharides in the core between serine and threonine and on the N-acetylgalactosamine carbohydrate chains, and N-glycosylation occurs on asparagine (Corfield 2015). The side chains of mucins are either branched or linear and up to 20 monosaccharides long and are linked with either  $\beta 1 \rightarrow 2$  or  $\beta 1 \rightarrow 3$  bonds (Krediet et al. 2009). The structural diversity of mucin is due to its carbohydrate composition (Thornton and Sheehan 2004). The ability of mucin to form a gel affects mucus function depending on its extensive glycosylation and polymerization ability (Younan et al. 1982). The Von Willebrand D (VWD) domain is involved in the polymerization of mucin monomers via intermolecular disulfide bonds (Ambort et al. 2012). The outer layer of the epidermis of fishes continuously produces new goblet cells because they cannot produce mucus more than once, but coral mucus goblet cells can continuously produce mucus from the same cells (Subramanian et al. 2007).

Corals require relatively fluid mucus for their ciliary feeding and to prevent the desiccation of coral polyps by air exposure during low tide (Brown and Bythell 2005; Bessell-Browne et al. 2017). Based on rheology, coral mucus is distributed as fluid mucus, string, web or flocs, and mucus sheets (Bessell-Browne et al. 2017). Distinct mucin characteristics are present across various layers within the coral surface mucus layer (SML) (Guppy et al. 2019). The mucin concentration and gel-forming properties of mucus are necessary for its protective and lubricating functions (Thornton and Sheehan 2004). Human mucus becomes hydrated and mucus contents turn it into a gel (Piludu et al. 2003). As polymeric mucins have been detected in coral mucus, similar to vertebrates (Jatkar et al. 2010a), it is possible that coral mucus also contains different types of mucin molecules that convert it from gel to liquid and liquid to gel as needed (Horricks et al. 2019). Because coral mucus is rich in plant-associated sugar arabinose, it may have some carbohydrate chains that differ from those of mammalian mucins (Wild et al. 2005). Human mucus changes its rheology from viscoelastic gel to viscous fluid at high

stress and returns to a viscous gel at low stress (Jatkar et al. 2010b) making it a lubricant and helping it move under various physiological conditions (Hill et al. 2022). A similar property may also be present in the coral SML, which remains to be investigated. However, corals can produce two types of mucus. The milked mucus has no gel-forming properties but can be converted to a gel by a higher concentration of mucin (Jatkar et al. 2010a), which is very similar to human respiratory mucus (Zetsche et al. 2016). Mucus sheets are sticky mucus layers derived from liquid mucus (Bessell-Browne et al. 2017). These are then transferred into the food web of the reef (Brown and Bythell 2005).

## Carbohydrates

The relative concentration of carbohydrates varies among coral species, yet their composition remains constant, with certain sugars being shared across multiple species, and this variability is intricately linked to host phylogeny, microbial associations, and symbiont composition (Hadaidi et al. 2019). The mucosal sugars glucose, mannose, xylose, and galactose are found in all coral species that have been analyzed, whereas arabinose, N-acetylglucosamine, ribose, fucose, and rhamnose are present in only a few species (Table 1). Thermal stress has been shown to induce changes in both the enrichment and abundance of various carbohydrate pools, with a notable impact on galactose levels (Hillyer et al. 2017). For example, a significant decrease in glucose, galactose, and mannose were described under rising temperature-stress conditions (Lee et al. 2016). Thus, mannose, glucose, and galactose can serve as markers for temperature stress, while N-acetyl glucosamine exhibits anti-inflammatory properties and reduces the production of free radicals (Salvatore et al. 2000). N-acetyl galactosamine also helps in the O-glycosylation of coral mucin molecules. Xylose and arabinose are types of plant sugars commonly utilized as energy sources, indicating that they may have come from the endosymbiont (Teplitski et al. 2016). Arabinose is widespread in *Acropora* spp., suggesting that it is transferred from algal symbionts to the host (Wild et al. 2005; Wild et al. 2010; Lee et al. 2016).

The carbohydrate composition of mucus differs among the corals *Acropora pharanois*, *Porites verrucosa*, and *Porites lobata* (Hadaidi et al. 2019). However, *Stylophora pistillata* and *Pocillopora verrucosa*, two pocilloporid corals, have similar carbohydrate (Neave et al. 2016) and zooxanthellae composition (LaJeunesse et al. 2018). Differences in mucus components could affect microbial community composition in coral reef habitats (Schottner et al. 2009). The most common mucus components observed in corals that occur across a large geographic range (e.g., *A. muricata*, *Pachyseris speciosa*, *Fungia fungites*, *Stylophora* sp., and *Pocillopora* sp.) are fucose, arabinose, glucose, galactose, the amino sugar N-acetylglucosamine, xylose, and mannose (Wild et al. 2005; Lee et al. 2016) (Table 1). Thus, mucus carbohydrates are essential substrates that support bacterial growth and contribute to the flux of dissolved organic matter (DOM) in seawater (Riemann and Azam 2002).

## **Fatty acids of mucus**

The precise understanding of the fatty acid (FA) composition and its role in coral mucus is still inadequate and remains one of the least studied aspects of coral mucus. It was reported that the FA contents of *Acropora* and *Porites* species were almost the same but the proportions were different not only in the mucus but also in comparison with the tissues (Ravindran et al. 2020). The most common and abundant FAs observed in the mucus of *Acropora* sp. in Discovery Bay, Jamaica, were palmitic acid and palmitoleic acid (Means and Sigleo 1986; Wild et al. 2005). These were also found more frequently in corals and zooxanthellae as structural and storage lipids (Patton et al. 1977; Means and Sigleo 1986). Other FAs, such as myristic acid and linoleic acid have also been observed in mucus, but in lower amounts (Wild et al. 2005). The most abundant saturated FAs in marine organisms are palmitic acid (C16:0), palmitoleic acid (C16:1), and myristic acid (C14:0), similar to those found in certain unicellular marine algae (Uslu et al. 2021). Even though the presence of FAs in mucus has been reported, their origins have not been documented.

FAs in coral mucus are essential for the health and functioning of coral reefs, providing energy, supporting microbial communities, and aiding in defense mechanisms (Bachok et al. 2006). Polyunsaturated fatty acids (PUFAs), especially n-3 and n-6 PUFAs, regulate metabolism, including growth, respiration, energy generation, energy reserves, membrane constituents, photosynthesis, planula larvae and egg production, biomass formation and general enzyme activity, survival, and stress resistance (Bachok et al. 2006). C20:5n-3 FAs are essential for sperm maturation, C20:4n-6 is essential for egg maturation and saturated (SAFA) and monounsaturated FAs (MUFA) are stored as triglycerides and wax esters (Grottoli et al. 2004; Tchernov et al. 2004; Treignier et al. 2008). In all of the previously studied coral mucus samples, higher concentrations of SAFAs were observed (Fig. 2). In addition, FAs in corals serve as markers for bacterial colonization and diversity, coral health, and food source tracking (Volkman et al. 1980). As the synthesis of FAs in corals and in coral mucus is directly coupled to photosynthesis, there is a significant correlation between dinoflagellate biomarkers and FA concentrations (Oku et al. 2003). Free fatty acids (FFAs) have been found to be involved in maintaining cell membrane stability and cell signaling (Binienda et al. 2013). FFAs are physiologically active as dietary components and participate in cell signaling (Zbigniew et al. 2013). FAs are also required for reproduction and membrane fluidity (Lin et al. 2012). However, the purpose and function of FAs present in mucus are not well understood and need to be thoroughly investigated in various genera and species of corals based on their habitat and other environmental parameters.

## **Mucus lipids**

Photosynthetic carbon is stored in the cytoplasm of zooxanthellae, in the form of lipids, which are then incorporated into the coral mucus, where it serves as an important source of energy for a variety of marine organisms (Crossland et al. 1980; Pasaribu et al. 2015). Many galactolipids synthesized by marine diatoms have been shown to induce apoptosis starting at micromolar concentrations in mammalian cells (Andrianasolo et al. 2008), suggesting the function of membrane lipids in regulating the symbiosis between corals and zooxanthellae. Coral mucus is composed of high-energy lipids which are also an important source of energy for fish (Wright et al. 2019). Upon thermal stress, corals may acquire more thermotolerant symbionts (Lajeunesse et al. 2010) having lipids with more SAFAs with a high melting point. This altered membrane lipid composition is observed in bleached corals, indicating differences in heat sensitivity among zooxanthellae lineages (Tchernov et al. 2004).

Zooxanthellae provides precursors and metabolites required for endogenous lipid synthesis in coral hosts mainly through photosynthesis (Oku et al. 2003; Iluz and Dubinsky 2015), and several studies have demonstrated the rapid conversion of photosynthetically fixed carbon to lipids (Radice et al. 2019). The presence of lipids (e.g., wax esters) in coral mucus has been demonstrated in significant amounts (Crossland 2021). Corals are rich in lipids, which are the biochemical link between the symbiotic mutualistic association of coral and zooxanthellae (Sikorskaya et al. 2022). Lipids are involved in coral physiological and biochemical processes and are an important source of long-term stored energy as mucus serves as an energy-rich source of nutrition for various organisms (Rodrigues et al. 2008). During bleaching, a change in FA composition and also a reduction in the rate of lipid production may affect the coral's metabolic needs (Grottoli et al. 2004). This reduced energy reserve increases disease susceptibility and mortality (Smith et al. 2020), which may be due to a change in the microbial community in the damaged tissues.

### **Proteins/peptides of mucus**

According to Benavides et al. (2017), a low C: N ratio in coral mucus indicates the presence of a large amount of protein. Previous studies on *Palythoa* sp. and *Fungia* sp. showed that their mucus contains 8-20% organic matter and 1-2% protein (Daumas and Thomassin 1977; Stabili et al. 2015). Coral mucus from *Acropora*, *Fungia*, and *Sarcophyton* species contains important enzymes such as peptidases, esterases, and glycosidases (Vacelet and Thomassin 1991; Vilas Bhagwat et al. 2023). Hydroxyproline, found in coral mucus, is crucial for collagen synthesis and maintaining the thermodynamic stability of collagen's triple-helical conformation and associated tissues (Ortega et al. 2018). Fish mucus, on the other hand, is known for its high protein content and enzymes such as lysozyme, alkaline phosphatase, complement, cathepsin B, transferrin, and C-reactive protein

(Subramanian et al. 2007; Zhao et al. 2008). The mucus of the zoanthid *Palythoa caribaeorum* contains lectins and proteases (Guarnieri et al. 2018). However, phenoloxidase (PO), peroxidases (POD), and antioxidant enzymes like superoxide dismutase, catalase, and glutathione peroxidase, which have been reported from coral tissues (Anithajothi et al. 2014), have not yet been reported in coral mucus. Thus, mucus reported with antimicrobial properties has to be investigated for its potential peptides and proteins.

### **Amino acids**

Corals and zooxanthellae can synthesize 20 different amino acids, with some produced by the coral polyps, others by the zooxanthellae, and yet others by both partners. However, different coral species have different amino acid synthesis abilities (Fitzgerald and Szmant 1997). Amino acids can also be taken up via plankton, detritus, and the water column. The acquired amino acids are used by the coral to synthesize proteins for tissue growth, calcification, and other processes (Grover et al. 2008; Shinzato et al. 2014). The production of methionine could be a result of the synergistic action of enzymes from both coral and zooxanthellae, indicating that these symbiotic partners have evolved to mutually benefit each other (Shinzato et al. 2014). Although corals acquire many amino acids, only 16 have been observed in the mucus, including aspartate, glutamine, serine, glycine, threonine, arginine, alanine, tyrosine, valine, phenylalanine, isoleucine, leucine, lysine, histidine and methionine, of several coral species (i.e., *Acropora formosa*, *Pachyseris peciosa*, *Fungia fungites*, *Sarcophyton* sp., *Lemnalia* sp., *Cespitularia* sp. *Platygyra amellina*, *Heteroxenia fuscesens*, *Oculina arbuscula* (Meikle et al. 1988; Ducklow and Mitchell 1979; Means and Sigleo 1986; Coddeville et al. 2011).

Analysis of mucus produced by *Palythoa* and *Fungia* species has shown that the amino acids serine, aspartic acid, glutamic acid, and glycine are present in high abundance, and these play important roles in a variety of biological processes, including protein synthesis, cellular signaling, and energy metabolism (Daumas and Thomassin 1977). Glycine is the most abundant amino acid found in both coral tissues and mucus, particularly in species such as *Montastraea cavernosa* (Daumas and Thomassin 1977). Glutamic acid which also acts as an antioxidant in many plants, animals, and bacteria, was more abundant in the mucus of *Fungia* sp., *Acropora* sp., *Platygyra* sp., *Sarcophyton* sp., and *Heteroxenia* sp., compared to serine, and methionine (Ducklow and Mitchell 1979). Alanine is generally observed in *Porites divaricata*, *Acropora cervicornis*, *Orbicella faveolata*, and the azooxanthellate corals, *Astrangia poculata* and *Tubastraea coccinea* (Fitzgerald and Szmant 1997). Similarly, *Fungia scutaria* was reported to utilize tyrosine, lysine, glycine and aspartic acid, from surrounding water (Stephens 1962). This suggests that corals acquire some amino acids through



heterotrophic feeding from the surrounding water. In addition, both host and zooxanthellae can synthesize amino acids and this varies from species to species.

### **Mucus antioxidants against oxidative stress**

A decrease in zooxanthellae density is due to oxidative stress causing coral bleaching (Downs et al. 2002; Weis 2008). Primarily, oxidative stress is generated during photosynthesis (Krieger-Liszkay 2005). Excessive production of reactive oxygen species (ROS) causes disorganization and rupture of the thylakoid membrane (Tchernov et al. 2004). Free radicals such as hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and hydroxyl radicals (-OH) degrades mucus glycoproteins at the histidine molecule, so that mucus property, molecular weight, and viscosity is affected (Dailah 2022). ROS imparts an adverse impact on the mucus defense properties which are then controlled by the mucus antioxidants (Suggett et al. 2008).

The presence of phenoloxidase (PO), an enzyme that activates the invertebrate defense system, was reported in coral secretory mucus (Rivera-Ortega and Thome 2018; Vilas Bhagwat et al. 2023). The mechanism of secretion of this enzyme into coral mucus is not known. However, the expulsion of similar molecules from tissue to the mucus was reported previously (Ritchie 2006). Thus, phenoloxidase is thought to be an intracellular enzyme, but similar activity was also observed in the mucus of other organisms such as in lancelets (Zhang and Li 2000) and some insect cuticles (Arakane et al. 2016). The pro-phenoloxidase (proPO) activates PO, resulting in the synthesis of melanin which in turn activates innate immune responses. The proPO system itself is activated by tyrosine and phenylalanine, which is synthesized by both the coral host as well as zooxanthellae (Fitzgerald and Szmant 1997). To reduce the impact of toxic by-products, such as reactive oxygen (RO) and other free radicals, antioxidant enzymes are up-regulated to offset self-harm (Gardner et al. 2016). However, the source and presence of such antioxidant enzymes and non-enzymatic antioxidants in coral mucus are not well studied (Vilas Bhagwat et al. 2023). Some enzymes, such as superoxide dismutase and glutathione peroxidase, have been reported as stress markers in corals (Gardner et al. 2016). Elevated PO activity triggers melanin production, serving as a signal for corals to react to pathogens, and concurrently functioning as a potent antioxidant against UV radiation (Palmer et al. 2011). The antioxidant capabilities of mucus against oxidative stress may determine the survival of the corals against bleaching and other stressors.

### **Antimicrobial activity of mucus and mucus-associated microbes**

Coral mucus from different habitats is composed of unique components that alter the presence of specific microbes on the surface of different coral species (Guppy and Bythell 2006; McShane et

al. 2021). The natural flora of healthy coral is presumably beneficial and any shift in this bacterial population may be an early indicator of coral stress, such as disease (Nithyanand et al. 2011). Therefore, the SML of coral is a selective media determined by host-microbe-environment interactions, where the commensal microbiota act against potential pathogens (Ritchie 2006; Alagely et al. 2011; Krediet et al. 2013; Teplitski et al 2016; Irudayarajan et al. 2023). The microbial gene abundance in the coral mucus is known to be involved in stress resistance, virulence, chemotaxis and motility, utilization of FAs and lipids, sulfur and nitrogen metabolism, and secondary metabolism making the mucus associated bacteria a first line of defense (Bythell and Wild 2011; Irudayarajan et al. 2023). Thus, some microbial communities in the SML are described as producing defensive antimicrobial chemicals (Orland and Kushmaro 2009; Irudayarajan et al. 2023). The Caribbean coral *Acropora palmata* demonstrated with antibacterial inhibition against the pathogen that causes white pox disease (Nissimov et al. 2009). Similarly, antibacterial activity was found in many cultivable bacteria of the coral mucus of six scleractinian corals from the Gulf of Eilat (Orland and Kushmaro 2009). Varied bacterial genera of both tissue and mucus of *A. digitifera*, including *Vibrio* sp. and *Bacillus* sp. have been elucidated for their antibacterial activity. *Bacillus* sp. synthesizes a significant peptide group of antibiotics including gramicidin, bacitracin and polymyxin B (Wiese et al. 2009). Thus, the highly specific mucus bacterial community is crucial for first line of defense of corals against diseases and other environmental stress factors (Kemp et al. 2015). Moreover, specific bacteriophages of the coral SML have been described to treat various bacterial infections as phage therapy (Nguyen et al. 2015; van Oppen 2019). The SML antimicrobial activity exhibited species specificity, while the mucus associated beneficial bacteria, such as *Pseudoalteromonas* sp. and *Roseobacter* sp., have been described for their antibacterial activity against a number of coral pathogens, including *V. shiloi* (Ortega and Thome 2018). As a result, beneficial microorganisms of corals (BMCs) are used as probiotic drugs in order to increase coral resistance to environmental stress, and the rotifers, *Brachionus plicatilis* are employed as carriers to impart BMCs into the coral *Pocillopora damicornis* (Assis et al. 2020).

The biochemicals present in the surface mucus of *Acropora palmata* and *Pseudodiploria strigosa* are responsible for regulating the coral-associated microbial community (Rivera-Ortega et al. 2018). Studies have shown that the mucus of *A. palmata* suppresses the growth of various bacterial strains, including *Bacillus subtilis*, *Staphylococcus aureus*, *Salmonella typhimurium*, and the white pox disease causative agent, *Serratia marcescens* (Ritchie 2006; Vilas Bhagwat et al. 2023). The mucus also exhibits antibacterial activity against both Gram-positive and Gram-negative bacteria (Ritchie 2006; Vilas Bhagwat et al. 2023). Similarly, healthy coral mucus obtained from *P. strigosa*

has been found to possess antimicrobial activity against *Serratia marcescens* and *Aurantimonas* sp. pathogens (Ortega and Thome 2018). In addition to its antimicrobial properties, *Galaxea fascicularis* mucus has been found to have potential health benefits for humans, such as apoptotic and anticancer activity, DNase activity, and activity against Topoisomerase I and II enzymes (Ding et al. 1999; Khalesi 2015). Homarine, an organic compound with antibiotic potential, has been documented in corals for its ability to potentially inhibit the growth of a range of bacteria and fouling diatoms (Maldonado et al. 2016). Fish mucus contains a variety of components, including disulfides, glycoconjugates, antimicrobial peptides (AMPs), lysozyme, cathepsin B, complement, alkaline phosphatase, C-reactive protein, transferrin, and others, which act as effective protection (Meyer et al. 2005; Subramanian et al. 2007; Zhao et al. 2008). AMPs in fish mucus are effective antimicrobial agents against human pathogens, including algae, bacteria, fungi, viruses, or parasites (Komatsu et al. 2009; Fernandes et al. 2010). Although the antimicrobial properties of coral mucus and associated bacteria have been mentioned in numerous papers, the specific components responsible for these properties are not well documented (Wright et al. 2019; Wijayanti et al. 2020), and further research is needed to better understand the immunological properties of coral mucus and the mechanisms by which it provides protection.

### **Mucus as a sunscreen and effective antioxidant under heat stress**

Mycosporine-like amino acids (MAAs) are responsible for shielding corals from the harmful effects of solar ultraviolet (UV) radiation by absorbing approximately 10 % of UV radiation. This mechanism is critical for the survival of corals, as exposure to high levels of UV radiation can cause damage to coral DNA, resulting in a range of negative impacts on coral health and the overall health of the surrounding ecosystem. By providing this important protection against UV radiation, MAAs play a key role in maintaining the health and resilience of coral populations (Teai et al. 1998; Kageyama and Waditee-Sirisattha 2019). Mycosporine-glycine, mycosporine-2glycine, palythine, mycosporine-methylamine-serine, and mycosporine-methylamine-threonine, which were characterized from *Lobophyllia*, *Acropora*, *Fungia*, and *Pocillopora* species, have all been reported from coral mucus (Teai et al. 1998). In addition to their ability to combat oxidative stress, MAAs can protect many marine organisms from the harmful effects of light (Shick and Dunlap 2002). Similarly, mycosporine-glycine showed an additional antioxidant property in coral mucus along with sunscreen, probably protecting endosymbionts from oxidative damage produced through photosynthesis (Teai et al. 1998). Biologically harmful UV rays between 310 and 360 nm are absorbed by MAAs. Endosymbionts and the coral hosts are both protected from the harmful effects of UV radiation by mucus MAAs. It was reported that the UV-absorbing molecules in the coral mucus of *Fungia fungites*

reduced with depth (Drollet et al. 1993). The reduction in UV radiation, light, and photosynthesis may be attributed to increased depth, as water absorbs and scatters UV radiation more effectively with increasing depth, leading to a decrease in UV radiation intensity.

Mycosporine-like amino acid concentrations vary depending on coral species and habitat. They have been found in both host tissues as well as in zooxanthellae (Shick and Dunlap 2002). MAAs are thermally stable, making them useful for scavenging reactive oxygen species under heat stress (Banaszak and Lesser 2009). MAAs play a role in cellular osmoregulation during salinity stress or in a high-salinity environment (Oren and Gunde-Cimerman 2007). The combination of mycosporine and glycine, the most common MAAs in coral hosts protects the coral symbiosis from oxidative damage (Banaszak et al. 2006). MAA also serves as a nitrogen source within the cell, which is required for growth and proliferation (Oren and Gunde-Cimerman 2007).

### **Mucus as a source of nutrients for marine organisms**

Mucus produced by coral is a valuable energy source with high glucose content. As glucose is readily available, it is a highly preferred energy source for microbes and other organisms (Ravindran et al. 2023). Thus, coral mucus is an important component of marine ecosystems, providing a rich source of energy for a wide group of organisms (Wild et al. 2010) (Table 2). Mucus energy levels are far greater than benthic photosynthetic algae. Further, a few soft corals have also been shown to feed on the mucus of other corals (Coffroth 1984). Similarly, fishes such as blue sprat (Eckes et al. 2015), damselfishes (Coles and Strathmann 1973; McMohan et al. 2016) and other benthic organisms such as an Acoelomorph worm (*Waminoa* sp.), bivalves (Shafir and Loya 1983), shrimp (*Coralliocaris superba*) (Horka et al. 2016), coral crabs (*Trapezia* sp.) (Shmuel et al. 2022) and some planktonic organisms, such as crown-of-thorns starfish larvae (Nakajima et al. 2016), copepods (*Acartia negligence*) (Richman 1975), and mysids, (*Mysidium integrum*) (Gottfried and Roman 1983) also exhibit feeding on coral mucus. Coral mucus was found to be the best nutritional source with an C:N ratio of mucus fluid of 6.9 to 13.7 and mucus sheet of 4.8 to 5.9 (Coffroth 1990) (Table 2). Since mucus sheets that stay on the coral surface can trap organic materials, their nutritional value is higher than reef benthic algae (~1) and reef algal detritus (~1) (Wilson et al. 2003). When compared with the other forms of energy sources, reef fish feces contribute to a nutritional value of ~10 (Bailey and Robertson 1982), phytoplankton ~3.5 (Renaud et al. 1999), and zooplankton ~5.5 (Ventura 2006). Furthermore, coral mucus acts as a vehicle for nutrients to reach the mouth (Goldberg 2018). Small reef fish and copepods utilize wax esters and triglycerides as energy sources, both of which are major components of mucus (Ho 2017). Thus, coral mucus helps in the fulfillment of energy requirements as well as the cycling of nutrients (Wild et al. 2004).

## **Role of mucus in reef ecosystem nutrient cycling**

When coral mucus is released into the water column, it is rapidly colonized by microorganisms that break down the organic compounds and release nutrients such as nitrogen ( $N_2$ ) and phosphorus (Van Oppen et al. 2019). These nutrients are then available for uptake by other organisms in the ecosystem, such as phytoplankton and zooplankton. Corals obtain  $N_2$  through predation and from the water column and excrete ammonia (Fig. 3) (Siboni et al. 2008). Further, Archaea use this ammonia, which plays an important role in ocean nitrification. During the daytime, when oxygen levels are high due to photosynthesis, certain types of Archaea convert ammonia into nitrite. Conversely, at night, when oxygen is low, different types of Archaea carry out denitrification to convert nitrite into nitrogen (Siboni et al. 2008 and 2012). Nitrogen-cycling microbes play a vital role in coral-algae symbiosis, especially under nutrient-rich or poor conditions for coral productivity (Radecker et al. 2015). Coral mucus contains high concentrations of dissolved organic nitrogen (DON), which can be readily taken up by bacteria for growth and metabolism. Some of these bacteria form symbiotic relationships with coral polyps, where they fix atmospheric nitrogen ( $N_2$ ) (Diaz 2017). Other bacteria in the mucus help to recycle nitrogen within the holobiont by breaking down waste products and releasing ammonium ( $NH_4^+$ ) and nitrate ( $NO_3^-$ ) into the surrounding water. These forms of nitrogen can then be taken up by other organisms in the reef ecosystem (Wild et al. 2004).

Up to 80% of mucus released by corals becomes dissolved organic carbon (DOC) (Wild et al. 2004). Sponges transfer dissolved organic matter (DOM) including DOC and DON to the next trophic level (Rix et al. 2017). Bacteria utilize mucus and break down organic compounds and release carbon dioxide ( $CO_2$ ) through respiration. This  $CO_2$  can then be taken up by photosynthetic organisms, such as algae and seagrasses, which use it to produce organic carbon through photosynthesis (Haas et al. 2016). This process helps to maintain the balance of carbon in the reef ecosystem. Coral mucus also contains significant amounts of dissolved organic phosphorus (DOP), which can be taken up by bacteria and recycled within the holobiont (Nakajima et al. 2015). Some of these bacteria are capable of breaking down complex organic phosphorus compounds, such as phospholipids, to release inorganic phosphorus ( $PO_4^{3-}$ ) into the surrounding water (Bourne et al. 2013). This inorganic phosphorus can then be taken up by algae and other organisms in the reef ecosystem. Mucus functions as nutrient trapping, thus, higher ammonia, nitrate, and phosphate in mucus than in surrounding water suggest its role in nutrient adsorption. In addition, coral mucus transports nutrients into reef sediment (Wild et al. 2005). Overall, coral mucus plays a crucial role in maintaining the health and productivity of reef ecosystems by facilitating nutrient cycling and supporting the diverse array of organisms that inhabit these ecosystems. Recycling these nutrients within the holobiont and the surrounding water helps to

maintain a balance of nutrients and supports the ecosystem's productivity. Additionally, the role of coral mucus in nutrient cycling highlights the importance of conserving coral reefs and protecting them from human activities that may disrupt this delicate balance.

## **Conclusion**

The coral surface mucus serves as an interface between the coral epithelium and the seawater environment. It provides a primary line of defense and protects coral surfaces from a variety of physiochemical and microbial invasions. It acts as a high-energy substrate for bacterial heterotrophic activity. As both host and endosymbiont provide mucus constituents, this symbiotic mucus synthesis is different from many other marine organisms. The exact mechanism of mucus synthesis, secretion, and sources of other important components present in mucus is unknown, and little information is available about the properties and rheological changes of mucus, which are not fully demonstrated. Only certain coral species have been studied for their mucus components, such as carbohydrates, fatty acids, amino acids, proteins, antioxidants, and antimicrobial compounds, but even among those, not all types have been thoroughly explored. Also, mucus defense properties, including antimicrobial peptides, antioxidants, and UV protectants (e.g., Mycosporine-like amino acids) and beneficial microbes, not well described for their roles in supporting coral health in various stress conditions. Coral mucus serves multiple ecological roles with their associated microorganisms and the surrounding environment that need to be studied to understand reef ecosystem functioning. Thus, the present review aids in understanding the role of coral mucus in coral reef ecosystems; however, more detailed studies of coral mucus components and their roles in the coral reef ecosystem by developing appropriate biochemical and molecular approaches may provide insights into protecting and restoring these valuable ecosystems.

## **Acknowledgments**

The authors thank the Director, CSIR-National Institute of Oceanography for the facilities. Thanks, Ministry of Environment, Forest and climate change (MoEFCC), Govt. of India, New Delhi and Forest Department, Principal Chief Conservator of Forests (PCCF) and Chief wildlife warden, Chennai-15, Tamil Nadu and Wildlife Warden, Gulf of Mannar Marine National Park, Tamil Nadu, India, for the entry and as well as for collecting the samples. his is NIO contribution #...

## **Funding**

This study was supported by the Department of Biotechnology (DBT), (Grant No. BT/PR15162/AAQ/3/752/2015) India to Dr. C.R..PVB was supported by CSIR (UGC) (Grant No.20/12/2015(ii)EU-V).. .

## Contributions

PVB-wrote the original draft, performed data analysis. CR- Conceptualization, fund acquisition, writing and editing the original draft, performed data analysis. LI- wrote the original draft, performed data analysis. All authors revised and approved the final manuscript.

## Corresponding author

Correspondence to Chinnarajan Ravindran – Email:cravindran@nio.org

## Ethics Declaration

## Conflicts of Interests

Authors have no conflict of interests.

**Ethical standards-** Not applicable

## References

- Alagely A, Krediet CJ, Ritchie KB, Teplitski M (2011) Signaling-mediated cross-talk modulates swarming and biofilm formation in a coral pathogen *Serratia marcescens*. *The ISME journal*. 5(10):1609-20.
- Ambort D, Johansson ME, Gustafsson JK, Ermund A, Hansson GC (2012) Perspectives on mucus properties and formation-lessons from the biochemical world. *Cold Spring Harbor Perspectives in Medicine* 2(11):014159.
- Andrianasolo EH, Haramaty L, Vardi A, White E, Lutz R, Falkowski P (2008) Apoptosis-inducing galactolipids from a cultured marine diatom, *Phaeodactylum tricorutum*. *Journal of Natural Products* 71(7):1197–1201.
- Anithajothi R, Duraikannu K, Umagowsalya G, Ramakritinan CM (2014) The presence of biomarker enzymes of selected scleractinian corals of Palk Bay, southeast coast of India. *BioMed Research International* 2014, 684874.
- Arakane Y, Noh MY, Asano T, Kramer KJ (2016) Tyrosine metabolism for insect cuticle pigmentation and sclerotization. *Extracellular composite matrices in arthropods*, pp. 165–220
- Assis JM, Abreu F, Villela HM, Barno A, Valle RF, Vieira R, Taveira I, Duarte G, Bourne DG, Høj L, Peixoto RS (2020) Delivering beneficial microorganisms for corals: rotifers as carriers of probiotic bacteria. *Frontiers in Microbiology*. 15(11):608506.
- Bachok Z, Mfilinge P, Tsuchiya M (2006) Characterization of fatty acid composition in healthy and bleached corals from Okinawa, Japan. *Coral Reefs* 25(4):545–554.
- Bailey TG, Robertson DR (1982) Organic and caloric levels of fish feces relative to its consumption by coprophagous reef fishes. *Marine Biology*, 69(1):45–50.

- Baker LJ, Reich HG, Kitchen SA, Grace Klings J, Koch HR, Baums IB, Muller EM, Thurber RV (2022) The coral symbiont *Candidatus Aquarickettsia* is variably abundant in threatened Caribbean acroporids and transmitted horizontally. *The ISME Journal*, 16(2):400-411.
- Bakshani CR, Morales-Garcia AL, Althaus M, Wilcox MD, Pearson JP, Bythell JC, Burgess JG (2018) Evolutionary conservation of the antimicrobial function of mucus: A first defense against infection. *npj Biofilms and Microbiomes*, 4(1):14.
- Banaszak AT, Barba Santos MG, LaJeunesse TC, Lesser MP (2006) The distribution of mycosporine-like amino acids (MAAs) and the phylogenetic identity of symbiotic dinoflagellates in cnidarian hosts from the Mexican Caribbean. *Journal of Experimental Marine Biology and Ecology*, 337(2):131–146.
- Banaszak AT, Lesser MP (2009) Effects of solar ultraviolet radiation on coral reef organisms. *Photochemical and Photobiological Sciences*, 8(9):1276–1294.
- Benavides M, Bednarz VN, Ferrier-Pagès C (2017) Diazotrophs: Overlooked key players within the coral symbiosis and tropical reef ecosystems? *Frontiers in Marine Science*, 4:10.
- Bessell-Browne P, Fisher R, Duckworth, A, Jones R (2017) Mucous sheet production in *Porites*: An effective bioindicator of sediment related pressures. *Ecological Indicators*, 77:276–285.
- Binienda ZK, Sarkar S, Silva-Ramirez S, Gonzalez C (2013) Role of free fatty acids in physiological conditions and mitochondrial dysfunction. *Food and Nutrition Sciences*, 04(9):6–15.
- Bishop DG, Kenrick JR (1980) Fatty acid composition of symbiotic Zooxanthellae in relation to their hosts. *Lipids*, 15(10):799.
- Bourne DG, Dennis PG, Uthicke S, Soo RM, Tyson GW, Webster N (2013) Coral reef invertebrate microbiomes correlate with the presence of photosymbionts. *The ISME journal*, 7(7):1452-1458.
- Brown BE, Bythell JC (2005) Perspectives on mucus secretion in reef corals. *Marine Ecology Progress Series*, 296:291–309.
- Bruckner AW (2015) White syndromes of western Atlantic reef-building corals. *Diseases of Coral*, 316–332.
- Burriesci MS, Raab TK, Pringle JR (2012) Evidence that glucose is the major transferred metabolite in dinoflagellate–cnidarian symbiosis. *Journal of Experimental Biology*, 215(19):3467-3477.
- Bythell JC, Wild C (2011) Biology and ecology of coral mucus release. *Journal of Experimental Marine Biology and Ecology*, 408(1–2):88–93.
- Coddeville B, Maes E, Ferrier-Pages C, Guerardel Y (2011) Glycan profiling of gel forming mucus layer from the scleractinian symbiotic coral *Oculina arbuscula*. *Biomacromolecules*, 12(6):2064–2073.
- Coffroth MA (1984) Ingestion and incorporation of coral mucus aggregates by a gorgonian soft coral. *Marine Ecology Progress Series*, 17(2):193–199.



- Coffroth MA (1990) Mucous sheet formation on poritid corals: An evaluation of coral mucus as a nutrient source on reefs. *Marine Biology*, 105(1):39–49.
- Coles SL, Strathmann R (1973) Observations on coral mucus “flocs” and their potential trophic significance 1. *Limnology and Oceanography*, 18(4):673–678.
- Corfield AP (2015) Mucins: A biologically relevant glycan barrier in mucosal protection. *Biochimica et Biophysica Acta*, 1850(1):236–252.
- Crossland CJ (1987) In situ release of mucus and DOC-lipid from the corals *Acropora variabilis* and *Stylophora pistillata* in different light regimes. *Coral Reefs*, 6(1):35–42.
- Crossland CJ (2021) Hermatypic coral, *Acropora* cf. *acuminata*. In *Endosymbiosis and cell biology: A synthesis of recent research. Proceedings of the International Colloquium on Endosymbiosis and Cell Research, Tübingen, April 1980 (Vol. 1, p. 163)*. Walter de Gruyter GmbH & Co KG.
- Crossland CJ, Barnes DJ, Borowitzka MA (1980) Diurnal lipid and mucus production in the staghorn coral *Acropora acuminata*. *Marine Biology*, 60(2–3):81–90.
- Dailah HG (2022) Therapeutic potential of small molecules targeting oxidative stress in the treatment of chronic obstructive pulmonary disease (COPD): A comprehensive review. *Molecules*, 27(17):5542.
- Daumas R, Thomassin BA (1977) Protein fractions in coral and zoantharian mucus, possible evolution in coral reef environment. In *Proceedings of the (3rd) Int. Coral Reef Symp. (Miami) 1* (pp. 517–525).
- Denny MW (1989) Invertebrate mucous secretions – Functional alternatives to vertebrate paradigms. *Symposia of the Society for Experimental Biology*, 43:337–366.
- Diaz L (2017) Response of the coral associated nitrogen fixing bacteria toward elevated water temperature. *Journal of Water Resources and Ocean Science*, 6(6):98–109.
- Ding JL, Fung FMY, Ng GWS, Chou LM (1999) Novel Bioactivities from a Coral, *Galaxea fascicularis*: DNase-like Activity and Apoptotic Activity Against a Multiple-Drug-Resistant Leukemia Cell Line. *Marine Biotechnology*, 1(4):328–336.
- Downs CA, Fauth JE, Halas JC, Dustan P, Bemiss J, Woodley CM (2002) Oxidative stress and seasonal coral bleaching. *Free Radical Biology and Medicine*, 33(4):533–543.
- Drollet JH, Teai T, Faucon M, Martin PMV (1997) Field study of compensatory changes in UV-absorbing compounds in the mucus of the solitary coral *Fungia repanda* (Scleractinia: Fungiidae) in relation to solar UV radiation, sea-water temperature, and other coincident physico-chemical parameters. *Marine and Freshwater Research*, 48(4):329–333.
- Ducklow HW, Mitchell R (1979) Bacterial populations and adaptations in the mucus layers on living corals 1. *Limnology and Oceanography*, 24(4):715–725.
- Eckes M, Dove S, Siebeck UE, Grutter AS (2015) Fish mucus versus parasitic gnathiid isopods as sources of energy and sunscreens for a cleaner fish. *Coral Reefs*, 34(3):823–833.

- Fang JHS (2015) Community analysis of coral mucus-associated bacteria and impact of temperature and CO<sub>2</sub> changes on them (Doctoral dissertation. Swinburne University of Technology (Sarawak campus).
- Fernandes-Alnemri T, Yu JW, Juliana C, Solorzano L, Kang S, Wu J, Datta P, McCormick M, Huang L, McDermott E, Eisenlohr L, Landel CP, Alnemri ES (2010) The AIM2 inflammasome is critical for innate immunity to *Francisella tularensis*. *Nature Immunology*, 11(5), 385–393.
- Fitzgerald LM, Szmant AM (1997) Biosynthesis of “essential” amino acids by scleractinian corals. *Biochemical Journal*, 322(1), 213–221.
- Gabay Y (2018) The cellular and physiological basis of host-symbiont specificity in a model cnidarian-dinoflagellate symbiosis (Doctoral dissertation, Open Access Te Herenga Waka-Victoria University of Wellington).
- Galli P, Montano S, Seveso D, Maggioni D (2021) Coral reef biodiversity of the Maldives. Atolls of the Maldives. *Nissology and Geography*. Rowman and Littlefield, Lanham, 196-212.
- Gardner SG, Nielsen DA, Laczka O, Shimmon R, Beltran VH, Ralph PJ, Petrou K (2016) Dimethylsulfonylpropionate, superoxide dismutase and glutathione as stress response indicators in three corals under short-term hyposalinity stress. *Proceedings. Biological Sciences*, 283(1824):20152418.
- Glasl B, Herndl GJ, Frade PR (2016) The microbiome of coral surface mucus has a key role in mediating holobiont health and survival upon disturbance. *ISME Journal*, 10(9):2280–2292.
- Goldberg WM (2018) Coral food, feeding, nutrition, and secretion: A review. In *Marine organisms as model systems in biology and medicine* (pp. 377–421).
- Gottfried M, Roman MR (1983) Ingestion and incorporation of coral-mucus detritus by reef zooplankton. *Marine Biology*, 72(3):211–218.
- Grottoli AG, Rodrigues LJ, Juarez C (2004) Lipids and stable carbon isotopes in two species of Hawaiian corals. *Porites compressa* and *Montipora verucosa*, following a bleaching event. *Marine Biology*, 145:621–631.
- Guarnieri MC, de Albuquerque Modesto JC, Pérez CD, Ottaiano TF, Ferreira RDS, Batista FP, de Brito MV, Campos IHMP, Oliva MLV (2018) Zoanthid mucus as new source of useful biologically active proteins. *Toxicon*, 143:96–107.
- Guppy R, Bythell JC (2006) Environmental effects on bacterial diversity in the surface mucus layer of the reef coral *Montastraea faveolata*. *Marine Ecology Progress Series*, 328:133-142.
- Guppy R, Brown B, Bythell JC (2019) Preserving the viscous coral surface mucus layer using low-acid glycol methacrylate (GMA) resin. *Coral Reefs*, 38(3):521-526.
- Haas AF, Fairouz MFM, Kelly LW, Nelson CE, Dinsdale EA, Edwards RA et al (2016) Global microbialization of coral reefs. *Nature Microbiology*, 1(7):16042.
- Hadaidi G, Gegner HM, Ziegler M, Voolstra CR (2019) Carbohydrate composition of mucus from scleractinian corals from the central Red Sea. *Coral Reefs*, 38(1):21–27.

- Hadaidi GA (2018) Coral-associated bacterial community dynamics in healthy, bleached, and disease states, (Doctoral Dissertation), KAUST Research Repository. <https://doi.org/10.25781/KAUST-0UN42>.
- Hafezi M, Giffin AL, Alipour M, Sahin O, Stewart RA (2020) Mapping long-term coral reef ecosystems regime shifts: A small island developing state case study. *Science of the Total Environment*, 716:137024.
- Hill DB, Button B, Rubinstein M, Boucher RC (2022) Physiology and pathophysiology of human airway mucus. *Physiological Reviews*, 102(4):1757–1836.
- Hillyer KE, Dias DA, Lutz A, Wilkinson SP, Roessner U, Davy SK (2017) Metabolite profiling of symbiont and host during thermal stress and bleaching in the coral *Acropora aspera*. *Coral Reefs*, 36:105-118.
- Ho TK (2017) Dredging land reclamation causing mucus development in massive spherical corals in the Spratly Islands, South China Sea: The effects on China's fishing industry. Poster session presented at Virginia Commonwealth University Poster Symposium for Undergraduate Research and Creativity, Richmond, VA.
- Horka I, De Grave S, Franssen CH, Petrusek A, Ďuriš Z (2016) Multiple host switching events shape the evolution of symbiotic palaemonid shrimps (Crustacea: Decapoda). *Scientific Reports*, 6(1):26486.
- Horricks RA, Herbinger CM, Lillie BN, Taylor P, Lumsden JS (2019) Differential protein abundance during the first month of regeneration of the Caribbean star coral *Montastraea cavernosa*. *Coral Reefs*, 38(1):45–61.
- Iluz D, Dubinsky Z (2015) Coral photobiology: New light on old views. *Zoology*, 118(2):71–78.
- Irudayarajan L, Ravindran C, Raveendran HP (2023) Antimicrobial activity of coral-associated beneficial bacteria against coral disease-causing microbial pathogens. *J Basic Microbiol*. doi: 10.1002/jobm.202300338. (Epub ahead of print).
- Jatkar AA, Brown BE, Bythell JC, Guppy R, Morris NJ, Pearson JP (2010a) Coral mucus: the properties of its constituent mucins. *Biomacromolecules*, 11(4):883-888.
- Jatkar AA, Brown BE, Bythell JC, Guppy R, Morris NJ, Pearson JP (2010b) Measuring mucus thickness in reef corals using a technique devised for vertebrate applications. *Marine biology*, 157:261-267.
- Jeffries HP (1972) Fatty acid ecology of a tidal Marsh 1. *Limnology and Oceanography*, 17(3):433–440.
- Kageyama H, Waditee-Sirisattha R (2019) Antioxidative, anti-inflammatory, and anti-aging properties of mycosporine-like amino acids: Molecular and cellular mechanisms in the protection of skin-aging. *Marine Drugs*, 17(4):222.
- Kemp DW, Rivers AR, Kemp KM, Lipp EK, Porter JW, Wares JP (2015) Spatial homogeneity of bacterial communities associated with the surface mucus layer of the reef-building coral *Acropora palmata*. *PLOS ONE*, 10(12):0143790.

- Khalesi MK (2015) Corals. Springer Handbook of Marine Biotechnology, 179-217.
- Komatsu K, Tsutsui S, Hino K, Araki K, Yoshiura Y, Yamamoto A, Nakamura O, Watanabe T (2009) Expression profiles of cytokines released in intestinal epithelial cells of the rainbow trout, *Oncorhynchus mykiss*, in response to bacterial infection. *Developmental and Comparative Immunology*, 33(4):499–506.
- Krediet CJ, Ritchie KB, Alagely A, Teplitski M (2013) Members of native coral microbiota inhibit glycosidases and thwart colonization of coral mucus by an opportunistic pathogen. *The ISME journal*, 7(5):980-990.
- Krediet CJ, Ritchie KB, Alagely A, Teplitski M (2013) Members of native coral microbiota inhibit glycosidases and thwart colonization of coral mucus by an opportunistic pathogen. *The ISME journal*. 7(5):980-90.
- Krediet CJ, Ritchie KB, Cohen M, Lipp EK, Sutherland KP, Teplitski M (2009) Utilization of mucus from the coral *Acropora palmata* by the pathogen *Serratia marcescens* and by environmental and coral commensal bacteria. *Applied and Environmental Microbiology*, 75(12):3851–3858.
- Krieger-Liszkay A (2005) Singlet oxygen production in photosynthesis. *Journal of Experimental Botany*, 56(411):337–346.
- Krueger T, Bodin J, Horwitz N, Loussert-Fonta C, Sakr A, Escrig S, Fine M, Meibom A (2018) Temperature and feeding induce tissue level changes in autotrophic and heterotrophic nutrient allocation in the coral symbiosis—A NanoSIMS study. *Scientific Reports*, 8(1):12710.
- Krupp DA (1982) The composition of the mucus from the mushroom coral *Fungia scutaria*. In *Proceedings of the 4th Int. Coral Reef Symp.*, 2 (9–73).
- Kuhl M, Cohen Y, Dalsgaard T, Jørgensen BB, Revsbech NP (1995) Microenvironment and photosynthesis of Zooxanthellae in scleractinian corals studied with microsensors for O<sub>2</sub>, pH and light. *Marine Ecology Progress Series*, 117:159–172.
- Kurihara H, Ikeda N, Umezawa Y (2018) Diurnal and seasonal variation of particle and dissolved organic matter release by the coral *Acropora tenuis*. *PeerJ*, 6:5728.
- LaJeunesse TC, Smith R, Walther M, Pinzón J, Pettay DT, McGinley M, Aschaffenburg M, Medina-Rosas P, Cupul-Magaña AL, Pérez AL, Reyes-Bonilla H, Warner ME (2010) Host–symbiont recombination versus natural selection in the response of coral–dinoflagellate symbioses to environmental disturbance. *Proceedings of the Royal Society B*, 277(1696):2925–2934.
- Lee ST, Dav SK, Tang SL, Kench PS (2016) Mucus sugar content shapes the bacterial community structure in thermally stressed *Acropora muricata*. *Frontiers in microbiology*, 7:371.
- Lewis RW (1976) Mucus globule membrane: An hypothesis concerning its role in determining the viscosity of mucus. *Journal of Theoretical Biology*, 61(1):21–25.
- Lin C, Wang LH, Fan TY, Kuo FW (2012) Lipid content and composition during the oocyte development of two gorgonian coral species in relation to low temperature preservation. *PLoS One*, 7(7):38689.

- Maldonado A, Johnson A, Gochfeld D, Slattery M, Ostrander GK, Bingham JP, Schlenk D (2016) Hard Coral (*Porites lobata*) extracts and homarine on Cytochrome P450 expression in Hawaiian butterflyfishes with different feeding strategies. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 179:57-63.
- Marshall AT, Wright OP (1993) Confocal laser scanning light microscopy of the extra-thecal epithelia of undecalcified scleractinian corals. *Cell and Tissue Research*, 272(3):533–543.
- Mascarelli PE, Bunkley-Williams L (1999) An experimental field evaluation of healing in damaged, unbleached and artificially bleached star coral, *Montastraea annularis*. *Bulletin of Marine Science*, 65(2):577–586.
- McMahon KW, Thorrold SR, Houghton LA, Berumen ML (2016) Tracing carbon flow through coral reef food webs using a compound-specific stable isotope approach. *Oecologia*, 180(3):809–821.
- McShane A, Bath J, Jaramillo AM, Ridley C, Walsh AA, Evans CM, Thornton DJ, Ribbeck K (2021) Mucus. *Current Biology*, 31(15):938–945.
- Means JC, Sigleo AC (1986) Contribution of coral reef mucus to the colloidal organic pool in the vicinity of Discovery Bay, Jamaica, WI. *Bulletin of Marine Science*, 39(1):110–118.
- Meikle P, Richards GN, Yellowlees D (1988) Structural investigations on the mucus from six species of coral. *Marine Biology*, 99(2):187–193.
- Meyer W, Seegers U, Schnapper A, Neuhaus H, Himstedt W, Toepfer-Petersen E (2007) Possible antimicrobial defense by free sugars on the epidermal surface of aquatic vertebrates. *Aquatic Biology*, 1(2):167–175.
- Mock A, Renwrantz L (1987) Lectin activity and a protease inhibitor in mucus from the skin of *Branchiostoma lanceolatum*. *Journal of Invertebrate Pathology*, 49(2):221–222.
- Muscatine L, Falkowski PG, Porter JW, Dubinsky Z (1984) Fate of photosynthetic fixed carbon in light- and shade-adapted colonies of the symbiotic coral *Stylophora pistillata*. *Proceedings of the Royal Society of London Series B*, 222(1227):181–202.
- Nakajima R, Nakatomi N, Kurihara H, Fox MD, Smith JE, Okaji K (2016) Crown-of-thorns starfish larvae can feed on organic matter released from corals. *Diversity*, 8(4):18.
- Nakajima R, Tanaka Y, Yoshida T, Fujisawa T, Nakayama A, Fuchinoue Y, Othman BHR, Toda T (2015) High inorganic phosphate concentration in coral mucus and its utilization by heterotrophic bacteria in a Malaysian coral reef. *Marine Ecology*, 36(3):835–841.
- Nguyen-Kim H, Bettarel Y, Bouvier T, Bouvier C, Doan-Nhu H, Nguyen-Ngoc L, Nguyen-Thanh T, Tran-Quang H, Brune J (2015) Coral mucus is a hot spot for viral infections. *Applied and Environmental Microbiology*, 81(17):5773–5783.
- Nissimov J, Rosenberg E, Munn CB (2009) Antimicrobial properties of resident coral mucus bacteria of *Oculina Patagonica*. *FEMS Microbiology Letters*, 292(2):210–215.

- Nithyanand P, Indhumathi T, Ravi AV, Pandian SK (2011) Culture independent characterization of bacteria associated with the mucus of the coral *Acropora digitifera* from the Gulf of Mannar. *World Journal of Microbiology and Biotechnology*, 27(6):1399–1406.
- Oku H, Yamashiro H, Onaga K (2003) Lipid biosynthesis from [14C]-glucose in the coral *Montipora digitata*. *Fisheries Science*, 69(3):625–631.
- Oren A, Gunde-Cimerman N (2007) Mycosporines and mycosporine-like amino acids: UV protectants or multipurpose secondary metabolites? *FEMS Microbiology Letters*, 269(1):1–10.
- Palmer CV, Traylor-Knowles NG, Willis BL, Bythell JC (2011) Corals use similar immune cells and wound-healing processes as those of higher organisms. *PLOS ONE*, 6(8):23992.
- Papina M, Meziane T, van Woesik R (2003) Symbiotic *Zooxanthellae* provide the host-coral *Montipora digitata* with polyunsaturated fatty acids. *Comparative Biochemistry and Physiology. Part B, Biochemistry and Molecular Biology*, 135(3):533–537.
- Pasaribu B, Weng LC, Lin IP, Camargo E, Tzen JT, Tsai CH, Ho SL, Lin MR, Wang LH, Chen CS, Jiang PL (2015) Morphological variability and distinct protein profiles of cultured and endosymbiotic *Zooxanthellae* cells isolated from *Exaiptasia Pulchella*. *Scientific Reports*, 5(1):15353.
- Patton JS, Abraham S, Benson AA (1977) Lipogenesis in the intact coral *Pocillopora capitata* and its isolated *Zooxanthellae*: Evidence for a light-driven carbon cycle between symbiont and host. *Marine Biology*, 44(3):235–247.
- Piggot AM, Fouke BW, Sivaguru M, Sanford RA, Gaskins HR (2009) Change in *Zooxanthellae* and mucocyte tissue density as an adaptive response to environmental stress by the coral, *Montastraea annularis*. *Marine Biology*, 156(11):2379–2389.
- Piludu M, Rayment SA, Liu B, Offner GD, Oppenheim FG, Troxler RF, Hand AR (2003) Electron microscopic immunogold localization of salivary mucins MG1 and MG2 in human submandibular and sublingual glands. *Journal of Histochemistry and Cytochemistry*, 51(1):69–79.
- Plass-Johnson JG, Cardini U, Van Hoytema N, Bayraktarov E, Burghardt I, Naumann MS, Wild C (2015) Coral bleaching. *Environmental Indicators*, 117–146.
- Que J (2015) The initial establishment and epithelial morphogenesis of the esophagus: A new model of tracheal–esophageal separation and transition of simple columnar into stratified squamous epithelium in the developing esophagus. *Wiley Interdisciplinary Reviews. Developmental Biology*, 4(4):419–430.
- Radecker N, Pogoreutz C, Voolstra CR, Wiedenmann J, Wild C (2015) Nitrogen cycling in corals: The key to understanding holobiont functioning? *Trends in Microbiology*, 23(8):490–497.
- Radice VZ, Brett MT, Fry B, Fox MD, Hoegh-Guldberg O, Dove SG (2019) Evaluating coral trophic strategies using fatty acid composition and indices. *PLOS ONE*, 14(9):0222327.
- Raina JB, Tapiolas D, Motti CA, Foret S, Seemann T, Tebben J, Willis BL, Bourne DG (2016) Isolation of an antimicrobial compound produced by bacteria associated with reef-building corals. *PeerJ*, 4:2275.

- Ravindran C, Lawrance I, Vasanth AJ (2022) Endosymbiotic *Symbiodinium* clades occurrence and influence on coral growth and resilience during stress. *Symbiosis* **86**, 261–272.
- Ravindran C, Bhagwat PV, Silveira KA, Shivaramu MS, Lele UP (2020) Characterization of coral associated ciliates and their interactions with disease lesion progression of Indian scleractinian corals. *Microbial Pathogenesis*, 149:104472.
- Ravindran C, Irudayarajan L, Raveendran HP (2023) Possible beneficial interactions of ciliated protozoans with coral health and resilience. *Applied and Environmental Microbiology*: e0121723.
- Renaud SM, Thinh LV, Parry DL (1999) The gross chemical composition and fatty acid composition of 18 species of tropical Australian microalgae for possible use in mariculture. *Aquaculture*, 170(2):147–159.
- Reshef L, Kore O, Loya Y, Zilber-Rosenberg I, Rosenberg E (2006) The coral probiotic hypothesis. *Environ. Microbiol.* 8:2068–2073
- Reynaud S, Ferrier-Pagès C (2019) Biology and ecophysiology of Mediterranean cold-water corals. In *Mediterranean cold-water corals: Past, present and future: Understanding the deep-sea realms of coral* (pp. 391–404).
- Richman S, Loya Y, Slobocckin LB (1975) The rate of mucus production by corals and its assimilation by the coral reef copepod *Acartia negligens* 1. *Limnology and Oceanography*, 20(6):918–923.
- Ritchie KB (2006) Regulation of microbial populations by coral surface mucus and mucus-associated bacteria. *Marine Ecology Progress Series*, 322:1–14.
- Rivera-Ortega J, Thomé PE (2018) Contrasting antibacterial capabilities of the surface mucus layer from three symbiotic cnidarians. *Frontiers in Marine Science*, 5:392.
- Rix L, de Goeij JM, van Oevelen D, Struck U, Al-Horani FA, Wild C, Naumann MS (2017) Differential recycling of coral and algal dissolved organic matter via the sponge loop. *Functional Ecology*, 31(3):778-789.
- Rosenberg, E (2021) Microbiomes. *Current Knowledge and Unanswered Questions*. Springer book.
- Salvatore S, Heuschkel R, Tomlin S, Davies SE, Edwards S, Walker-Smith J, ..., Murch SH (2000) A pilot study of N-acetyl glucosamine, a nutritional substrate for glycosaminoglycan synthesis, in paediatric chronic inflammatory bowel disease. *Alimentary pharmacology & therapeutics*, 14(12):1567-1579.
- Schottner S, Hoffmann F, Wild C, Rapp HT, Boetius A, Ramette A (2009) Inter-and intra-habitat bacterial diversity associated with cold-water corals. *The ISME journal*, 3(6):756-759.
- Shafir A, Loya A (1983) Consumption and assimilation of coral mucus by the burrowing mussel *Lithophaga lessepsiana*. In *Proceedings of the Int Conf Mar Sci. Red Sea*, 9 (pp. 135–140).
- Sharma D, Ravindran, C. 2020. Diseases and pathogens of marine invertebrate corals in Indian reefs. *J. of Invertebrate pathology*. 173: 107373.

- Shick JM, Dunlap WC, Pearse JS, Pearse VB (2002) Mycosporine-like amino acid content in four species of sea anemones in the genus *Anthopleura* reflects phylogenetic but not environmental or symbiotic relationships. *Biological Bulletin*, 203(3):315–330.
- Shmuel Y, Ziv Y, Rinkevich B (2022) Coral-inhabiting Trapezia crabs forage on demersal plankton. *Frontiers in Marine Science*, 9:1652.
- Shnit-Orland M, Kushmaro A (2009) Coral mucus-associated bacteria: A possible first line of defense. *FEMS Microbiology Ecology*, 67(3):371–380.
- Siboni N, Ben-Dov E, Sivan A, Kushmaro A (2008) Global distribution and diversity of coral associated Archaea and their possible role in the coral holobiont nitrogen cycle. *Environmental Microbiology*, 10(11):2979-2990.
- Siboni N, Ben-Dov E, Sivan A, Kushmaro A (2012) Geographic specific coral-associated ammonia-oxidizing archaea in the northern Gulf of Eilat (Red Sea). *Microbial ecology*, 64:18-24.
- Sikorskaya TV, Ermolenko EV, Efimova KV (2022) Lipids of Indo-Pacific gorgonian corals are modified under the influence of microbial associations. *Coral Reefs*, 41(2):277–291.
- Smith HA, Conlan JA, Pollock FJ, Wada N, Shore A, Hung JYH, Aeby GS, Willis BL, Francis DS, Bourne DG (2020) Energy depletion and opportunistic microbial colonisation in white syndrome lesions from corals across the Indo-Pacific. *Scientific Reports*, 10(1):19990.
- Stabili L (2019) The mucus of marine invertebrates: Cnidarians, polychaetes, and echinoderms as case studies. In *Enzymatic technologies for marine polysaccharides* (pp. 151–162). CRC Press.
- Stabili L, Schirosi R, Parisi MG, Piraino S, Cammarata M (2015) The mucus of *Actinia equina* (Anthozoa, Cnidaria): An unexplored resource for potential applicative purposes. *Marine Drugs*, 13(8):5276–5296.
- Stephens GC (1962) Uptake of organic material by aquatic invertebrates. I. Uptake of glucose by the solitary coral, *Fungia scutaria*. *Biological Bulletin*, 123(3):648–659.
- Subramanian S, MacKinnon SL, Ross NW (2007) A comparative study on innate immune parameters in the epidermal mucus of various fish species. *Comparative Biochemistry and Physiology. Part B, Biochemistry and Molecular Biology*, 148(3):256–263.
- Suggett DJ, Warner ME, Smith DJ, Davey P, Hennige S, Baker NR (2008) Photosynthesis and production of hydrogen peroxide by Zooxanthellae (pyrrhophyta) phylotypes with different thermal tolerances 1. *Journal of Phycology*, 44(4):948–956.
- Tchernov D, Gorbunov MY, De Vargas C, Narayan Yadav SN, Milligan AJ, Häggblom M, Falkowski PG (2004) Membrane lipids of symbiotic algae are diagnostic of sensitivity to thermal bleaching in corals. *Proceedings of the National Academy of Sciences of the United States of America*, 101(37):13531–13535.
- Teai T, Drollet JH, Bianchini JP, Cambon A, Martin PMV (1998) Occurrence of ultraviolet radiation-absorbing mycosporine-like amino acids in coral mucus and whole corals of French Polynesia. *Marine and Freshwater Research*, 49(2):127–132.



- Teplitski M, Krediet CJ, Meyer JL, Ritchie KB (2016). Microbial interactions on coral surfaces and within the coral holobiont. *The Cnidaria, Past, Present and Future: The World of Medusa and Her Sisters*, 331-346.
- Thompson JR, Rivera HE, Closek CJ, Medina M (2014) Microbes in the coral holobiont: Partners through evolution, development, and ecological interactions. *Frontiers in Cellular and Infection Microbiology*, 4:176.
- Thornton DJ, Sheehan JK (2004) From mucins to mucus: Toward a more coherent understanding of this essential barrier. *Proceedings of the American Thoracic Society*, 1(1):54–61.
- Tranter PRG, Nicholson DN, Kinchington D (1982) A description of spawning and post-gastrula development of the cool temperate coral, *Caryophyllia smithi*. *Journal of the Marine Biological Association of the United Kingdom*, 62(4):845–854.
- Treignier C, Grover R, Ferrier-Pagés C, Tolosa I (2008) Effect of light and feeding on the fatty acid and sterol composition of *Zooxanthellae* and host tissue isolated from the scleractinian coral *Turbinaria reniformis*. *Limnology and Oceanography*, 53(6):2702–2710.
- Uslu L, Sayin S, Naz M, Taskin E, Soyler O, Saygili I, Cetin Z, Dinler ZM, Isik O (2021) Proximate analysis and fatty acid profile of some brown macroalgae collected from the Northeastern Mediterranean coast. *Fresenius Environment Bulletin*, 30:9433–9437.
- Vacelet E, Thomassin BA (1991) Microbial utilization of coral mucus in long term in situ incubation over a coral reef. *Hydrobiologia*, 211(1):19–32.
- Van Oppen MJ, Blackall LL (2019) Coral microbiome dynamics, functions and design in a changing world. *Nature Reviews Microbiology*, 17(9):557-567.
- Vaughan EJ, Wilson SK, Howlett SJ, Parravicini V, Williams GJ, Graham NAJ (2021) Nitrogen enrichment in macroalgae following mass coral mortality. *Coral Reefs*, 40(3):767–776.
- Ventura M (2006) Linking biochemical and elemental composition in freshwater and marine crustacean zooplankton. *Marine Ecology Progress Series*, 327:233–246.
- Vilas Bhagwat P, Ravindran C, Irudayarajan L (2023) Characterization of the defense properties of healthy and diseased coral mucus. *J Invertebr Pathol*. 201:108001.
- Volkman JK, Johns RB, Gillan FT, Perry GJ, Bavor Jr HJ (1980) Microbial lipids of an intertidal sediment—I. Fatty acids and hydrocarbons. *Geochimica et Cosmochimica Acta* 44 (8):1133-1143.
- Weis VM (2008) Cellular mechanisms of Cnidarian bleaching: Stress causes the collapse of symbiosis. *Journal of Experimental Biology*, 211(19):3059–3066.
- Wiese J, Thiel V, Nagel K, Staufenberg T, Imhoff JF (2009) Diversity of antibiotic-active bacteria associated with the brown alga *Laminaria saccharina* from the Baltic Sea. *Marine Biotechnology*, 11(2):287–300.
- Wijayanti DP, Sabdono A, Dirgantara D, Widyananto PA, Sibero MT, Bhagooli R, Hidaka M (2020) Antibacterial activity of acroporid bacterial symbionts against White Patch Disease in

- Karimunjawa Archipelago, Indonesia. *The Egyptian Journal of Aquatic Research*, 46(2):187-193.
- Wild C, Huettel M, Klueter A, Kremb SG, Rasheed MY, Jorgensen BB, (2004) Coral mucus functions as an energy carrier and particle trap in the reef ecosystem. *Nature*, 428(6978):66-77.
- Wild C, Naumann M, Niggel W, Haas A (2010) Carbohydrate composition of mucus released by scleractinian warm- and cold-water reef corals. *Aquatic Biology*, 10(1):41–45.
- Wild C, Woyt H, Huettel M (2005) Influence of coral mucus on nutrient fluxes in carbonate sands. *Marine Ecology Progress Series*, 287:87–98.
- Wilson SK, Bellwood DR, Choat JH, Furnas MJ (2003) Detritus in the epilithic algal matrix and its use by coral reef fishes. *Oceanography and Marine Biology*, 41:279–310.
- Wright RM, Strader ME, Genuise HM, Matz M (2019) Effects of thermal stress on amount, composition, and antibacterial properties of coral mucus. *Peer J*, 7:6849.
- Younan F, Pearson J, Allen A, Venables C (1982) Changes in the structure of the mucous gel on the mucosal surface of the stomach in association with peptic ulcer disease. *Gastroenterology*, 82(5 Pt 1):827–831.
- Zetsche EM, Baussant T, Meysman FJ, Van Oevelen D (2016) Direct visualization of mucus production by the cold-water coral *Lophelia pertusa* with digital holographic microscopy. *PLOS ONE*, 11(2):0146766.
- Zhang S, Li G (2000) Presence of phenoloxidase and prophenoloxidase in the epidermal cells and the epidermis mucus of the lancelet *Branchiostoma belcheritsingtauense*. *Ophelia*, 52(3):207–212.
- Zhao X, Findly RC, Dickerson HW (2008) Cutaneous antibody-secreting cells and B cells in a teleost fish. *Developmental and Comparative Immunology*, 32(5):500-508.

## Figure legends

**Fig. 1 - Coral mucus and properties.** Coral mucus synthesis occurs in coral polyps from mucocyte cells. The mucus has a combination of nutrients provided by both the coral and endosymbiont. The released mucus acts as a first line of defense, with various properties against microbial pathogens, UV radiation and other stressors. that influence the health of the corals.

**Fig. 2 - Fatty acid contents of mucus.** Percent occurrence of polyunsaturated fatty acids (PUFA), monounsaturated fatty acids (MUFA), and saturated fatty acids (SAFA) in coral mucus (data calculated from the means of fatty acids reported in Sigleo (1986) and Ravindran et al. (2020)).

**Fig. 3 – Coral mucus in reef nutrient cycling.** Corals release mucus rich in dissolved organic matter (DOM) to the surrounding water, which is utilized by reef organisms whose respiration returns carbon dioxide and other nutrients required for photosynthesis by the endosymbiont zooxanthellae. Also, endosymbiont derived organic compounds are released into the mucus and nutrient cycling is activated through various microbes. (DOM- dissolved organic matter, DON- dissolved organic nitrogen, DIP- dissolved inorganic phosphate, DOC- dissolved organic carbon, DOP- dissolved organic phosphate, NH<sub>4</sub>- Ammonia, NO<sub>2</sub>- nitrite, N<sub>2</sub>- nitrogen).

**Table 1 Carbohydrate content of various coral species mucus.**

<b>S.no.</b>	<b>Carbohydrate</b>	<b>Coral species</b>	<b>Reference</b>
1.	Arabinose	<i>Acropora</i> sp., <i>Fungia</i> sp., <i>Pachyseris speciose</i> , <i>Fungia fungites</i> , <i>Sarcophyton</i> sp., <i>Lemnalina</i> sp., <i>Cespitularia</i> sp., <i>Porites lobata</i> ., <i>Pachyseris</i> sp., <i>Lophelia</i> sp.	Meikle et al. 1988; wild et al. 2005; Wild et al. 2010; Hadaidi et al. 2019; Campana et al. 2022.
2.	Fucose	<i>Acropora</i> sp., <i>Fungia</i> sp., <i>Pachyseris speciosa</i> , <i>Fungia fungites</i> , <i>Sarcophyton</i> sp., <i>Lemnalina</i> sp., <i>Cespitularia</i> sp., <i>Galaxea fascicularis</i> , <i>Porites lobata</i> , <i>Ctenactis</i> sp., <i>Pocillopora</i> sp., <i>Pachyseris</i> sp., <i>Lophelia</i> sp.	Meikle et al. 1988; wild et al. 2005; Wild et al. 2010; Sneed et al. 2015; Lee et al. 2016; Hadaidi et al. 2019.
3.	Galactose	<i>Acropora</i> sp., <i>Fungia</i> sp., <i>Acropora formosa</i> , <i>Pachyseris speciose</i> , <i>Fungia fungites</i> , <i>Sarcophyton</i> sp., <i>Lemnalina</i> sp., <i>Cespitularia</i> sp., <i>Porites lobata</i> , <i>Galaxea fascicularis</i> , <i>Stylophora pistillata</i> , <i>Pocillopora verrucose</i> , <i>Ctenactis</i> sp., <i>Pachyseris</i> sp., <i>Lophelia</i> sp.	Meikle et al. 1988; Wild et al. 2005; Wild et al. 2010; Lee et al. 2016; Hadaidi et al. 2019.
4.	Glucose	<i>Acropora</i> sp., <i>Fungia</i> sp., <i>Acropora formosa</i> , <i>Fungia fungites</i> , <i>Sarcophyton</i> sp., <i>Lemnalina</i> sp., <i>Cespitularia</i> sp., <i>P. lobata</i> , <i>G. fascicularis</i> , <i>S. pistillata</i> , <i>P. verrucose</i> , <i>Ctenactis</i> sp., <i>Pocillopora</i> sp., <i>Stylophora</i> sp., <i>Pachyseris</i> sp., <i>Lophelia</i> sp.	Meikle et al. 1988; Wild et al. 2005; Wild et al. 2010; Lee et al. 2016; Hadaidi et al. 2019.
5.	Mannose	<i>Acropora</i> sp., <i>Fungia</i> sp., <i>Acropora formosa</i> , <i>Pachyseris speciose</i> , <i>Fungia fungites</i> , <i>Sarcophyton</i> sp., <i>Lemnalina</i> sp., <i>Cespitularia</i> sp., <i>P. lobata</i> , <i>G. fascicularis</i> , <i>S. pistillata</i> , <i>P. verrucose</i> , <i>Ctenactis</i> sp., <i>28ungi asp.</i> , <i>Pocillopora</i> s p., <i>Pachyseris</i> sp., <i>Pachyseris</i> sp., <i>Lophelia</i> sp.	Meikle et al. 1988; Wild et al. 2005; Wild et al. 2010; Lee et al. 2016; Hadaidi et al. 2019.
6.	N-acetyl galactosamine	<i>Acropora formosa</i> , <i>Lemnalina</i> sp., <i>Pachyseris speciose</i> , <i>Fungia fungites</i> , <i>Sarcophyton</i> sp., <i>Cespitularia</i> sp.	Meikle et al. 1988; Corfield 2015; Shane et al. 2021.
7.	N-acetyl glucosamine	<i>Acropora</i> sp., <i>Acropora Formosa</i> , <i>Pachyseris speciose</i> , <i>Fungia fungites</i> , <i>Cespitularia</i> sp. <i>Sarcophyton</i> sp., <i>Lemnalina</i> sp., <i>A. pharaonis</i> , <i>P. lobata</i> , <i>G. fascicularis</i> , <i>S. pistillata</i> , <i>Ctenactis</i> sp., <i>Fungia</i> sp., <i>Pachyseris</i> sp., <i>Lophelia</i> sp.	Meikle et al. 1988; Salvatore et al. 2000; Wild et al. 2005; Wild et al. 2010; Hadaidi et al. 2019.
8.	Ribose	<i>A. pharaonis</i> , <i>P. lobata</i>	Wild et al. 2010.
9.	Rhamnose	<i>A. pharaonis</i> , <i>P. lobata</i> , <i>Acropora</i> sp., <i>Madrepora</i> sp., <i>Lophelia</i> sp.	Wild et al. 2010; Hadaidi et al. 2019.
10	Xylose	<i>Fungia fungites</i> , <i>Sarcophyton</i> sp., <i>Lemnalina</i> sp., <i>Acropora</i> sp., <i>Fungia</i> sp., <i>Lophelia</i> sp.	Meikle et al. 1988; Wild et al. 2010; Krediet et al. 2013; Hadaidi et al. 2019.

**Table 2 C/N ratio in fresh coral mucus.**

<b>Hard corals:</b>	<b>C/N ratio</b>	<b>Reference</b>
<i>Acropora</i> sp.	8.5	(Coles and Strathmann 1973)
<i>Acropora formosagrands</i>	5.9	(Vacelet and Thomassin 1991)
<i>Fungia scutaria</i>	7.5	(Coles and Strathmann 1973)
<i>Fungia scutaria</i>	6.9	(Krupp 1982).
<i>Fungia echinata</i>	5.6	(Vacelet and Thomassin 1991)
<i>Porites compressa</i>	6.0	(Coles and Strathmann 1973)
<i>Goniopora lobata</i>	5.9	(Coles and Strathmann 1973)
<i>Platygyra lamellina</i>	6.6	(Pascal 1981)
<i>P. astreoides</i>	5.9	(Coffroth 1990)
<i>P. furcate</i>	4.1	(Coffroth 1990)
<i>P. australiensis</i>	5.9	(Coffroth 1990)
<i>P. lutea</i>	5.9	(Coffroth 1990)
<i>P. lobata</i>	5	(Coffroth 1990)
<i>P. murrayensis</i>	5.5	(Coffroth 1990)
<i>Porites</i> sp.	4.8	(Coffroth 1990)
<b>Soft corals:</b>		
<i>Heteroxenia fuscescens</i>	7.3	(Pascal 1981)
<i>Sarcophyton</i> cf. <i>glaucum</i>	5.6	(Vacelet and Thomassin 1991)
<i>Briarium asbestinum</i>	8.9	(Pascal 1981)

Figure 1




Figure 2




Figure 3

